

The Arabidopsis *CLAVATA2* Gene Encodes a Receptor-like Protein Required for the Stability of the *CLAVATA1* Receptor-like Kinase

Sangho Jeong, Amy E. Trotochaud, and Steven E. Clark¹

Department of Biology, University of Michigan, Ann Arbor, Michigan 48109-1048

The *CLAVATA2* (*CLV2*) gene regulates both meristem and organ development in Arabidopsis. We isolated the *CLV2* gene and found that it encodes a receptor-like protein (RLP), with a presumed extracellular domain composed of leucine-rich repeats similar to those found in plant and animal receptors, but with a very short predicted cytoplasmic tail. RLPs lacking cytoplasmic signaling domains have not been previously shown to regulate development in plants. Our prior work has demonstrated that the *CLV1* receptor-like kinase (RLK) is present as a disulfide-linked multimer in vivo. We report that *CLV2* is required for the normal accumulation of *CLV1* protein and its assembly into protein complexes, indicating that *CLV2* may form a heterodimer with *CLV1* to transduce extracellular signals. Sequence analysis suggests that the charged residue in the predicted transmembrane domain of *CLV2* may be a common feature of plant RLPs and RLKs. In addition, the chromosomal region in which *CLV2* is located contains an extremely high rate of polymorphism, with 50 nucleotide and 15 amino acid differences between Landsberg *erecta* and Columbia ecotypes within the *CLV2* coding sequence.

INTRODUCTION

The shoot meristem is responsible for the formation of all above-ground organs in higher plants. This task is accomplished by balancing two processes: the maintenance of a population of undifferentiated cells at the center of the shoot meristem and the direction of progeny cells on the flanks of the meristem toward organ formation and eventual differentiation. Mutations in several genes in Arabidopsis have been shown to disrupt this balance (reviewed in Clark, 1997), including *SHOOT MERISTEMLESS* (*STM*), *WUSCHEL* (*WUS*), and the genes comprising the *CLAVATA* loci (*CLV1*, *CLV2*, and *CLV3*) (Leyser and Furner, 1992; Barton and Poethig, 1993; Clark et al., 1993, 1995; Laux et al., 1996; Kayes and Clark, 1998). Plants carrying mutations at the *CLV* loci accumulate undifferentiated cells in the shoot and flower meristems, and it has therefore been proposed that the *CLV* genes either inhibit the division of these cells or promote their transition toward a differentiated state (reviewed in Clark, 1997).

The phenotypes of *clv1* and *clv3* mutants are identical, and these two genes function in the same pathway to regulate meristem development. In addition, the phenotypes of the *clv1* and *clv3* mutants and the gene expression patterns for

CLV1 and *CLV3* appear to be specific to the shoot and flower meristems (Clark et al., 1993, 1995, 1997; Fletcher et al., 1999). Compared with *clv1* and *clv3* mutants, all *clv2* mutant alleles display weak phenotypes at the shoot and flower meristems (Kayes and Clark, 1998). *clv2* mutations also affect the development of several organ types. These include the pedicel, which is elongated in *clv2* mutants, and the anthers and valves of the flower, both of which are reduced in *clv2* mutants. Genetic analysis indicates that *CLV2* functions in the same pathway as *CLV1* and *CLV3* in the regulation of meristem development but in a separate pathway in the regulation of organ development (Kayes and Clark, 1998).

CLV1 and *CLV3* appear to encode components of a signal transduction pathway. The predicted *CLV1* protein is a receptor-like kinase (RLK), with extracellular leucine-rich repeats (LRRs), a single transmembrane domain, and a functional intracellular serine protein kinase domain (Clark et al., 1997; Williams et al., 1997; Stone et al., 1998). *CLV3* encodes a potential ligand in the form of a small, secreted protein (Fletcher et al., 1999). Evidence for the role of *CLV3* in the activation of *CLV1* has come from analysis of *CLV1* protein complexes in vivo (Trotochaud et al., 1999). *CLV1* is found in two protein complexes in cauliflower and Arabidopsis. One complex is an inactive disulfide-linked multimer of 185 kD, and the other, which is presumed to be the active complex, is a 450-kD complex that includes the 185-kD

¹To whom correspondence should be addressed. E-mail clarks@umich.edu; fax 734-647-0884.

CLV1 multimer as well as additional proteins. Inferences as to the function of these two complexes (active and inactive) are based on analysis of CLV1 complex formation in mutant plants. In a *clv1* mutant with a partial loss of in vitro kinase activity (*clv1-7*), a reduction of the proportion of CLV1 in the 450-kD complex was observed, whereas in a *clv1* mutant with a complete loss of in vitro kinase activity (*clv1-10*), a complete loss of the 450-kD complex was observed. In *clv3* mutants, a complete loss of the 450-kD complex was also observed, indicating that CLV3 is required for CLV1 activation (Trotochaud et al., 1999).

Additional components of the 450-kD complex have been identified (Trotochaud et al., 1999). One component is the kinase-associated protein phosphatase (KAPP), which was shown previously to interact with several protein kinases (Stone et al., 1994; Braun et al., 1997) and to be a negative regulator of CLV1 signal transduction (Williams et al., 1997; Stone et al., 1998). The 450-kD complex also contains a Rho GTPase-related protein that presumably relays signals to downstream components (Trotochaud et al., 1999).

Because *CLV2* functions in the same pathway as *CLV1* and *CLV3* to regulate meristem development, we considered it a reasonable hypothesis that *CLV2* may code for an additional component of one or both of the CLV1 complexes. However, the *CLV2* protein must be unique in function because inactivating *CLV2* results in a weaker phenotype than does inactivating *CLV1* or *CLV3* and because *CLV2* regulates the development of several different organ types independently of *CLV1* and *CLV3* (Kayes and Clark, 1998). Thus, we undertook to isolate the *CLV2* gene. We report here that it encodes a receptor-like protein (RLP) with predicted extracellular LRRs but only a very short cytoplasmic domain. Consistent with a role in CLV1 protein function, we demonstrate that *CLV2* is required for CLV1 protein accumulation. Possible roles for *CLV2* are as a heterodimer partner for *CLV1* to detect extracellular signals within the shoot meristem and as a heterodimer partner with other RLKs to regulate the development of organs such as the pedicel and gynoecium.

RESULTS

CLV2 Cloning

The *clv2-5* mutant allele was isolated from T-DNA insertional mutagenized populations (Feldmann, 1991; Kayes and Clark, 1998). By outcrossing to wild-type *Arabidopsis*, we observed tight linkage between the *clv2* phenotype and kanamycin resistance (data not shown). The *clv2-5* genomic region surrounding the T-DNA insertion was cloned and sequenced. A complex, highly rearranged T-DNA insertion was found to have disrupted the open reading frame (ORF) of a gene encoding an RLP (Figure 1). Sequences for this




Figure 1. *CLV2* Encodes an RLP.

(A) The genomic organization at the *CLV2* locus, indicating the site of T-DNA insertion in the *clv2-5* allele. The original T-DNA structure is shown at the top, with its domains color coded. The best estimate of the T-DNA structure after insertion is shown in the middle. The orientations of the colored arrows represent the orientations of the corresponding domains after insertion. Restriction sites used for sequencing are indicated (E, EcoRI; H, HindIII; S, SacI). The region from the wild-type *Col* ecotype genome used to complement *clv2* is also shown. LB, left border; Amp^r, ampicillin resistance; Tn₉₀₃, transposon 903; Kan^r, kanamycin resistance; Tet^r, tetracycline resistance; RB, right border.

(B) to (J) Regions of the predicted *CLV2* protein, separated into the signal peptide (B), the cysteine-pair regions (C) and (G), the LRRs (D) and (F), the "island" (E), the acidic domain (H), the transmembrane domain (I), and the cytoplasmic stop transfer sequence (J). Red amino acids indicate residues that are polymorphic among different ecotypes. Dashes indicate gaps introduced into the sequence to maximize alignment. The GenBank accession numbers for the *CLV2* gene and protein sequence from *Col*, *Ler*, and *Ws* are AF177672, AF177673, and AF177674, respectively.

Table 1. *clv2* Alleles

Allele	Lesion ^a	Predicted Effect
<i>clv2-1</i>	C→T at 97	Gln-33→stop codon
<i>clv2-2</i>	1-bp deletion at 1281	Frameshift
<i>clv2-3</i>	61-bp deletion from 1095 to 1155	Frameshift
<i>clv2-4</i>	C→T at 97	Gln-33→stop codon
<i>clv2-5</i>	T-DNA insertion at 2158	Frameshift

^aNucleotides are numbered from the predicted translation start site.

ORF were compared among all five *clv2* alleles and their corresponding wild-type ecotypes. This revealed lesions leading to premature termination codons in all mutant alleles, suggesting that this ORF corresponded to the *CLV2* gene (Table 1) and that the *clv2* mutations are likely null alleles. To confirm that this ORF was *CLV2*, we transformed a 4.9-kb fragment from the wild-type genomic region into *clv2-3* mutants (Figure 1A). Of 47 primary transformants, 44 exhibited a complete suppression of the mutant phenotype (data not shown).

Predicted CLV2 Protein Structure

The *CLV2* gene appears to contain a single exon that codes for a 720-amino acid protein with a characteristic signal peptide (Figures 1B to 1J). 5' Rapid amplification of cDNA ends (RACE) analysis indicated a single major transcription initiation site 153 bp upstream of the predicted start ATG (see Methods). An expressed sequence tag (GenBank accession numbers AA586072 and T41629) for this gene contains the predicted stop codon, 3' untranslated region, and polyadenylation tail, indicating that no unidentified exons exist downstream.


The majority of the predicted *CLV2* protein consists of LRRs. The alignment for these repeats is shown in Figures 1D and 1F. The LRRs are flanked by cysteine pairs that are each separated by seven residues, a motif common to extracellular LRR proteins (Jones and Jones, 1996) (Figures 1C and 1G). The *CLV2* protein shares similarity with a growing family of receptors and RLKs found in both plants and animals (Jiang et al., 1995; Lease et al., 1998). However, the *CLV2* protein lacks a kinase domain. Receptor-like LRR proteins without any cytoplasmic signaling domain are relatively rare, and only one family has been previously described in plants. This is the *Cf* family of genes from tomato, which provides resistance to specific races of the fungus *Cladosporium fulvum* (Jones et al., 1994; Dixon et al., 1996). Both the *CLV2* and the *Cf* RLPs have a very short predicted cytoplasmic domain and an island of non-LRR sequence between repeats 4 and 5, counting from the transmembrane

domain (Figures 1E and 1J). *CLV2* and *Cf* proteins also share an acidic domain immediately adjacent to the transmembrane domain (Figure 1H).

CLV2 Is Expressed in Many Tissues

Because *CLV2* is required for the proper development of many organ types, we predicted that *CLV2* should be expressed in many different tissues. Total RNA was collected from apices (shoot meristems and unopened flower meristems), opened flowers, siliques, 4-day-old seedlings, and rosette leaves and used for RNA gel blot analysis. Whereas the highest expression was detected in apices, as expected, *CLV2* mRNA was detected in all tissues (Figure 2A).

To determine whether the weaker signal detected in some of the organ types was the result of hybridization to a related sequence, we tested the Arabidopsis genome for cross-hybridizing sequences. Low-stringency DNA gel blot hybridization using the entire *CLV2* gene as a probe detected only the *CLV2* genomic sequence (Figure 2B), indicating that no closely related genes exist in Arabidopsis. Moreover, no highly similar DNA sequences were identified by BLAST analysis (Altschul et al., 1997).

**Figure 2.** *CLV2* Is Expressed in All Tissues Tested.

(A) Twenty micrograms of total RNA isolated from wild-type Arabidopsis 4-day-old seedlings (lane 1), rosette leaves (lane 2), siliques (lane 3), opened flowers (lane 4), and apices (shoot meristems and unopened flowers; lane 5) was used to prepare an RNA gel blot that was probed consecutively with a portion of the *CLV2* coding region and an 18S rDNA control.

(B) Genomic DNA from wild-type Col (lane 1), *Ler* (lane 2), and *Ws* (lane 3) ecotypes was digested with *EcoRI* and used to generate a DNA gel blot probed with the entire *CLV2* coding region. This was washed at low stringency. Note that the *Ws* genome has a polymorphic *EcoRI* site at position 433 in the coding region, resulting in two hybridizing bands.

CLV2 Is Required for CLV1 Protein Accumulation

Because *CLV2* functions in the same pathway as *CLV1* to regulate meristem development (Kayes and Clark, 1998), we hypothesized that the *CLV2* gene product could interact with the *CLV1* RLK. We have previously demonstrated that *CLV1* is found in two distinct protein complexes in vivo: a disulfide-linked 185-kD complex and a larger 450-kD complex that includes the 185-kD multimer, the protein phosphatase KAPP, and a Rho GTPase-related protein (Trotochaud et al., 1999). We have proposed that the 185-kD complex is the inactive form of *CLV1* and may represent a *CLV1* hetero-




Figure 3. *CLV2* Is Required for *CLV1* Protein Stability.

(A) Total extracts from wild-type *Ler* and *clv2-3* plants were separated by SDS-PAGE, and the presence of *CLV1* was detected by protein gel blot analysis.

(B) Total extracts from wild-type *Ler*, *clv2-2*, and *clv2-3* plants were tested using ELISA assays for the amount of *CLV1* (open bars) and KAPP protein (solid bars) present. Standard error is indicated.

(C) Total extracts from wild-type *Ler* (solid circles) and *clv2-3* (open squares) plants were separated by gel chromatography and tested using ELISA assays for the amount of *CLV1* protein present in each fraction.

dimer with a second RLK. That the 450-kD *CLV1* complex is absent in *clv1* and *clv3* mutant backgrounds and is associated with a Rho-related signaling protein suggest that the 450-kD complex contains the active form of *CLV1*. If *CLV2* is a component of either complex, then eliminating the *CLV2* protein should alter complex size and/or accumulation.

To test this hypothesis, we first examined *CLV1* protein levels in wild-type and *clv2-3* plants by protein gel blotting, as shown in Figure 3A. We did not detect cross-reaction of the *CLV1* antiserum to extracts from *clv2-3* plants, suggesting a significant reduction in *CLV1* protein in these plants. To quantify this difference precisely, we performed ELISA assays on extracts from wild-type, *clv2-2*, and *clv2-3* plants, revealing a >90% reduction in *CLV1* accumulation in *clv2* plants (Figure 3B). In a control procedure, the same extracts from *clv2* plants were shown to have a normal amount of KAPP protein (Figure 3B). Because we have previously shown that *clv2* mutants have normal, if not higher, levels of *CLV1* transcription (Kayes and Clark, 1998), any changes in *CLV1* protein accumulation must be due to post-transcriptional mechanisms. ELISA analysis of gel chromatography fractions revealed that the *CLV1* protein that is present in *clv2* mutants is primarily found in a novel, higher molecular mass complex of ~600 kD (Figure 3C).

Unprecedented Polymorphisms

While determining the sequences of the *clv2* mutant alleles, we observed a very high level of sequence polymorphism between various *Arabidopsis* ecotypes. We observed 50 nucleotide polymorphisms between Landsberg *erecta* (*Ler*) and Columbia (*Col*) and 54 between *Ler* and Wassilewskija (*Ws*) (Table 2). In total, 68 sites were found to be polymorphic. These polymorphisms have a significant affect on the predicted amino acid sequences of the *CLV2* protein from the various ecotypes. Red residues in Figure 1 indicate those that are polymorphic between the ecotypes. Most are clustered in a short region of sequence in the N-terminal portion of the predicted protein. The changes predominantly alter those residues that are not part of the LRR consensus. The three consensus residues that are polymorphic are subjected to conservative substitutions: residue 484 is valine (*Ler* and *Col*) and isoleucine (*Ws*); residue 532 is phenylalanine (*Col*), leucine (*Ler*), and isoleucine (*Ws*); and residue 739 is valine (*Ler*) and isoleucine (*Col* and *Ws*).

To determine whether the polymorphisms might reflect selection for modified forms of *CLV2*, we examined the amount of sequence polymorphism in the coding sequences of the adjacent serpin ORF and in the intergenic region between the *CLV2* and serpin ORFs. These regions also contained a very high degree of sequence polymorphism. The serpin ORF contained 20 sites that were polymorphic between the *Ler*, *Col*, and *Ws* ecotypes over the 1068 bp of the predicted coding sequences. These polymorphisms included two deletions/insertions that are predicted to disrupt

Table 2. Sequence Polymorphisms in the *CLV2* Gene^a

Nt ^b	Col	Ler	Ws	Nt	Col	Ler	Ws
36	T	C	C	693	G	A	G
84	C	T	T	696	A	G	A
138	C	T	T	723	T	A	T
196	C	T	T	730	A	G	A
231	C	T	A	739	A	C	A
239	T	T	C	813	A	G	A
243	C	T	T	816	T	C	T
259	A	G	G	819	C	T	C
324	C	T	T	853	A	G	A
329	C	G	C	1095	C	C	T
348	C	C	T	1110	T	T	A
372	G	G	A	1119	C	C	T
374	C	C	T	1120	G	T	G
375	A	A	T	1128	C	C	G
390	G	A	G	1131	C	C	A
400	G	A	G	1176	G	A	G
411	A	T	T	1182	A	T	T
414	C	T	C	1239	A	G	A
415	G	C	G	1335	C	T	C
423	T	C	T	1344	G	A	G
424	G	A	G	1347	G	A	G
426	A	G	G	1392	T	C	C
442	A	G	A	1398	T	C	T
465	T	C	T	1410	G	A	G
469	G	A	G	1413	T	C	T
477	C	C	T	1452	A	G	A
483	G	A	G	1500	A	A	G
484	G	G	A	1563	T	T	A
486	C	T	T	1605	T	G	G
502	A	G	G	1621	T	T	C
532	T	C	A	1626	T	T	C
534	T	T	A	1715	A	G	A
680	A	G	A	1854	T	C	T
687	C	G	C	1945	G	G	A

^aThe *CLV2* gene (2163 bp) was sequenced from three Arabidopsis ecotypes, namely, Col, Ler, and Ws. Sixty-eight sites were found to be polymorphic among the three ecotypes.

^bNucleotide (Nt) positions are numbered starting from the translational start.

the reading frame in the Ler and Ws ecotypes. The introns of the serpin gene contain 17 polymorphisms over 305 bp of sequences, including an alteration of a base at a splice acceptor site. The 554-bp intergenic region between the reading frames for CLV2 and the serpin ORF contain nine polymorphisms, including a 251-bp deletion in the Ws ecotype compared with Ler and Col.

DISCUSSION

The identification of the *CLV2* gene revealed a predicted RLP that represents a novel class of developmental regula-

tors. CLV2 is the first RLP lacking a cytoplasmic signaling domain that has been shown to regulate development in plants. Previously, the only known functional RLPs were those encoded by the *Cf* gene family from tomato, which is required for recognition of specific races of the fungus *C. fulvum* (Jones et al., 1994; Dixon et al., 1996).

CLV2 Function

How does the identification of the *CLV2* gene sequence help us to understand its role in development? *CLV2* operates in the same pathway as *CLV1* and *CLV3*, whose gene products appear to function as components of a signal transduction cascade. *CLV1* encodes an RLK (Clark et al., 1997), whereas *CLV3* encodes a small, secreted protein (Fletcher et al., 1999). *CLV3* is also required for the activation of the CLV1 protein in vivo (Trotochaud et al., 1999). Because *CLV2* functions in the same pathway as *CLV1* and *CLV3*, we would expect CLV2 to function as a component of this signal transduction cascade. Based on the predicted CLV2 protein structure, one possible role for CLV2 in CLV1/CLV3 signaling could be to act as a heterodimer partner with CLV1. This would be consistent with the size of the inactive complex, which is ~185 kD. However, size determination by column chromatography can be influenced by several variables, and the actual mass of the components of this complex may add up to significantly more or less than 185 kD. Thus, although the size of the inactive CLV1 complex is consistent with a CLV1/CLV2 heterodimer, it does not rule out other possibilities, such as a CLV1 homodimer.

Additional evidence for CLV1/CLV2 heterodimers came from analysis of CLV1 protein amounts in the *clv2* mutant background. We observed that >90% of the CLV1 protein was absent in these plants, despite the fact that we have previously shown that *CLV1* is transcribed at normal, if not higher, levels in *clv2* plants (Kayes and Clark, 1998). Thus, the loss of CLV1 protein in *clv2* mutants reflects post-transcriptional regulation. In many other systems, when one member of a heterodimer pair is absent, the partner protein is degraded in the endoplasmic reticulum (Hurtley and Helenius, 1989). The *clv2* mutation affected the accumulation of both the 185- and 450-kD CLV1 complexes, which suggests that CLV2 is a component of both complexes. Interestingly, the CLV1 protein that is present in *clv2* mutants is primarily found in a novel, higher molecular weight complex of ~600 kD (Figure 3B). If this remaining CLV1 protein is active, then it would explain why all *clv2* mutant alleles are weak in phenotype compared with *clv1* and *clv3* mutants. The recent cloning of *CLV3*, revealing a predicted small, secreted protein, is consistent with a role for CLV3 as the ligand for CLV1 (Fletcher et al., 1999). If this is the case, then efficient ligand binding may require both CLV1 and CLV2. However, the evidence for CLV1/CLV2 heterodimer formation is circumstantial. Definitive proof will require an analysis of coimmunoprecipitation between CLV1 and CLV2.

If CLV1 forms a heterodimer with CLV2, then binding would likely be mediated by disulfide linkage between conserved cysteine pairs flanking the LRR regions. We have previously shown that all CLV1 protein in vivo is in a disulfide-linked multimer of 185 kD (Trotochaud et al., 1999). Charge interaction in the extracellular sequences immediately adjacent to the putative transmembrane domain could also play a role in heterodimer formation. In this region, CLV1 is basic, whereas CLV2 is acidic. This type of charge interaction in an extracytoplasmic peritranmembrane (EPTM) domain was suggested as a mechanism for major histocompatibility complex II $\alpha\beta$ heterodimer formation (Cosson and Bonifacino, 1992). Because CLV2 functions in a pathway separate from CLV1 in the regulation of organ development, one possibility is that CLV2 forms a heterodimer with a separate RLK in these tissues.

How LRR-containing RLPs that lack a cytoplasmic domain can transmit signals across the plasma membrane is unknown. Heterodimer formation between CLV1 and CLV2 provides a possible mechanism by which this class of RLPs may function as signaling components. That is, these RLPs may form heterodimers with an RLK to form a functional complex with both a receptor domain and a cytoplasmic signaling domain. Interestingly, Xa21, a rice RLK that provides resistance to a specific race of *Xanthomonas* spp, was shown to function without a transmembrane or cytoplasmic domain (Wang et al., 1998). Perhaps the truncated form of Xa21 functions in a manner analogous to CLV2.

Transmembrane Charge

An interesting aspect of the deduced CLV2 amino acid sequence is the presence of an aspartic acid residue in the

predicted transmembrane domain. There are only a few examples of single-pass transmembrane RLPs containing a charged residue in the transmembrane domain (Harrison, 1996). The best-studied example is the T-cell receptor (TCR) complex. The components of the TCR complex contain charged residues in their transmembrane domains that have been shown to be critical for protein-protein interactions and complex formation (Cosson et al., 1991).

To address the potential significance of the charged residues in the transmembrane and EPTM domains in plant RLP and RLK function, we analyzed 54 additional RLKs and RLPs identified from the Arabidopsis genome database. Of the sequences analyzed, 43 encoded apparent RLKs, whereas 12 (including CLV2) encoded RLPs (Table 3). Six of the RLKs and seven of the RLPs contained charged residues in their predicted transmembrane domain. These were evenly split between basic and acidic residues, with most of the RLKs containing acidic residues and most of the RLPs containing basic residues. Thus, charged transmembrane domain residues appear to be a common feature of plant RLPs and RLKs. Our analysis also indicated that all RLKs have basic EPTM domains and that all RLPs have acidic EPTM domains (Table 3). The opposite charges of the EPTM residues may mediate heterodimer formation between RLPs and RLKs. Although CLV1 lacks a corresponding charged residue in its predicted transmembrane domain, the charged amino acid in CLV2 could mediate interaction with an RLK in its regulation of organ development or could mediate interaction with other components of the CLV1 signaling complex.

CLV2 Polymorphisms

The region of chromosome 1 in which CLV2 is located has the highest level of polymorphism reported to date for single-copy gene regions of Arabidopsis. The only other reports of highly polymorphic regions in Arabidopsis are the gene clusters of disease resistance loci (Botella et al., 1998; McDowell et al., 1998). The 68 polymorphic sites over 2163 bp represented at least an order of magnitude increase over that typically found among the Col, Ler, and Ws ecotypes for other genes. For example, the CLV1 gene, also located on chromosome 1, has no polymorphic sites between the Ler and Col ecotype over 2940 bp of coding sequence. The ratio of synonymous to nonsynonymous changes (2.4:1) and the proportion of third-position to first- and second-position changes (2:1) found between Ler and Ws suggest that CLV2 remains under selection (Table 4). Indeed, despite the amino acid polymorphisms, the CLV2 alleles found in each of the three ecotypes are functionally similar, based on the similarity of their phenotypes when mutated (Kayes and Clark, 1998).

This high amount of polymorphism likely does not reflect selection for altered versions of CLV2. This conclusion is based on the observation that the adjacent gene and the

Table 3. Charged Residues in Predicted Transmembrane Domains of RLPs and RLKs

Type ^a	Total ^b	No Charge ^b	His ^b	Asp ^b	Lys ^b	EPTM ^c	
						Basic	Acidic
RLK	43	37	0	5	1	43	0
RLP	12	5	6	1	0	0	12

^a The predicted CLV2 amino acid sequence was used in a BLAST analysis (Altschul et al., 1997) to identify fully sequenced homologous genes (partial clones and expressed sequence tags were not included).

^b The number of predicted gene products in each category containing no charged residues (No charge), histidine (His), aspartic acid (Asp), and lysine (Lys) in their putative transmembrane domains.

^c The first 10 residues from the predicted transmembrane domain were assessed to determine if the putative EPTMs were basic or acidic overall.

Table 4. Summary of Polymorphisms in the *CLV2* Gene

Ecotypes	Total ^a	Ts ^b	Tv ^c	1st ^d	2nd	3rd	Syn ^e	Nsyn ^f
Col vs. <i>Ler</i>	50	41	9	13	3	34	35	15
Col vs. <i>Ws</i>	34	23	11	7	2	25	27	7
<i>Ler</i> vs. <i>Ws</i>	54	40	14	13	5	36	38	16

^a Total number of polymorphic sites.

^b Transitions.

^c Transversions.

^d Polymorphisms at the first, second, and third codon positions.

^e Synonymous changes.

^f Nonsynonymous changes.

intergenic region contain similar levels of polymorphism. Thus, the changes in the *CLV2* sequence indicate that this region of chromosome 1 is subjected to a very high level of mutagenesis. Why this region is mutagenic is unclear. It is unlikely due to high levels of methylation, because the ratio of transitions to transversions (2.9:1 between *Ler* and *Ws*) is not unusually high. Other genes that have been shown to have high levels of polymorphism in *Arabidopsis* (although not as high as *CLV2*) include *AP3* (*APETALA3*; Purugganan and Suddith, 1999), *RPS2* (*RESISTANCE TO PSEUDOMONAS SYRINGAE2*; Caicedo et al., 1999), and *CAL* (*CAULIFLOWER*; Purugganan and Suddith, 1998). Interestingly, the highly rearranged T-DNA insertion in the *clv2-5* line (Figure 1A) may reflect the genomic instability in this region.

If the high level of polymorphism is consistent in other ecotypes, then sequencing of *CLV2* in additional ecotypes may provide a wealth of structure-function information. The polymorphisms detected in the *Ler*, *Col*, and *Ws* ecotypes suggest that LRR repeats 1 to 4 may not be functionally as important as the remaining portion of the protein, because the majority of first- and second-position substitutions occur in this region.

METHODS

CLV2 Cloning

A λ genomic library was generated using DNA isolated from *clv2-5* homozygous *Arabidopsis thaliana* plants and was screened by using the neomycin phosphotransferase *nptII* sequence from the T-DNA as a probe. Five hybridizing λ clones were isolated, subcloned, and partially sequenced. Plant DNA on each side of the T-DNA insertion was used to probe a λ genomic library made from wild-type Columbia (*Col*). Two λ clones hybridizing to both flanking probes were isolated and analyzed. The single expressed sequence tag for the *CLAVATA2* gene (GenBank accession numbers AA586072 and T41629) was obtained from the *Arabidopsis* Biological Resource Center (Ohio State University, Columbus, OH).

Sequence Analysis

For all sequencing, both DNA strands were sequenced. For *clv2* allele sequencing, genomic DNA was isolated from each mutant, and the entire coding region was amplified by polymerase chain reaction (PCR). The PCR products were gel-purified and used as templates for direct PCR sequencing with a genetic analyzer (model ABI310; Perkin-Elmer, Norwalk, CT). For analysis of polymorphisms, the *CLV2* gene was amplified by PCR from Landsberg *erecta* (*Ler*) and Wassilewskija (*Ws*) genomic DNA and directly sequenced. The corresponding *Col* genomic region was not sequenced because it was available from the database and it matched the sequence from the *clv2-3* allele, which was isolated from the *Col* ecotype. The intergenic region and the serpin gene from *Ler* and *Ws* were sequenced the same way as for *CLV2*. The *Col* serpin sequence from the database was partially verified by sequencing the first exon to confirm a deletion/insertion site that disrupts the open reading frame in the *Ler* and *Ws* ecotypes.

DNA Gel Blot Analysis

The entire *CLV2* gene was used as a probe. Hybridization was conducted overnight at 65°C in 0.25 M sodium phosphate, pH 7.4, 1 mM EDTA, 1% casein, and 7% SDS. The filter was washed twice with 2 × SSPE (1 × SSPE is 0.18 M NaCl, 10 mM NaH₂PO₄, and 10 mM EDTA, pH 7.4) and 0.1% SDS at room temperature for 10 min each and then with 0.5 × SSPE and 0.1% SDS at 50°C for 20 min.

Rapid Amplification of cDNA Ends

For 5' rapid amplification of cDNA ends (RACE), total RNA was isolated from shoot apices using the RNeasy plant mini kit (Qiagen, Chatsworth, CA), and mRNA was purified using the PolyATract mRNA isolation system (Promega). 5' RACE PCR was performed with a 5' RACE kit (Boehringer Mannheim) according to the manufacturer's instructions. Primers used for reverse transcription and PCR reactions were as follows: primer 1, 5'-GCTAAACCTGTCCAGTTA-GAA-3'; primer 2, 5'-TCGAGATCAGGAAGCTGAGAT-3'; and primer 3, 5'-TCAACGGAGAAAGATGAAGTT-3'. A major PCR product was gel-purified and sequenced by direct PCR sequencing with an ABI310 genetic analyzer.

RNA Gel Blot Analysis

Plant tissues were collected from shoot apices with unopened flowers, opened flowers, rosette leaves, green siliques, and 4-day-old seedlings and were immediately frozen in liquid nitrogen. Total RNAs were extracted from each tissue using the RNeasy plant mini kit. Twenty micrograms of total RNA was used for blotting. The NorthernMax kit (Ambion, Austin, TX) was used for RNA gel blot hybridization, according to the manufacturer's instructions. A 945-bp fragment was amplified using primers 5'-TGGATCCCAGATTCA-CCCATCTCT-3' and 5'-AGAATTCTCATTCGCCAGAGAGATTGTT-3' from the *CLV2* coding region and cloned into the pCR2.1 vector (Stratagene, La Jolla, CA) to generate an RNA probe with the ribo-probe in vitro transcription system (Promega).

Plant Growth Conditions

Seeds were sown on a 1:1:1 mix of top soil:perlite:vermiculite and imbibed for 5 days at 4°C. Plants were grown at 20°C under 900 foot-candles of constant cool white fluorescent light and watered with 0.05 g/L 20-20-20 fertilizer (Knox Fertilizer Co., Knox, IN). Seedlings used for RNA analysis were grown on plates containing Murashige and Skoog salts (Murashige and Skoog, 1962), pH 5.7, 0.8 % agar, and 1% sucrose.

Plant Transformation

The 4.9-kb genomic fragment was subcloned into binary vector pPZP 222 (Hajdukiewicz et al., 1994) and transformed into *Agrobacterium tumefaciens* GV3101. Wild-type *Ler* and *clv2-3* plants were transformed by the vacuum infiltration method (Bechtold et al., 1993). Transformants were selected on plates with gentamycin, and segregation of *clv2-3* plants in the next generation was observed.

Protein Analysis

All protein work was performed as described by Trotochaud et al. (1999).

ACKNOWLEDGMENTS

We thank David Bay for photographic assistance, Robert Williams for identifying the *clv2-5* allele, and Jeffrey Kayes for performing linkage analysis. This work was supported by the U.S. Department of Energy (Grant No. DE-FG02-99ER14972).

Received July 1, 1999; accepted August 4, 1999.

REFERENCES

- Altschul, S.F., Madden, T.L., Schäffer, A.A., Zhang, J., Zhang, Z., Miller, W., and Lipman, D. (1997). Gapped BLAST and PSI-BLAST: A new generation of protein database search programs. *Nucleic Acids Res.* **25**, 3389–3402.
- Barton, M.K., and Poethig, R.S. (1993). Formation of the shoot apical meristem in *Arabidopsis thaliana*: An analysis of development in the wild type and *shoot meristemless* mutant. *Development* **119**, 823–831.
- Bechtold, N., Ellis, J., and Pelletier, G. (1993). In planta *Agrobacterium*-mediated gene transfer by infiltration of adult *Arabidopsis thaliana* plants. *C. R. Acad. Sci. Paris Sci. Vie* **316**, 1194–1199.
- Botella, M.A., Parker, J.E., Frost, L.N., Bittner-Eddy, P.D., Beynon, J.L., Daniels, M.J., Holub, E.B., and Jones, J.D.G. (1998). Three genes of the *Arabidopsis* *RPP1* complex resistance locus recognize distinct *Peronospora parasitica* avirulence determinants. *Plant Cell* **10**, 1847–1860.
- Braun, D.M., Stone, J.M., and Walker, J.C. (1997). Interaction of the maize and *Arabidopsis* kinase interaction domain with a subset of receptor-like protein kinases: Implications for transmembrane signaling in plants. *Plant J.* **12**, 83–95.
- Caicedo, A.L., Schaal, B.A., and Kunkel, B.N. (1999). Diversity and molecular evolution of the *RPS2* resistance gene in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **98**, 302–306.
- Clark, S.E. (1997). Organ formation at the vegetative shoot meristem. *Plant Cell* **9**, 1067–1076.
- Clark, S.E., Running, M.P., and Meyerowitz, E.M. (1993). *CLAVATA1*, a regulator of meristem and flower development in *Arabidopsis*. *Development* **119**, 397–418.
- Clark, S.E., Running, M.P., and Meyerowitz, E.M. (1995). *CLAVATA3* is a specific regulator of shoot and floral meristem development affecting the same processes as *CLAVATA1*. *Development* **121**, 2057–2067.
- Clark, S.E., Williams, R.W., and Meyerowitz, E.M. (1997). The *CLAVATA1* gene encodes a putative receptor kinase that controls shoot and floral meristem size in *Arabidopsis*. *Cell* **89**, 575–585.
- Cosson, P., and Bonifacino, J.S. (1992). Role of transmembrane domain interactions in the assembly of class II MHC molecules. *Science* **258**, 659–662.
- Cosson, P., Lankford, S.P., Bonifacino, J.S., and Klausner, R.D. (1991). Membrane protein association by potential intramembrane charge pairs. *Nature* **351**, 414–416.
- Dixon, M.S., Jones, D.A., Keddie, J.S., Thomas, C.M., Harrison, K., and Jones, J.D.G. (1996). The tomato *Cf-2* disease resistance locus comprises two functional genes encoding leucine-rich repeat proteins. *Cell* **84**, 451–459.
- Feldmann, K.A. (1991). T-DNA insertion mutagenesis in *Arabidopsis*: Mutational spectrum. *Plant J.* **1**, 71–82.
- Fletcher, J.C., Brand, U., Running, M.P., Simon, R., and Meyerowitz, E.M. (1999). Signaling of cell fate decisions by *CLAVATA3* in *Arabidopsis* shoot meristems. *Science* **283**, 1911–1914.
- Hajdukiewicz, P., Svab, Z., and Maliga, P. (1994). The small versatile pPZP family of *Agrobacterium* binary vectors for plant transformation. *Plant Mol. Biol.* **25**, 989–994.
- Harrison, P.T. (1996). Protein:protein interactions in the lipid bilayer. *Mol. Membr. Biol.* **13**, 67–79.
- Hurtley, S.M., and Helenius, A. (1989). Protein oligomerization in the endoplasmic reticulum. *Annu. Rev. Cell Biol.* **5**, 277–307.
- Jiang, X., Dreano, M., Buckler, D.R., Cheng, S., Ythier, A., Wu, H., Hendrickson, W.A., and Tayar, N.E. (1995). Structural predictions for the ligand-binding region of glycoprotein hormone receptors and the nature of hormone-receptor interactions. *Structure* **3**, 1341–1353.
- Jones, D.A., and Jones, J.D.G. (1996). The roles of leucine-rich repeat proteins in plant defences. *Adv. Bot. Res. Inc. Adv. Plant Pathol.* **24**, 89–167.
- Jones, D.A., Thomas, C.M., Hammond-Kosack, K.E., Balint-Kurti, P.J., and Jones, J.D. (1994). Isolation of the tomato *Cf-9* gene for resistance to *Cladosporium fulvum* by transposon tagging. *Science* **266**, 789–793.
- Kayes, J.M., and Clark, S.E. (1998). *CLAVATA2*, a regulator of meristem and organ development in *Arabidopsis*. *Development* **125**, 3843–3851.

- Laux, T., Mayer, K.F.X., Berger, J., and Jürgens, G.** (1996). The *WUSCHEL* gene is required for shoot and floral meristem integrity in *Arabidopsis*. *Development* **122**, 87–96.
- Lease, K., Ingham, E., and Walker, J.C.** (1998). Challenges in understanding RLK function. *Curr. Opin. Plant Biol.* **1**, 388–392.
- Leyser, H.M.O., and Furner, I.J.** (1992). Characterization of three shoot apical meristem mutants of *Arabidopsis thaliana*. *Development* **116**, 397–403.
- McDowell, J.M., Dhandaydham, M., Long, T.A., Aarts, M.G.M., Goff, S., Holub, E.B., and Dangi, J.L.** (1998). Intragenic recombination and diversifying selection contribute to the evolution of downy mildew resistance at the *RPP8* locus of *Arabidopsis*. *Plant Cell* **10**, 1861–1874.
- Murashige, T., and Skoog, F.** (1962). A revised medium for rapid growth and bioassays with tobacco tissue culture. *Physiol. Plant.* **15**, 473–497.
- Purugganan, M.D., and Suddith, J.I.** (1998). Molecular population genetics of the *Arabidopsis CAULIFLOWER* regulatory gene: Non-neutral evolution and naturally occurring variation in floral homeotic function. *Proc. Natl. Acad. Sci. USA* **95**, 8130–8134.
- Purugganan, M.D., and Suddith, J.I.** (1999). Molecular population genetics of floral homeotic loci: Departures from the equilibrium-neutral model at the *APETALA3* and *PISTILLATA* genes of *Arabidopsis thaliana*. *Genetics* **151**, 839–848.
- Stone, J.M., Collinge, M.A., Smith, R.D., Horn, M.A., and Walker, J.C.** (1994). Interaction of a protein phosphatase with an *Arabidopsis* serine/threonine receptor kinase. *Science* **266**, 793–795.
- Stone, J.M., Trotochaud, A.E., Walker, J.C., and Clark, S.E.** (1998). Control of meristem development by CLAVATA1 receptor kinase and kinase-associated protein phosphatase interactions. *Plant Physiol.* **117**, 1217–1235.
- Trotochaud, A.E., Hao, T., Guang, W., Yang, Z., and Clark, S.E.** (1999). The CLAVATA1 receptor-like kinase requires CLAVATA3 for its assembly into a signaling complex that includes KAPP and a Rho-related protein. *Plant Cell* **11**, 393–405.
- Wang, G.-L., Ruan, D.-L., Song, W.-Y., Sideris, S., Chen, L., Pi, L.-Y., Zhang, S., Zhang, Z., Fauquet, C., Gaut, B.S., Whalen, M.C., and Ronald, P.C.** (1998). *Xa21D* encodes a receptor-like molecule with a leucine-rich repeat domain that determines race-specific recognition and is subject to adaptive evolution. *Plant Cell* **10**, 765–779.
- Williams, R.W., Wilson, J.M., and Meyerowitz, E.M.** (1997). A possible role for kinase-associated protein phosphatase in the *Arabidopsis* CLAVATA1 signaling pathway. *Proc. Natl. Acad. Sci. USA* **94**, 10467–10472.

The Arabidopsis *CLAVATA2* Gene Encodes a Receptor-like Protein Required for the Stability of the *CLAVATA1* Receptor-like Kinase

Sangho Jeong, Amy E. Trotochaud and Steven E. Clark

Plant Cell 1999;11;1925-1933

DOI 10.1105/tpc.11.10.1925

This information is current as of October 15, 2019

References	This article cites 31 articles, 19 of which can be accessed free at: /content/11/10/1925.full.html#ref-list-1
Permissions	https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&ciissn=1532298X&WT.mc_id=pd_hw1532298X
eTOCs	Sign up for eTOCs at: http://www.plantcell.org/cgi/alerts/ctmain
CiteTrack Alerts	Sign up for CiteTrack Alerts at: http://www.plantcell.org/cgi/alerts/ctmain
Subscription Information	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: http://www.aspb.org/publications/subscriptions.cfm