

Feedback Regulation of *GA5* Expression and Metabolic Engineering of Gibberellin Levels in Arabidopsis

Yun-Ling Xu,^{a,1} Li Li,^{a,2} Douglas A. Gage,^b and Jan A. D. Zeevaart^{a,3}

^aMichigan State University—Department of Energy Plant Research Laboratory, Michigan State University, East Lansing, Michigan 48824-1312

^bDepartment of Biochemistry, Michigan State University, East Lansing, Michigan 48824-1319

The gibberellin (GA) 20-oxidase encoded by the *GA5* gene of Arabidopsis directs GA biosynthesis to active GAs, whereas that encoded by the *P16* gene of pumpkin endosperm leads to biosynthesis of inactive GAs. Negative feedback regulation of *GA5* expression was demonstrated in stems of Arabidopsis by bioactive GAs but not by inactive GA. In transgenic Arabidopsis plants overexpressing *P16*, there was a severe reduction in the amounts of C₂₀-GA intermediates, accumulation of large amounts of inactive GA₂₅ and GA₁₇, a reduction in GA₄ content, and a small increase in GA₁. However, due to feedback regulation, expression of *GA5* and *GA4*, the gene coding for the subsequent 3 β -hydroxylase, was greatly increased to compensate for the effects of the *P16* transgene. Consequently, stem height was only slightly reduced in the transgenic plants.

INTRODUCTION

Gibberellins (GAs) are tetracyclic diterpenoid compounds that play an important role in many aspects of plant growth and development, such as promotion of cell division and extension, seed germination, stem growth, flowering, and fruit set (Crozier, 1983). The pathway of GA biosynthesis has been elucidated in extensive biochemical studies, and the enzymes involved have been characterized.

The GA biosynthetic pathway can be divided into three stages according to the type of reactions and enzymes involved (Graebe, 1987; Hedden and Kamiya, 1997; Kende and Zeevaart, 1997; Phillips, 1998). Stage 1 involves the cyclization of geranylgeranyl diphosphate to *ent*-copalyl diphosphate, which in turn is converted to *ent*-kaurene. The enzymes that catalyze these reactions are called *ent*-copalyl diphosphate synthase and *ent*-kaurene synthase, which in Arabidopsis are encoded by the *GA1* (Sun and Kamiya, 1994) and *GA2* (Yamaguchi et al., 1998b) genes, respectively.

In the second stage, the P450 monooxygenase *ent*-kaurene oxidase, encoded by *GA3* of Arabidopsis (Helliwell et al., 1998), sequentially oxidizes C-19 of *ent*-kaurene via *ent*-kaurenol and *ent*-kaurenal to *ent*-kaurenoic acid, which is further oxidized to *ent*-7 α -hydroxy kaurenoic acid. This is

followed by contraction of the B ring with extrusion of C-7 to give GA₁₂-aldehyde.

The first step of stage 3 involves oxidation of GA₁₂-aldehyde to GA₁₂. Further metabolism of GA₁₂ differs among species with respect to the position and sequence of oxidative steps. Common in higher plants is the early 13-hydroxylation pathway, which involves hydroxylation at C-13 to give GA₅₃. Both GA₁₂ and GA₅₃ are substrates for the multifunctional enzyme GA 20-oxidase in parallel pathways (GA₁₂→GA₁₅→GA₂₄→GA₉ and GA₅₃→GA₄₄→GA₁₉→GA₂₀) in which C-20 is successively oxidized and eliminated. In addition, small amounts of the 20-carboxylic acids GA₂₅ and GA₁₇ are produced (Figure 1).

Three GA 20-oxidases have been isolated from Arabidopsis (Phillips et al., 1995), of which one, *GA5*, is highly expressed in elongating stems (Phillips et al., 1995; Xu et al., 1997). Next, GA₉ and GA₂₀ are converted by 3 β -hydroxylase (encoded by *GA4* in Arabidopsis) to the biologically active forms GA₄ and GA₁, respectively. Finally, these bioactive GAs are deactivated by 2 β -hydroxylation: GA₁ to GA₈ and GA₄ to GA₃₄ (Figure 1). Oxidations in stage 3 of the pathway are catalyzed by dioxygenases that use 2-oxoglutarate and molecular oxygen as cosubstrates.

With the elucidation of the GA biosynthetic pathway comes the question: how is GA biosynthesis regulated? Graebe (1987) has suggested that GA biosynthesis is activated and deactivated at different developmental stages and in different organs of higher plants. With the cloning of several GA biosynthetic genes (see Hedden and Kamiya, 1997), it is now possible to investigate the regulation of GA

¹Current address: Department of Entomology, Michigan State University, East Lansing, MI 48824-1115.

²Current address: U.S. Plant, Soil, and Nutrition Laboratory, Cornell University, Tower Road, Ithaca, NY 14853.

³To whom correspondence should be addressed. E-mail zeevaart@pilot.msu.edu; fax 517-353-9168.




Figure 1. Stage 3 of the GA Biosynthetic Pathway in Arabidopsis.

The multifunctional enzyme GA 20-oxidase (GA5) oxidizes and eliminates C-20 of GA_{12} and GA_{53} in two parallel pathways to the C_{19} -GAs GA_9 and GA_{20} , respectively. The products of these reactions are converted by 3 β -hydroxylase (GA4) to bioactive GAs: GA_9 to GA_4 and GA_{20} to GA_1 . These active GAs are inactivated by 2 β -hydroxylase to GA_{34} and GA_8 , respectively. A GA 20-oxidase of pumpkin endosperm, P16, oxidizes C-20 exclusively to 20-carboxylic acids: GA_{12} to GA_{25} and GA_{53} to GA_{17} . All GAs with C-20 as a carboxyl group are biologically inactive.

biosynthesis at the molecular level in response to both endogenous and environmental signals. In Arabidopsis, it has been found that the amount of *GA5* transcript is increased by long days, but *GA4* expression is not under photoperiodic control (Xu et al., 1997). Expression of both the *GA4* and *GA5* genes is sensitive to feedback regulation in Arabidopsis. Treatment with GA reduced the amount of both transcripts in hypocotyls, leaves, and floral shoots (Chiang et al., 1995; Phillips et al., 1995; Xu et al., 1995; Cowling et al., 1998). However, expression of *GA3* is not subject to feedback regulation by GA (Helliwell et al., 1998). Here, we

report on the expression of *GA5* in the elongating stems of GA-deficient and GA-insensitive mutants after the application of active and inactive GAs.

To obtain further insight into the feedback regulation of GA biosynthesis, we also expressed in Arabidopsis a gene coding for an enzyme that only oxidizes C-20 of GAs. GA 20-oxidases normally oxidize and eliminate C-20 of C_{20} -GAs, thus producing C_{19} -GAs (Figure 1). However, a GA 20-oxidase encoded by a cDNA isolated from pumpkin endosperm, here called *P16*, oxidizes C-20 to a carboxyl group, thus giving rise to the biologically inactive com-




Figure 2. Feedback Regulation by GA of *GA5* Expression in Stems of Wild-Type Arabidopsis and GA-Deficient Mutants.

(A) Effects of GA_4 on the accumulation of the *GA5* transcript in *ga4* and *ga5* mutants. Wild-type Landsberg *erecta* plants and the mutants *ga4* and *ga5* were sprayed with a solution of 100 μ M GA_4 . The upper 2-cm sections of the stems were harvested 24 hr later.

pounds GA_{25} and GA_{17} (Figure 1; Xu et al., 1995). It was of interest, therefore, to determine whether the constitutive expression of the pumpkin 20-oxidase, *P16*, would lower the level of active GAs and thus reduce stem growth. Our results demonstrate that expression of the transgene *P16* stimulates production of GA_{25} and GA_{17} , but this is accompanied by increased transcript levels of *GA4* and *GA5* so that an adequate level of active GAs is maintained for stem growth to occur at close to its normal rate.

RESULTS

Negative Feedback Regulation of *GA5* Expression in the Stems of Arabidopsis Plants

The *ga4* and *ga5* mutants are impaired in the final steps of the GA biosynthetic pathway (Figure 1; Koornneef and van der Veen, 1980) so that these mutants have reduced levels of the 3β -hydroxy-GAs GA_1 and GA_4 (Talon et al., 1990a). We found that the level of *GA5* transcript in elongating stems of these mutants was considerably higher than in stems of Landsberg *erecta* plants. However, when GA_4 was applied, the level of *GA5* transcript was reduced in the mutants as well as in stems of wild-type plants (Figure 2A). These results demonstrate that expression of *GA5* in the stems of Arabidopsis is under negative feedback regulation by GA. Thus, the reduced stature of *ga4* and *ga5* plants is due to lower levels of active GAs and is accompanied by a higher level of *GA5* mRNA. This negative feedback regulation of *GA5* transcript level is exerted by active GAs, such as GA_3 and GA_4 , but not by inactive GA_{17} (Figure 2B).

The GA-insensitive mutant, *gai*, has a semidwarf stature (Koornneef et al., 1985) and increased levels of active GAs

Poly(A)⁺-enriched RNA (2 to 3 μ g) isolated from stems was separated by electrophoresis. The blot was hybridized successively with antisense *GA5* and actin mRNA. The ratio of *GA5*/actin mRNA in the different treatments is indicated at the top.

(B) Effects of GA_3 , GA_4 , and GA_{17} on the accumulation of the *GA5* transcript in the *ga4* mutant. *ga4* plants were sprayed with a solution of 100 μ M GA_3 , GA_4 , or GA_{17} . The upper 2-cm sections of the stems were harvested 24 hr later. Poly(A)⁺-enriched RNA (2 to 3 μ g) isolated from stems was separated by electrophoresis. The blot was hybridized successively with antisense *GA5* and actin mRNA. The ratio of *GA5*/actin mRNA in the different treatments is indicated at top.

(C) Effects of GA_4 on the accumulation of the *GA5* transcript in the *gai* mutant. *gai* plants were sprayed with a solution of 100 μ M GA_4 . The upper 2-cm sections of the stems were harvested 24 hr later. Poly(A)⁺-enriched RNA (2 to 3 μ g) isolated from stems was separated by electrophoresis. The blot was hybridized successively with antisense *GA5* and actin mRNA. The ratio of *GA5*/actin is indicated at the top.

compared with wild-type plants (Talon et al., 1990b). It was found in these experiments that *GA5* mRNA accumulated to a higher level in stems of the *gai* mutant than in stems of wild-type plants (Figure 2C). Thus, in contrast to the GA-deficient *ga4* and *ga5* mutants, a higher level of active GAs in the *gai* mutant is accompanied by a higher level of *GA5* mRNA. However, application of GA_4 reduced accumulation of *GA5* mRNA in stems of the *gai* mutant (Figure 2C).

Generation and Characterization of Transgenic Arabidopsis Plants Overexpressing *P16*

In the GA biosynthetic pathway, successive oxidation of C-20 of C_{20} -GAs leads to the loss of this carbon atom as CO_2 and to the formation of C_{19} -GAs, which subsequently are converted to biologically active GAs (Figure 1). In addition, C_{20} -GAs can also be converted to inactive C-20 carboxylic acids. The GA 20-oxidase encoded by *P16* from pumpkin endosperm is functionally different from other GA 20-oxidases in that it converts C_{20} -GAs predominantly to the inactive C-20 carboxylic acids GA_{17} and GA_{25} (Lange et al., 1994; Xu et al., 1995). We introduced the *P16* gene under control of the cauliflower mosaic virus 35S promoter (Figure 3) into Arabidopsis to determine whether this would result in redirecting the flow in the GA biosynthetic pathway to inactive forms (Figure 1), in the expectation that this would result in plants with reduced stem height.

Molecular Analysis of Transgenic Plants

To confirm that the transgenic plants did carry the *P16* gene, we performed DNA gel blot analysis with genomic DNA that

had been digested with BamHI. Upon electrophoresis and hybridization of the blot with ^{32}P -labeled *P16*, the digests of all three transgenic lines gave a single band (Figure 4), indicating that each transgenic line carried a single copy of the *P16* gene. The fact that there is a BamHI site at the junction of the cauliflower mosaic virus 35S promoter and the transgene excludes the possibility that the transgenic plants carried tandem repeats. Hybridized bands among the three transgenic lines differed in size, indicating that *P16* was integrated at different sites of the genome.

To determine whether the transgene was transcribed, we analyzed mRNA from stems by RNA gel blotting. As shown in Figure 5A, the expression level of *P16* in plants of lines 5A and 5H was very high; in plants of line 5D, *P16* mRNA was barely detectable. Protein gel blot analysis was also performed to check for the formation of P16 protein in the transgenic plants. Figure 5B shows that plants of lines 5A and 5H accumulated the P16 protein, whereas 5D had no detectable amount of P16 protein. This indicates that *P16* in line 5D is not well expressed in its integrated chromosomal matrix.

Upregulation of *GA5* and *GA4* Expression in Transgenic Plants

To determine expression of the *GA5* and *GA4* loci in the transgenic plants, we conducted RNA gel blotting with mRNA from stems. The results presented in Figure 5A show that *GA5* and *GA4* transcript levels had increased markedly in lines 5A and 5H but not in line 5D. This demonstrates that the expression of the native genes for both GA 20-oxidase (*GA5*) and 3 β -hydroxylase (*GA4*) was significantly upregulated in two of the three lines of transgenic plants.




Figure 3. Structure of the pBI3516 Plasmid Used for Transformation of Arabidopsis.

Schematic presentation of the construct p35S-*P16* used for expression of GA 20-oxidase from pumpkin endosperm in Arabidopsis. The β -glucuronidase (*GUS*) gene was removed from pBI121 by digestion with BamHI and SstI, and the *P16* gene was inserted in the same sites. The cauliflower mosaic virus (CaMV) 35S promoter and the nopaline synthase terminator signal (*NOS-t*) are also indicated. The resulting plasmid was designated pBI3516.




Figure 4. Genomic DNA Gel Blot Analysis of Wassilewskija and Three Transgenic Lines.

Genomic DNA from wild-type Wassilewskija (Ws) plants and the transgenic lines 5A, 5D, and 5H was digested with BamHI and hybridized with ^{32}P -labeled *P16* cDNA. All three transgenic lines show a single band that hybridizes with *P16*, indicating that each line contains a single copy of the transgene, because *P16* does not contain a BamHI restriction site. Numbers at right indicate the positions of the markers in kilobases.

Phenotype of Transgenic Plants

Stem height of plants of lines 5A and 5H was only slightly reduced, whereas that of plants of lines 5D was not reduced at all compared with that of wild-type plants (Figure 6). No other morphological differences were observed between wild-type and transgenic plants.

Analysis of GAs in Transgenic Plants

A quantitative analysis of the endogenous GAs showed that there were large decreases in the 20-methyl (GA_{53}), 20-hydroxy lactone (GA_{44}), and 20-aldehyde (GA_{24} and GA_{19}) GAs (Table 1) and large relative increases in the 20-carboxylic acids GA_{25} and GA_{17} (Table 2) in all three transgenic lines but especially in lines 5A and 5H. These results demonstrate that the *P16* transgene is not only expressed at the mRNA

level but that an active enzyme is produced with the same catalytic properties in *Arabidopsis* as reported for the recombinant protein (Lange et al., 1994; Xu et al., 1995).

The non-13-hydroxylated C_{19} -GAs GA_9 and GA_4 were considerably reduced in the transgenic plants, especially in lines 5A and 5H. By contrast, the 13-hydroxylated C_{19} -GAs GA_{20} , GA_1 , and GA_8 showed slight increases. This trend was observed in three separate experiments. Much less *P16*




Figure 5. Expression of *GA5*, *P16*, *GA4*, and Actin mRNA in Stems of Wassilewskija and Three Transgenic Lines.

(A) RNA gel blot analysis. Poly(A)⁺-enriched RNA (2 to 3 μg) isolated from stems was separated by electrophoresis. The blot was hybridized successively with *Arabidopsis GA5* and *GA4* antisense RNA, antisense *P16* of pumpkin, and *Arabidopsis actin*. The ratio of *GA5*/actin mRNA in plants of wild-type Wassilewskija (Ws) and in three transgenic lines is indicated at top.

(B) Protein gel blot analysis. Approximately 30 μg of protein isolated from leaves was loaded per lane and analyzed by electrophoresis. The blot was cross-reacted with an antiserum raised against the *P16* protein purified from pumpkin endosperm.




Figure 6. Stem Growth of Wassilewskija and Three Transgenic Lines Expressing the *P16* Transgene.

The plants were grown initially under SD conditions and later transferred to LD conditions consisting of a 9-hr photoperiod of high photon flux density, which was extended with light from incandescent lamps for 15 hr, for a total of 24 hr a day. Days are numbered from the time of transfer to LD conditions. Twenty plants of ecotype Wassilewskija (*Ws*) and 20 plants of each transgenic line were measured. Error bars indicate the standard error of the mean.

mRNA accumulated in line 5D than in lines 5A and 5H (Figure 5A), and these differences in *P16* transcript levels were correlated with the observed changes in GA levels (Tables 1 and 2).

As shown in Table 2, there were very large relative increases in the levels of GA_{17} and GA_{25} in transgenic plants expressing *P16* compared with the levels found in wild-type plants. The greater increases in GA_{17} than in GA_{25} should not be interpreted to mean that more GA_{17} than GA_{25} is produced in the transgenic plants. In fact, the opposite is most likely the case, because the non-13-hydroxylated pathway is the main GA biosynthetic pathway in Arabidopsis (Talon et al., 1990a). Consequently, more GA_{25} than GA_{17} is found in wild-type Arabidopsis; thus, normalization of the data (see footnotes in Table 2) to those of the wild type starts with a much higher base value for GA_{25} than it does for GA_{17} .

DISCUSSION

In these experiments, we have demonstrated that negative feedback regulation of *GA5* expression operates at the level

of transcription in elongating stems of Arabidopsis and presumably plays an important role in maintaining the plant's height within a certain range. Expression of *GA5* increased when the endogenous levels of active GAs were reduced by either genetic (GA-deficient mutants; Figures 2A and 2B) or chemical (GA biosynthesis inhibitors; data not shown) methods. Application of GA_4 or GA_3 reduced the accumulation of *GA5* mRNA (Figure 2B). However, neither application of GA_{17} (Figure 2B) nor increased endogenous levels of GA_{17} or GA_{25} (Table 2) reduced accumulation of *GA5* mRNA, clearly demonstrating that active GAs are required not only for growth but also for feedback regulation of *GA5* expression. Similarly, it has been shown in Arabidopsis that GAs that promote hypocotyl elongation can downregulate *GA4* transcript abundance, whereas inactive GAs and GA analogs cannot (Cowling et al., 1998).

Both GA_4 and GA_3 promoted stem elongation and reduced the accumulation of *GA5* mRNA (Figure 2B). Yet GA_3 has not been identified in Arabidopsis (Talon et al., 1990a). This indicates that the receptor for transduction of the signal that regulates *GA5* expression and growth is not highly specific. In this respect, the signal transduction pathway appears quite similar, if not the same, for GAs regulating stem elongation and *GA5* expression. This raises the question of whether *GA5* expression is regulated by changes in stem growth itself or more directly by GA signal transduction. In GA-deficient mutants impaired in the genes for enzymes that catalyze the final steps in the pathway, a correlation exists between increased expression of *GA5*, a decrease in the levels of active GAs, and reduced stem height (Talon et al., 1990a; Phillips et al., 1995; Xu et al., 1995). However, this correlation does not hold in the GA-insensitive mutant *gai*, which has a semidwarf phenotype. More *GA5* transcript accumulated in this mutant than in wild-type plants (Figure 2C;

Table 1. GA Content of Arabidopsis Ecotype Wassilewskija and of Three Transgenic Lines Transformed with *P16* of Pumpkin Endosperm

GAs	Ws ^a	5A ^a	5D ^a	5H ^a
Non-13-OH-GAs				
GA_{24}	59.0	0.1	4.8	0.0
GA_9	2.0	0.5	2.4	0.5
GA_4	6.2	1.7	2.1	1.3
13-OH-GAs				
GA_{53}	13.0	0.2	0.9	0.2
GA_{44}	1.8	0.4	1.8	0.3
GA_{19}	11.8	2.1	7.6	1.7
GA_{20}	0.2	0.3	0.4	0.3
GA_1	0.2	0.3	0.3	0.3
GA_8	19.6	25.0	21.2	24.3

^a Nanograms of different GAs per gram dry weight of Wassilewskija (*Ws*) and three transgenic lines.

Table 2. Increase in Levels of GA₂₅ and GA₁₇ in Three Transgenic Lines Transformed with *P16* of Pumpkin Endosperm Compared with That of Ecotype Wassilewskija

GAs	Ws ^a	5A ^a	5D ^a	5H ^a
Non-13-OH-GA				
GA ₂₅ : <i>m/z</i> 312/316 ^b	1	65	58	75
13-OH-GA				
GA ₁₇ : <i>m/z</i> 492/436 ^c	1	378	244	643

^a Fold increase in transgenic plants compared with Wassilewskija (Ws).

^b *m/z* 312 of GA₂₅/*m/z* 316 of ²H-labeled GA₂₄, normalized to 1 for Ws.

^c *m/z* 492 of GA₁₇/*m/z* 436 of ²H-labeled GA₁₉, normalized to 1 for Ws.

Peng et al., 1997), despite the higher levels of active GAs in the mutant, indicating that GAI plays a role in the regulation of *GA5* expression. In previous work, treatment of *gai* plants with GA₄ caused a slight increase in GA5 mRNA in leaves (Xu et al., 1995). However, here we observed that application of a high concentration of GA₄ decreased *GA5* RNA in stems of *gai* plants (Figure 2C). This difference in *GA5* expression may be due to differential regulation of *GA5* in stems and leaves of *gai* plants. It is also important to note that the *gai* mutant is not completely insensitive to GA. In the double mutant *gai gai 1*, in which the GA level is reduced, the phenotype becomes more extreme dwarf (Koorneef et al., 1985). A similar extreme phenotype is seen after treatment with the GA biosynthesis inhibitor paclobutrazol (Peng et al., 1997).

Using GA biosynthesis inhibitors to control plant stature is a common practice in agriculture. Molecular cloning of genes for GA biosynthetic enzymes (Hedden and Kamiya, 1997) makes it possible to target plant stature for genetic engineering. Because several studies have shown that steps catalyzed by GA 20-oxidase may be limiting GA biosynthesis (Coles et al., 1996; Hedden and Kamiya, 1997; Xu et al., 1997), GA 20-oxidase was chosen as an engineering target. The P16 protein, a fusion GA 20-oxidase encoded by a cDNA from pumpkin endosperm, produces inactive tricarboxylic acid GAs as its major products *in vitro* (Lange et al., 1994; Xu et al., 1995). It was expected that overexpression of *P16* would redirect the GA biosynthetic flow to inactive GA₁₇ and GA₂₅, leading to a reduction in active GA₁ and GA₄ (Figure 1). Accumulation of GA₁₇ and GA₂₅ was indeed found in transgenic plants overexpressing *P16* (Table 2), but only small changes in stem height were observed; *GA5* and *GA4* mRNA accumulated to very high levels in these plants (Figure 5A). Thus, GA homeostasis was maintained by upregulating expression of the native genes in the transgenic plants.

Analysis of the endogenous GA content in *P16*-overexpressing plants showed a severe reduction in the amounts of C₂₀-GA intermediates with a large accumulation of inactive tricarboxylic acid GAs (Table 2). Although there was a substantial decrease in the amount of active GA₄, only a small effect on stem growth was observed in lines 5A and 5H, and

no effect at all was observed in line 5D (Table 1 and Figure 6). The small increases in GA₁ and substantial decreases in GA₄ in the transgenic plants further support the conclusion that GA₄ is the major bioactive GA for stem elongation in *Arabidopsis* (Xu et al., 1997). Apparently, endogenous GA₄ is saturating for stem growth in wild-type plants, and a considerable reduction in its content is required before it becomes limiting for growth. Another possibility is that sensitivity to GA increases as the levels of active GAs are reduced. This would mean that in the *P16*-overexpressing plants, GA sensitivity is increased so that their stature is maintained relatively unchanged, despite reduced levels of GA₄.

P16 was expressed much less at the mRNA and protein levels in line 5D than in lines 5A and 5H (cf. Figures 5A and 5B). Nevertheless, the C₂₀-GAs GA₂₄ and GA₅₃ were much reduced in plants of line 5D compared with wild-type plants, whereas GA₁₉ showed only a slight decrease and GA₄₄ remained unchanged (Table 1). Considering the very low levels of C₂₀-GAs and the very high levels of GA₁₇ and GA₂₅ in plants of lines 5A and 5H (Table 2), it appears that the *P16* enzyme had reached saturating levels in these two transgenic lines, although this was clearly not the case in line 5D. Although no *P16* protein was detectable by immunoblotting in plants of line 5D, an adequate amount of enzyme must have been produced to cause the large increases in GA₁₇ and GA₂₅ (Table 2).

Reduced plant height is a desirable trait in modern agriculture. In principle, two strategies can be envisioned by which stem height in plants can be reduced by metabolic engineering of GA levels: (1) by expressing GA biosynthetic genes in the antisense orientation and (2) by overexpressing genes that deactivate GAs. In *Arabidopsis*, we overexpressed *P16* from pumpkin endosperm, which catalyzes the production of inactive 20-carboxylic acids, and observed only a slight decrease in stem height due to feedback regulation that resulted in much increased transcript levels of the native *GA4* and *GA5* genes. Thus, upregulation of *GA4* and *GA5* expression, and presumably of the corresponding enzymes as well, compensated for the increased deactivation of GAs by *P16*. Other target enzymes for genetic engineering of reduced GA levels and plant stature include 2β-hydroxylase and GA-conjugating enzymes. Aside from feedback regulation, the success or failure of this approach depends on the affinity of two competing enzymes, with one activating and the other deactivating a common substrate, such as GA₉, to either GA₄, an active GA, or GA₅₁, an inactive GA.

METHODS

Plant Material and Gibberellin Treatment

Plants (*Arabidopsis thaliana*) were grown as described previously (Talon et al., 1990a). All mutants used were in ecotype Landsberg

erecta. Short-day (SD) conditions consisted of 9 hr of light from fluorescent tubes and incandescent lamps (150 to $200 \mu\text{mol m}^{-2} \text{sec}^{-1}$) at 23°C , followed by 15 hr of darkness at 20°C . The plants were grown under SD conditions for 4 weeks and then transferred to long-day (LD) conditions. LD conditions consisted of the same photoperiod as did SD conditions but were extended with 15 hr of low-intensity light from incandescent lamps ($10 \mu\text{mol m}^{-2} \text{sec}^{-1}$) at 20°C . When required, plants were sprayed with an aqueous solution of $100 \mu\text{M}$ of the gibberellins GA_4 , GA_3 , or GA_{17} containing 5% (v/v) ethanol and 0.05% Tween 20.

For DNA and protein extraction, rosette leaves were harvested before stem elongation. For mRNA extraction, the upper 2-cm portion of the stem was harvested when the stems were 2 to 5 cm long. All material was harvested at the end of the main light period.

Plants to be used for GA measurements were initially grown in SD conditions for 6 weeks from sowing and then exposed to 3 LD. At the time of harvest, all plants had stems with flower buds. The entire plant (rosette leaves and stems with leaves and flower buds) was collected, frozen in liquid N_2 , and lyophilized.

Construction of Plasmid and Transformation of Arabidopsis

Plasmid pP16, which contains the GA 20-oxidase cDNA from pumpkin endosperm (Xu et al., 1995), was digested with BamHI and SstI, and *P16* was inserted into the same sites of pBI121 (Jefferson et al., 1987) (Clontech, Palo Alto, CA). The resulting plasmid was designated pBI3516 (Figure 3). This plasmid was introduced into *Agrobacterium tumefaciens* LBA4404 (Clontech).

For transformation of Arabidopsis ecotype Wassilewskija, cotyledons were removed from 7-day-old seedlings and cocultured with LBA4404 harboring pBI3516. Transgenic plants (T_0) were regenerated from these cotyledon explants, as described previously (Barghchi et al., 1994). All analyses were performed on T_2 plants, which had been tested for homozygosity in the T_3 generation.

DNA Isolation and Gel Blot Analysis

Genomic DNA was extracted as described previously (see Focus, Volume 12, published by Bethesda Research Laboratories). Digested DNA samples were separated by electrophoresis in a 0.8% agarose gel and transferred to nylon membranes. The blots were prehybridized and hybridized in $0.5 \text{ M Na}_2\text{HPO}_4$, pH 7.2, 7% SDS, and 1 mM EDTA at 68°C and washed in $40 \text{ mM Na}_2\text{HPO}_4$, pH 7.2, 5% SDS, and 1 mM EDTA at 68°C . They were then exposed to a PhosphorImager screen (Molecular Dynamics, Sunnyvale, CA).

mRNA Isolation and RNA Gel Blot Analysis

Total RNA was extracted by using an Extract-A-Plant RNA isolation kit (Clontech). mRNA was then purified by using a PolyAtract mRNA isolation system (Promega). RNA gel blots were prepared by electrophoresis of 2 to 3 μg of poly(A)⁺ RNA in agarose gels in the presence of formaldehyde (Sambrook et al., 1989), followed by transfer to nylon membranes. The blots were probed with ^{32}P -labeled antisense RNA. Hybridization was performed as given for DNA gel blotting except that the temperature was 70°C . The blots were washed in $0.1 \times \text{SSC}$ ($1 \times \text{SSC}$ is 0.15 M NaCl and $0.015 \text{ M sodium citrate}$) at 70°C and exposed to a PhosphorImager screen. The relative amounts of mRNA were determined with a PhosphorImager. Each blot was se-

quentially probed with *GA5*, *GA4* (Chiang et al., 1995), *P16*, or actin (clone 40F11 from Dr. T. Newman, Michigan State University) anti-sense RNA probes after the blots were stripped. The ratio of *GA5*/actin mRNA was used to compare *GA5* mRNA levels in different treatments. Each experiment was repeated at least twice with similar results.

The hybridization conditions used did not cause cross-hybridization with transcripts of homologs of *GA5* (Phillips et al., 1995; Y.-L. Xu, unpublished results). A homolog of *GA4*, *GA4H*, has been isolated that was predominantly expressed during Arabidopsis seed germination. Cross-hybridization between the mRNAs of these two genes was below the level of detection (Yamaguchi et al., 1998a).

Protein Extraction and Protein Gel Blot Analysis

Protein from pumpkin (*Cucurbita maxima*) endosperm was precipitated with ammonium sulfate, and the 40 to 70% fraction with GA 20-oxidase activity was purified by anion exchange, gel filtration, and hydrophobic chromatography. The partially purified enzyme preparation was subjected to semipreparative native PAGE, and gel sections were assayed for enzyme activity. Proteins in the zone with the highest enzyme activity were eluted and used for immunization of rabbits. Polyclonal antibodies were obtained that immunoprecipitated the GA 20-oxidase activity. These antibodies were used to isolate the *P16* cDNA clone, which encodes a GA 20-oxidase, by immunoscreening a cDNA library made from pumpkin endosperm (Xu et al., 1995).

Proteins were extracted from Arabidopsis rosette leaves by using 62.5 mM Tris-HCl buffer at pH 6.8. Protein concentration was estimated by the Bradford (1976) method. Cold (-20°C) acetone was then added to a final concentration of 80% (v/v) to precipitate proteins. Precipitated proteins were dissolved in loading buffer (62.5 mM Tris-HCl , pH 6.8, 5% SDS, 10 mM DTT, and 10% glycerol).

Extracted proteins were electrophoresed on 10% SDS-polyacrylamide gels. After electrophoresis, the polypeptides were electrotransferred to a nitrocellulose filter. The filter was developed with the ProtoBlot II AP system (Promega).

Quantification of GAs

The GAs were extracted, purified, and quantified by the use of deuterated GAs as internal standards (Talon et al., 1990a), except for the tricarboxylic acids GA_{25} and GA_{17} , which were not available in deuterated form. Because these GAs were collected in the same fractions as their corresponding 20-aldehydes, GA_{24} and GA_{19} , they were quantitatively related to the internal standards ^2H -labeled GA_{24} and ^2H -labeled GA_{19} by gas chromatography-selected ion monitoring. The ions used for calculating the increases in GA_{25} and GA_{17} in transgenic plants in comparison with Wassilewskija represent major fragments in the mass spectra of GA_{25} , GA_{17} , and ^2H -labeled GA_{24} and GA_{19} (see footnotes in Table 2).

ACKNOWLEDGMENTS

The original seeds of the Arabidopsis ecotype Wassilewskija (CS915) were supplied by the Arabidopsis Biological Resource Center (Ohio

State University, Columbus). The authors are grateful to Marlene Cameron and Kurt Stepnitz for help in preparing the figures. This work was supported by the U.S. Department of Energy (Grant No. DE-FG02-91ER20021), the U.S. Department of Agriculture (Grant No. 94-37304-0956), and a grant from the National Institutes of Health (Grant No. RR-00480) to the Michigan State University–National Institutes of Health Mass Spectrometry Facility.

Received November 24, 1998; accepted March 9, 1999.

REFERENCES

- Barghchi, M., Turgut, K., Scott, R., and Draper, J. (1994). High-frequency transformation from cultured cotyledons of *Arabidopsis thaliana* ecotypes "C24" and "Landsberg erecta." *Plant Growth Regul.* **14**, 61–67.
- Bradford, M.M. (1976). A rapid sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein–dye binding. *Anal. Biochem.* **72**, 248–254.
- Chiang, H.-H., Hwang, I., and Goodman, H.M. (1995). Isolation of the *Arabidopsis* *GA4* locus. *Plant Cell* **7**, 195–201.
- Coles, J.P., Phillips, A.L., Valero, D., and Hedden, P. (1996). Genetic manipulation of gibberellin biosynthesis in *Arabidopsis*. *Plant Physiol.* **111** (suppl.), 49 (abstr.).
- Cowling, R.J., Kamiya, Y., Seto, H., and Harberd, N.P. (1998). Gibberellin dose–response regulation of *GA4* gene transcript levels in *Arabidopsis*. *Plant Physiol.* **117**, 1195–1203.
- Crozier, A. (1983). *The Biochemistry and Physiology of Gibberellins*, Vol. 2. (New York: Praeger).
- Graebe, J.E. (1987). Gibberellin biosynthesis and control. *Annu. Rev. Plant Physiol.* **38**, 419–465.
- Hedden, P., and Kamiya, Y. (1997). Gibberellin biosynthesis: Enzymes, genes and their regulation. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **48**, 431–460.
- Helliwell, C.A., Sheldon, C.C., Olive, M.R., Walker, A.R., Zeevaart, J.A.D., Peacock, W.J., and Dennis, E.S. (1998). Cloning of the *Arabidopsis* *ent*-kaurene oxidase gene *GA3*. *Proc. Natl. Acad. Sci. USA* **95**, 9019–9024.
- Jefferson, R.A., Kavanagh, T.A., and Bevan, M.W. (1987). GUS fusions: β -Glucuronidase, a sensitive and versatile gene fusion marker in higher plants. *EMBO J.* **6**, 3901–3907.
- Kende, H., and Zeevaart, J.A.D. (1997). The five "classical" plant hormones. *Plant Cell* **9**, 1197–1210.
- Koornneef, M., and van der Veen, J.H. (1980). Induction and analysis of gibberellin sensitive mutants in *Arabidopsis thaliana* (L.) Heynh. *Theor. Appl. Genet.* **58**, 257–263.
- Koornneef, M., Elgersma, A., Hanhart, C.J., van Loenen-Martinet, E.P., van Rijn, L., and Zeevaart, J.A.D. (1985). A gibberellin insensitive mutant of *Arabidopsis thaliana*. *Physiol. Plant.* **65**, 33–39.
- Lange, T., Hedden, P., and Graebe, J. (1994). Expression cloning of a gibberellin 20-oxidase, a multifunctional enzyme involved in gibberellin biosynthesis. *Proc. Natl. Acad. Sci. USA* **91**, 8552–8556.
- Peng, J., Carol, P., Richards, D.E., King, K.E., Cowling, R.J., Murphy, G.P., and Harberd, N.P. (1997). The *Arabidopsis* *GA1* gene defines a signaling pathway that negatively regulates gibberellin responses. *Genes Dev.* **11**, 3194–3205.
- Phillips, A.L. (1998). Gibberellins in *Arabidopsis*. *Plant Physiol. Biochem.* **36**, 115–124.
- Phillips, A.L., Ward, D.A., Uknes, S., Appleford, N.E.L., Lange, T., Huttly, A.K., Gaskin, P., Graebe, J.E., and Hedden, P. (1995). Isolation and expression of three gibberellin 20-oxidase cDNA clones from *Arabidopsis*. *Plant Physiol.* **108**, 1049–1057.
- Sambrook, J., Fritsch, E.F., and Maniatis, T. (1989). *Molecular Cloning: A Laboratory Manual*, 2nd ed. (Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press).
- Sun, T.-p., and Kamiya, Y. (1994). The *Arabidopsis* *GA1* locus encodes the cyclase *ent*-kaurene synthetase A of gibberellin biosynthesis. *Plant Cell* **6**, 1509–1518.
- Talon, M., Koornneef, M., and Zeevaart, J.A.D. (1990a). Endogenous gibberellins in *Arabidopsis thaliana* and possible steps blocked in the biosynthetic pathways of the semidwarf *ga4* and *ga5* mutants. *Proc. Natl. Acad. Sci. USA* **87**, 7983–7987.
- Talon, M., Koornneef, M., and Zeevaart, J.A.D. (1990b). Accumulation of C_{19} -gibberellins in the gibberellin-insensitive dwarf mutant *gai* of *Arabidopsis thaliana* (L.) Heynh. *Planta* **182**, 501–505.
- Xu, Y.-L., Li, L., Wu, K., Peeters, A.J.M., Gage, D.A., and Zeevaart, J.A.D. (1995). The *GA5* locus of *Arabidopsis thaliana* encodes a multifunctional gibberellin 20-oxidase: Molecular cloning and functional expression. *Proc. Natl. Acad. Sci. USA* **92**, 6640–6644.
- Xu, Y.-L., Gage, D.A., and Zeevaart, J.A.D. (1997). Gibberellins and stem growth in *Arabidopsis thaliana*: Effects of photoperiod on expression of the *GA4* and *GA5* loci. *Plant Physiol.* **114**, 1471–1476.
- Yamaguchi, S., Smith, M.W., Brown, R.G.S., Kamiya, Y., and Sun, T.-p. (1998a). Phytochrome regulation and differential expression of gibberellin 3 β -hydroxylase genes in germinating *Arabidopsis* seeds. *Plant Cell* **10**, 2115–2126.
- Yamaguchi, S., Sun, T.-p., Kawaide, H., and Kamiya, Y. (1998b). The *GA2* locus of *Arabidopsis thaliana* encodes *ent*-kaurene synthase of gibberellin biosynthesis. *Plant Physiol.* **116**, 1271–1278.

Feedback Regulation of GA5 Expression and Metabolic Engineering of Gibberellin Levels in *Arabidopsis*

Yun-Ling Xu, Li Li, Douglas A. Gage and Jan A. D. Zeevaart
Plant Cell 1999;11;927-935
DOI 10.1105/tpc.11.5.927

This information is current as of November 27, 2020

References	This article cites 22 articles, 14 of which can be accessed free at: /content/11/5/927.full.html#ref-list-1
Permissions	https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&ciissn=1532298X&WT.mc_id=pd_hw1532298X
eTOCs	Sign up for eTOCs at: http://www.plantcell.org/cgi/alerts/ctmain
CiteTrack Alerts	Sign up for CiteTrack Alerts at: http://www.plantcell.org/cgi/alerts/ctmain
Subscription Information	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: http://www.aspb.org/publications/subscriptions.cfm