

RESEARCH ARTICLE

# *slender* Rice, a Constitutive Gibberellin Response Mutant, Is Caused by a Null Mutation of the *SLR1* Gene, an Ortholog of the Height-Regulating Gene *GAI/RGA/RHT/D8*

Akira Ikeda,<sup>a,b,1</sup> Miyako Ueguchi-Tanaka,<sup>a</sup> Yutaka Sonoda,<sup>a,1</sup> Hidemi Kitano,<sup>a</sup> Masaji Koshioka,<sup>c</sup> Yuzo Futsuhara,<sup>b</sup> Makoto Matsuoka,<sup>a</sup> and Junji Yamaguchi<sup>a,1,2</sup>

<sup>a</sup> BioScience Center and Graduate School of Bioagricultural Sciences, Nagoya University, Chikusa-ku, Nagoya 464-8601, Japan

<sup>b</sup> Department of Agriculture, Meijo University, Tenpaku, Nagoya 468-8503, Japan

<sup>c</sup> National Research Institute of Vegetables, Ornamental Plants, and Tea, 60 Kusawa, Ano, Age, Mie 514-2328, Japan

The rice *slender* mutant (*slr1-1*) is caused by a single recessive mutation and results in a constitutive gibberellin (GA) response phenotype. The mutant elongates as if saturated with GAs. In this mutant, (1) elongation was unaffected by an inhibitor of GA biosynthesis, (2) GA-inducible  $\alpha$ -amylase was produced by the aleurone layers without gibberellic acid application, and (3) endogenous GA content was lower than in the wild-type plant. These results indicate that the product of the *SLR1* gene is an intermediate of the GA signal transduction pathway. *SLR1* maps to *OsGAI* in rice and has significant homology with height-regulating genes, such as *RHT-1Da* in wheat, *D8* in maize, and *GAI* and *RGA* in *Arabidopsis*. The *GAI* gene family is likely to encode transcriptional factors belonging to the *GRAS* gene superfamily. DNA sequence analysis revealed that the *slr1-1* mutation is a single basepair deletion of the nuclear localization signal domain, resulting in a frameshift mutation that abolishes protein production. Furthermore, introduction of a 6-kb genomic DNA fragment containing the wild-type *SLR1* gene into the *slr1-1* mutant restored GA sensitivity to normal. These results indicate that the *slr1-1* mutant is caused by a loss-of-function mutation of the *SLR1* gene, which is an ortholog of *GAI*, *RGA*, *RHT*, and *D8*. We also succeeded in producing GA-insensitive dwarf rice by transforming wild-type rice with a modified *SLR1* gene construct that has a 17–amino acid deletion affecting the DELLA region. Thus, we demonstrate opposite GA response phenotypes depending on the type of mutations in *SLR1*.

## INTRODUCTION

Gibberellins (GAs) have an important role in the regulation of many physiologic processes in the growth and development of plants, such as seed germination, shoot/stem elongation, and flower development. Changes in both GA concentration and tissue sensitivity to GA influence these processes. The molecular mechanisms by which the GA signal is transduced into morphologic and biochemical changes in plants, however, are largely unknown.


Many studies on the detection of and response to GA are focused on identifying the mutants that affect these processes. GA response mutants isolated from various plant species fall into two phenotypic categories: elongated *slen-*

*der*-type mutants and GA-unresponsive dwarf mutants (Hooley, 1994; Swain and Olszewski, 1996). The *slender* mutants have constitutive activation of their GA response, and the dwarf mutants are deficient in GA detection or signal transduction. The *slender* mutant of barley, which is homozygous for the recessive *sln1* alleles, is characterized by a rapid growth rate and long leaf sheaths (Foster, 1977). *slender* barley does not respond to growth retardants such as ancymidol (Lanahan and Ho, 1988) and paclobutrazol (Crocker et al., 1990). During germination, aleurone layers of *slender* barley synthesize  $\alpha$ -amylase in the absence of exogenous GA (Lanahan and Ho, 1988). The mutant contains a smaller amount of active GAs compared with wild-type plants, indicating that GAs are not involved in the growth of *slender* barley. Thus, the *slender* mutation might be important for elucidating the GA signal transduction pathway.

In rice, many dwarf mutants have been characterized and classified as dwarf and GA deficient, dwarf and GA insensitive, or dwarf due to other reasons (Mitsunaga et al., 1994).

<sup>1</sup> Current address: Division of Biological Sciences, Graduate School of Science, Hokkaido University, N10, W8, Sapporo 060-0810, Japan.

<sup>2</sup> To whom correspondence should be addressed. E-mail jiyama@sci.hokudai.ac.jp; fax 81-11-706-2737.



**Figure 1.** Phenotype of *slender* Rice and Its Original Wild Type, Nipponbare.

**(A)** *slender* rice was isolated as a tall mutant from rice cv Nipponbare. The plants homozygous for the recessive *slender* (*slr1-1*) gene segregated at a 1:3 ratio in 2-week-old plants.

**(B)** Shoot elongation of wild type treated with (middle) or without (left) 10  $\mu\text{M}$  GA<sub>3</sub> and *slender* without GA<sub>3</sub> treatment (right). The photograph was taken 1 week after germination.

**(C)** An elongated internode of a 2-week-old *slender* plant. The arrowhead indicates the third internode.

**(D)** Root morphology of 1-week-old plants: wild type (left), *slender* (middle), and wild type treated with 10  $\mu\text{M}$  GA<sub>3</sub> (right).

A *slender*-type mutant has not been reported in rice, however, except for the *awaodori* (*ao-1*) mutant (Nakamura, 1992).

Several GA signaling intermediates have been identified by mutation analyses in a number of plant species (Taylor, 1998), and some corresponding genes were recently cloned. In Arabidopsis, *SPINDLY* (*SPY*) acts as a negative regulator of the GA response. The deduced amino acid sequence of *SPY* suggests that the protein is an *O*-linked *N*-acetylglucosamine transferase that might glycosylate other molecules involved in GA signaling (Thornton et al., 1999). A second GA signaling intermediate from Arabidopsis is encoded by the *GA-INSENSITIVE* (*GAI*) gene, which has also been cloned (Peng et al., 1997). The original *gai* allele causes dwarfism and behaves genetically as a gain-of-function mutation (Peng and Harberd, 1993). Recent molecular analyses confirmed that this allele encodes a constitutively active mutant protein that has apparently lost its ability to respond to GA (Peng et al., 1997). Additionally, extragenic suppressors of GA mutants, defined by *rga* mutants, have been identified (Silverstone et al., 1997). *RGA* encodes a protein similar to that of the *GAI* gene (Silverstone et al., 1998). The *gai* mutant allele that results in the GA-insensitive dwarf mutation contains a 51-bp in-frame deletion of 17 amino acids in the DELLA domain (Peng et al., 1997). Peng et al. (1999) examined GA-insensitive dwarf mutants from other species and demonstrated that the mutations of reduced height (*RHT-B1* and *RHT-D1* in wheat and *D8* in maize) are caused by an N-terminal truncation near the DELLA domain. These results indicate that the DELLA domain is important to GA detection in the GA signal transduction pathway.

In wheat, some dwarf plants differ from tall plants in their lack of GA response. These dwarf mutants have a lodging-resistant characteristic and have resulted in increased wheat grain yields around the world since the 1960s. These so-called "green revolution" dwarf cultivars, which are derived from the Japanese variety Norin 10, have a phenotype with a reduced response to GA that is caused by mutations in the (*Rht-B1* and *Rht-D1*) gene, which is the ortholog of *GAI* (Peng et al., 1999).

In this study, we isolated and characterized the *slender*-type mutant in rice (*slr1-1*). Biochemical analysis revealed that the mutation results in a constitutive GA response. Consequently, we demonstrated that the *slr1-1* mutant contains a loss-of-function mutation in the *SLR1* gene, which is an ortholog of *GAI* and *RGA* in Arabidopsis, *RHT* in wheat, and *D8* in maize.

## RESULTS

### Isolation of *slender* Rice

A mutant with greatly accelerated extension growth was isolated from the rice cv Nipponbare after treatment with  $\gamma$  irra-

diation. The phenotype of the mutant, designated *slender* (*slr1-1*), has very rapid extension growth in the seedling and is sterile. M2 progeny tests of heterozygotes yielded a segregation of 311 normal and 89 *slender* plants ( $\chi^2$  [3:1] = 1.47;  $0.1 < P < 0.25$ ), indicating monofactorial recessive inheritance of the *slender* characteristic.

### *slender* Rice Behaves as if It Were Continually Saturated with GAs

One of the best known actions of GA is the stimulation of shoot growth in rice (Murakami, 1968; Matsukura et al., 1998). Germinating seeds of progeny from the heterozygote segregated a shoot length phenotype (Figure 1A). Shoots of the *slender* phenotype were more than twofold longer than those of the wild-type plant and were similar to those of wild-type plants treated with gibberellic acid ( $GA_3$ ; Figure 1B). Basal internodes in *slender* rice elongate concurrently during seedling growth, which is usually observed in the wild-type plant treated with  $GA_3$  (Figure 1C). The mutant also had a reduced number and root length compared with the wild-type plant (Figure 1D). Matsukura et al. (1998) reported that the promotion of leaf sheath growth by  $GA_3$  is due mainly to cell elongation. The average longitudinal and transverse lengths of the epidermal cells at the apical portion of the second leaf sheath were approximately twofold and 0.5-fold, respectively, the lengths of epidermal cells in the wild type (Table 1). The overall results indicate that *slender* rice behaves as if it were continually saturated with GAs.


The anatomic characteristics of *slender* rice are similar to those previously reported for the *ao-1* mutant in rice (Nakamura, 1992) and the *slender* mutant in barley (Favret and Malvarez, 1975; Foster, 1977).

*slender* and wild-type plants were grown on agar plates with or without uniconazole, a GA biosynthesis inhibitor (Izumi et al., 1984; Mitsunaga and Yamaguchi, 1993), for up to 15 days. The height of wild-type plants was decreased significantly with uniconazole throughout the experimental period, but the height of *slender* plants was unaffected (Figure 2). This observation suggests that the elongation of *slender* is independent of endogenous GA levels.

**Table 1.** Characteristics of Epidermal Cells in the Second Leaf Sheath of *slender* Rice and its Wild Type<sup>a</sup>

Position	Plant	Cell Length ( $\mu\text{m}$ )	Cell Width ( $\mu\text{m}$ )	Length/Width
Basal	Wild type	69.3 $\pm$ 11.8	19.7 $\pm$ 1.9	3.5
	<i>slender</i>	88.2 $\pm$ 5.8	15.8 $\pm$ 1.3	5.7
Apical	Wild type	55.9 $\pm$ 11.1	17.5 $\pm$ 2.4	3.2
	<i>slender</i>	107.9 $\pm$ 10.3	9.2 $\pm$ 0.7	11.7

<sup>a</sup>The second leaf sheath was dissected from 1-week-old *slender* and wild-type plants and stained with Safranin. Values are means  $\pm$  SE ( $n = 20$ ).



**Figure 2.** Effect of Uniconazole on Shoot Elongation in *slender* Rice and Its Wild Type.

**(A)** Seed of wild-type (1 and 2) and *slender* rice (3 and 4) were germinated for 6 days with (2 and 4) or without (1 and 3) 6.9  $\mu$ M uniconazole.

**(B)** Shoots of wild-type and *slender* rice were measured from 3 to 15 days after germination. Results from wild-type (open squares) and *slender* rice (open circles) without uniconazole and wild-type (closed

### *slender* Mutation Results in Modulation of Endogenous GAs and $\alpha$ -Amylase Production


Endogenous levels of GAs were estimated using a combination of HPLC purification and gas chromatography–mass spectrometry analysis (Figure 3). The level of GA<sub>1</sub>, the active GA molecule in vegetative tissues of rice, was lower in the shoots of *slender* rice than in the wild type. Similarly, the levels of GA<sub>19</sub> and GA<sub>20</sub>, both of which are inactive precursors of the GA<sub>1</sub> molecule, were lower in *slender* than in the wild type (Figure 3), indicating that the mutation decreases the endogenous GA level in the shoot.

For further characterization of *slender* rice, agar plate assays for amylases were conducted using the progeny seeds from selfing a plant heterozygous for the *slender* allele. The embryoless half-seeds were placed on the starch plate with or without 1  $\mu$ M GA<sub>3</sub> for 2 days, and the starch was stained with iodine (Figure 4A). Production and secretion of  $\alpha$ -amylase from wild-type embryoless half-seeds were observed as cleared zones (plaques) only on the plate containing GA<sub>3</sub> (Figure 4A, +GA<sub>3</sub>). Some *slender* half-seeds, however, produced amylase even in the absence of exogenous GA<sub>3</sub> (Figure 4A, –GA<sub>3</sub>). The ratio of the seeds requiring GA<sub>3</sub> for amylase secretion to seeds requiring no GA<sub>3</sub> was less than 3:1 (30:7 actual values). This suggests that the M2 progeny were segregated into *slender* and normal characteristics for  $\alpha$ -amylase production, because the aleurone layer consists of triploid tissues. These results demonstrate that exogenous GA<sub>3</sub> is not necessary for the synthesis and secretion of amylases from the embryoless half-seeds of *slender* rice.

We confirmed that the secreted amylase was GA-inducible  $\alpha$ -amylase (Figures 4B and 4C). Immunoblotting using antisera against rice  $\alpha$ -amylase (isoform A) revealed that the amylases secreted from the embryoless half-seeds of the *slender* mutant without GA<sub>3</sub> application are identical to those of wild-type seeds with GA<sub>3</sub> application (Figure 4B, lanes 2 and 3). Isoform analysis using an isoelectric focusing gel revealed that the main amylase is an isoform A, the gene product of *RAmy1A* (Figure 4C; Mitsui et al., 1996; Yamaguchi, 1998). The transcript of the rice  $\alpha$ -amylase gene *RAmy1A* is abundant in aleurone layer cells during germination and it is promoted by GA<sub>3</sub> (Itoh et al., 1995; Sugimoto et al., 1998). These results indicate that  $\alpha$ -amylase is produced independently of GAs from the aleurone layer cells of *slender* rice.

These findings (GA-mediated phenomena, shoot growth, and  $\alpha$ -amylase production) suggest that *slender* rice (*slr1-1*) is a constitutive GA response mutant. A similar conclusion was reported for the barley *slender* mutant (Chandler, 1988; Lanahan and Ho, 1988). Based on these results, it is likely

squares) and *slender* rice (closed circles) with 6.9  $\mu$ M uniconazole are shown. Data are means ( $n = 10$ ). Variation in width was <10% of the reported data.



**Figure 3.** GA Content of Shoots in Wild-Type and *slender* Rice.

Ten-day-old shoots were measured for GA content by gas chromatography–mass spectrometry analysis. Ten-day-old shoots were harvested from wild-type and *slender* rice and the GA content was determined. Samples were performed on 10 independent plants. FW, fresh weight. Bars indicate  $\pm$ SE.


that the *SLR1* gene is an intermediate of the GA signaling pathway.

### Molecular Cloning of the *SLR1* Gene

Nipponbare (a Japonica rice) containing *slr1-1* was crossed to Kasalath (an Indica rice), and segregation of *slr1-1* and *OsGAI* (Ogawa et al., 2000) was examined to determine if *SLR1* and *GAI* are linked. *GAI* in Arabidopsis is an intermediate of the GA signaling pathway. Direct sequencing was performed to determine the difference of the *GAI* homolog between Nipponbare and Kasalath. Base differences at position 1025 from the translation initiation site and their phenotypes were as follows: A, Nipponbare with a slender phenotype; G, Kasalath with a normal phenotype; and a mixture of A and G in the heterozygous plant having a normal phenotype. F<sub>2</sub> progeny tests of the heterozygotes yielded a segregation of 49 normal and 16 *slender* plants, and genotypes based on the base change indicated 16 Nipponbare homozygous (Japonica), 33 heterozygous (Japonica plus Indica), and 16 Kasalath homozygous (Indica) plants. These results indicate tight linkage between the *slender* mutation (*slr1-1*) and *OsGAI*.

We determined the open reading frame (ORF) sequence of the rice *GAI* homolog (data not shown). This sequence


was identical to that of *OsGAI* reported previously (Ogawa et al., 2000). We propose that *OsGAI* be renamed *SLR1* after its mutant phenotype and so use the term *SLR1* in place of *OsGAI* throughout this article. The deduced *SLR1* protein shares a high overall identity with RHT-D1a in wheat (77.2%), D8 in maize (80.3%), and RGA and *GAI* in Arabidopsis (41.2 and 47.2%, respectively). This protein contains



**Figure 4.**  $\alpha$ -Amylase Production from Embryoless Half-Seeds of Wild-Type and *slender* Rice.

(A) Embryoless half-seeds of wild-type and *slender* rice were placed on starch plates containing 1  $\mu$ M GA<sub>3</sub> (+GA<sub>3</sub>) or no GA<sub>3</sub> (–GA<sub>3</sub>) for 2 days, and starch was detected by staining with iodine.

(B) and (C) Immunoblot and zymogram pattern of  $\alpha$ -amylase expressed in embryoless half-seeds of wild-type and *slender* rice. (B) shows immunochromatological detection of  $\alpha$ -amylase protein using  $\alpha$ -amylase antibody. (C) shows zymogram pattern of  $\alpha$ -amylase activity. Lanes 1, wild type without GA<sub>3</sub> application; lanes 2, wild type with GA<sub>3</sub> application; lanes 3, *slender* rice without GA<sub>3</sub> application. Embryoless half-seeds of *slender* rice secreted isoform A (molecular mass  $\sim$ 43 kD; pI 4.5), which is a GA<sub>3</sub>-inducible  $\alpha$ -amylase in rice (Yamaguchi, 1998).



**Figure 5.** The Mutation Positions of the *slr1* Alleles.

Scheme of the deduced amino acid sequence for SLR1. Arrows indicate mutation of the *slr1* alleles in the position of the SLR1 ORF.

the DELLA motif (see Figure 5), which is believed to be a GA response domain, a putative nuclear localization signal (NLS motif), the VHIID domain, and the LXXLL motif.

#### Molecular Analysis of the *slender* Mutant Alleles and Mapping of the *SLR1* Gene


Three tall rice mutants (*slr1-2*, *slr1-3*, and *slr1-4*) were determined to be alleles of *slr1-1*. To identify the molecular basis of these mutations, the DNA sequences of the four mutant alleles were determined and compared with those of the *SLR1* gene. The *slr1-1* allele contained one base deletion at Leu<sup>289</sup>, a putative NLS region, that alters the N-terminal region of the protein that it encodes (Figure 5). The other three alleles (*slr1-2*, *slr1-3*, and *slr1-4*) contained premature stop codons (Figure 5). These results indicate that the *SLR1* gene is associated with the slender phenotype.

We mapped the *SLR1* gene on the rice genome using restriction fragment length polymorphism analysis. Linkage analysis was performed with digested genomic DNA from recombinant inbred lines of crosses between Asominori and IR24. Consequently, chromosome 3 contained a single *SLR1* gene (Figure 6). From the partial linkage maps of wheat and rice chromosomes, the rice *GAI* homolog was predicted to be on chromosome 3 (Peng et al., 1999). This result is consistent with the *SRL1* gene being a *GAI* homolog.

#### Molecular Complementation of the *slender* Mutant by the Wild-Type *SLR1* Gene

To confirm that the *SLR1* gene corresponds to the *slender* mutation, we performed complementation analysis of the *slender* mutant using the wild-type *SLR1* gene. Transformation of the *slender* mutant with a control vector that carried

no rice genomic DNA had no apparent effect on the phenotype (data not shown). When a 6-kb DNA fragment containing the entire wild-type *SLR1* gene was introduced, however, the normal phenotype was restored in most plants that were resistant to hygromycin (Figure 7, NT). GA<sub>3</sub> and uniconazole were applied to the transformants to determine if normal sensitivity to the GA response was restored. The application of GA<sub>3</sub> increased and uniconazole decreased



**Figure 6.** Mapping of the *SLR1* Gene on Rice Chromosome 3.

The map position of the *SLR1* gene is shown relative to its physical flanking markers C1351 and R19. The numbers at left are the distance of the markers (in centimorgans), and the number in parentheses denotes the position of the *SLR1* gene according to the 71 recombinant inbred lines from a cross between rice cultivars Asominori (a Japonica rice) and IR24 (an Indica rice).

plant height (Figure 7, +GA<sub>3</sub> and +Unico), indicating that the introduction of the wild-type *SLR1* gene complemented the *slender* mutation. Based on these results, we concluded that the *slender* mutant is caused by a loss-of-function mutation of the *SLR1* gene, which is identical to the rice *GAI* homolog.

#### Truncation of DELLA, a Putative GA Response Motif in the *SLR1* Gene, Leads to the Dwarf Phenotype

In *Arabidopsis*, the GA-insensitive dwarf mutation allele *gai* contains a 51-bp deletion of the *GAI* ORF. This in-frame deletion results in the absence of 17 amino acid residues in the DELLA motif of the *GAI* protein. We attempted to produce GA-insensitive dwarf rice using the truncated *SLR1* gene construct *pSLRtr*, which also contains a 17-amino acid deletion (39-DELLAALGYKVRSSDMA-55) in the DELLA motif (Figure 8A). When the *pSLRtr* construct was introduced into the normal Japonica rice (cv Nipponbare) to produce ~200 plants, >90% of transgenic rice resulted in a typical dwarf plant (Figure 8B, *pSLRtr*). The transgenic plant had retarded growth in the leaf sheath and blade and thickening in the roots, which are typical responses for a GA-deficient plant. The dwarf transgenic plant had no response, however, to exogenously applied GA<sub>3</sub>. Introduction of the truncated *SLR1* gene led to a wide range of plant heights, from severe, extremely dwarf phenotypes to mild phenotypes (Figure 8C).

## DISCUSSION

### *slender* (*slr1-1*) Is a Constitutive GA Response Mutant in Rice

In this study, *slender* rice, which is caused by a single recessive mutation, was considered to be a constitutive GA response phenotype for the following reasons. The mutant shoot was more than twofold taller than the wild type, similar to wild-type shoots treated with high amounts of GA<sub>3</sub>. The height of *slender* was unaffected by the application of uniconazole, an inhibitor of GA biosynthesis, and endogenous levels of GAs were lower in *slender* than in the wild type. Finally, GA-inducible  $\alpha$ -amylase, the *RAmy1A* protein, was produced without GA<sub>3</sub> application.

There are similar recessive *slender*-type mutants in barley (*slender* [*sln1*]; Chandler, 1988; Lanahan and Ho, 1988) and in pea (*la crys*; Potts et al., 1985). These mutants behave as if saturated with GAs and are not responsive to either exogenous GAs or GA biosynthesis inhibitors. On the other hand, the *SPY* mutant in *Arabidopsis* (Jacobsen and Olszewski, 1993) and the *pro* mutant in tomato (Jones, 1987) respond to applied GAs. These two types of mutants might be due to the mutation of genes that have different functions or that




**Figure 7.** Complementation Analysis of the Slender Phenotype.

A 6-kb wild-type genomic DNA fragment containing the entire *SLR1* gene was used for transformation of the *slender* mutant (*slr1-1*). The transgenic plant had a normal phenotype without hormone treatment (NT). Application of 10  $\mu$ M GA<sub>3</sub> (+GA<sub>3</sub>) led to increased height, and treatment with 6.9  $\mu$ M uniconazole (+Unico) led to reduced height.

encode different intermediates in the GA signal transduction pathway.

Embryoleless half-seeds of *slender* produced GA-inducible  $\alpha$ -amylase without GA<sub>3</sub>, indicating that the *SLR1* protein might be a negative regulator that blocks transcription of the  $\alpha$ -amylase gene (see below). The induction of  $\alpha$ -amylase by GAs in cereal grains represents a classic model system for studying the mode of action of GAs (Jones and Jacobsen, 1991).  $\alpha$ -Amylase induction by GA occurs mainly at the transcriptional steps, which might be mediated through abscisic acid, Ca<sup>2+</sup> (Schuurink et al., 1996), cyclic GMP (Penson et al., 1996), or sugar signaling (Perata et al., 1997), and signals communication among them (Toyofuku et al., 2000). GA-dependent transcriptional activation of the  $\alpha$ -amylase gene requires the GA response *cis*-element complex and the *GAMYb* protein as a *trans*-acting factor (Gubler et al., 1995). Further study is needed to clarify the targeting site of the *SLR1* protein through the GA signaling pathway.



**Figure 8.** Truncation of the DELLA Motif in *SLR1* Leads to a Dwarf Phenotype.

### The *SLR1* Gene Is an Ortholog of the Height-Regulating Gene *GAI/RGA/RHT/D8*

We succeeded in the molecular cloning of the *SLENDER* gene (*SLR1*), which was originally derived from the rice *GAI* homolog. Molecular analysis of *slender* mutant alleles revealed that four mutant alleles contain nonsense or stop codon mutations in the *SLR1* gene. Introduction of a 6-kb genomic DNA fragment for the wild-type *SLR1* gene into the *slr1-1* mutant restored the normal phenotype and sensitivity to the GA response.

*GAI*, *SLR1*, *D8*, *RHT-1Da*, and *RGA* seem to be transcriptional factors belonging to the GRAS gene superfamily, which has a putative NLS, an Ser/Thr-rich repeat region, and the LXXLL motif. Indeed, in a transient assay using onion epidermal cells, *RGA-GFP* (green fluorescent protein) and *OsGAI-GFP* fusion proteins were localized exclusively in the nucleus (Silverstone et al., 1998; Ogawa et al., 2000). Transactivation assays indicate that the *OsGAI* protein has a transcriptional activator or coactivator (Ogawa et al., 2000).

Sequence comparison among members of the *GAI* family identified a putative Src homology 2 (SH2) phosphotyrosine binding domain (Peng et al., 1999). The SH2 domain is present in a family of transcription factors called STATs (signal transducers and activators of transcription) in animals (Darnell, 1997). The function of this domain is to mediate the binding of STATs to various receptor tyrosine kinases. The STATs are then activated by the receptor kinase and translocate from the cytoplasm to the nucleus. If this putative SH2 domain is functional in plants, then tyrosine phosphorylation might have a role in GA signaling (Silverstone and Sun, 2000).

The *slr1-4* mutation was detected at the extreme C-terminal end (Trp<sup>620</sup>) of the *SLR1* protein (deduced ORF of 625 amino acids). This mutant did not have any phenotypic differences from the other alleles. These results indicate that even the loss of six amino acid residues at the C-terminal end, which are not a conserved motif among family members, abolishes the protein function in spite of the presence of consensus motifs NLS, LXXLL, and SH2.

**(A)** The construct *pSLRtr* was used for the transformation of rice plants. The *Actin1* promoter (*pAct1*) was recovered from vector pBI101-Hm3. Transformation with the *SLRtr* sequence resulted in the DELLA motif truncation form (39-DELLAALGYKVRSSDMA-55) of the *SLR1* ORF. *nos*, nopaline synthase.


**(B)** Construct *pSLRtr* **(A)** was introduced into the normal Japonica rice Nipponbare (Control). Introduction of the truncated *SLR1* gene resulted in the typical dwarf plant (*pSLRtr*). The transgenic plant shows retarded growth in the leaf sheath and blade.

**(C)** Introduction of the truncated *SLR1* gene resulted in a range of plant heights. Construct *pSLRtr* **(A)** was introduced into the wild-type cv Nipponbare (Control). The *pSLRtr* transgenic plants show a wide range of plant heights, from weak to severe dwarf phenotypes.

### SLR1 Is a Negative Regulator Involved in the GA Signaling Pathway

The *slr1-1*, *slr1-2*, and *slr1-3* mutations truncate the protein, suggesting that these mutations are loss-of-function mutations. Therefore, SLR1 protein must be a negative regulator of the GA signaling process, because lack of its negative regulation results in a constitutive GA response phenotype (Figure 9).

Consistent with the similarity of the phenotypes of *Rht-B1b*, *Rht-D1b*, and *D8* to the semidominant *gai* mutant, the genetic lesions in these mutants are also similar to those in *gai* (Peng et al., 1999). The lesions in three *D8* mutant alleles have been identified. Both *D8-1* and *D8-2023* contain small internal deletions in the DELLA domain. The *D8-Mpl* allele has a 330-bp deletion at the N terminus that begins in the 59-untranslated amino acid region and extends through Val<sup>84</sup>. Because of the gain-of-function (semidominant) nature of this mutant, it is proposed that it still encodes a truncated D8 protein that might be initiated at Met<sup>106</sup> (Peng et al., 1999). *Rht-B1b* and *Rht-D1b* both contain a base substitution that introduces a stop codon in the DELLA domain. Because there is a methionine a short distance after these stop codons, it is possible that translational reinitiation occurs to synthesize a protein lacking the original N terminus (Peng et al., 1999). These results support the hypothesis that the DELLA domain in the N terminus of the GAI protein family is responsible for modulating the activity of these proteins in response to the GA signal. Presumably, the effect of either the internal deletions or the N-terminal truncations is to lock the protein into a conformation that can no longer, or can only weakly, respond to the GA signal (Silverstone and Sun, 2000).



**Figure 9.** Model of the GA Signal Transduction Pathway in Rice, Arabidopsis, and Wheat.

Arrows indicate the GA signal transduction pathway from GA reception (GA) to various actions of GA, such as shoot elongation,  $\alpha$ -amylase induction, and so on. SLR1 in rice, GAI and RGA in Arabidopsis, and RHT-A1, -B1, and -D1 proteins in wheat negatively regulate the GA signaling pathway.

### The Distinct Manner of Mutation on the GAI Ortholog Leads to Opposite Phenotypes in Terms of GA Sensitivity

In Arabidopsis, two orthologous proteins, GAI and RGA, have a redundant GA response; therefore, even if one of the proteins is inactive, the plant still has a normal phenotype (Figure 9). In this case, a dominant negative mutation such as the DELLA truncation is the only way to produce an abnormal phenotype, such as the GA-insensitive dwarf. There is a similar redundancy in wheat, with its gene duplication, because wheat possesses three sets of chromosomes (A, B, and D genomes).

In rice, the *slender* mutant results in a loss-of-function mutation of the *SLR1* gene (Figure 9). Because the *SLR1* gene is likely to be nonredundant, the loss-of-function mutation results in clearly abnormal phenotypes in terms of the GA response, that is, the slender and constitutive GA response phenotype. We demonstrated that the distinct manner of mutation on the orthologous genes leads to opposite phenotypes in terms of the GA response: slender, a constitutive GA response, and a GA-insensitive dwarf.

### Dwarfism in Rice and the Alternative Green Revolution

World wheat grain yields increased substantially in the 1960s and 1970s because farmers rapidly adopted the new varieties and cultivation methods of the so-called green revolution. The new varieties are shorter (dwarf or semidwarf) and increase grain yield by means of lodging resistance. These wheat cultivars are short because they contain the DELLA-truncated *RHT* gene (Peng et al., 1999).

With rice, a similar green revolution was realized by the introduction of the distinct gene *sd-1* from *GAI/RHT*. IR8 was established by crossing *Dee-geo-woo-gen* (*sd-1*) from a Chinese cultivar to *Peta* from an Indonesian cultivar at the International Rice Research Institute (Los Baños, Philippines; Hargrove et al., 1980). In Japan, the semidwarf cultivar *Reimei* (*d-49*; allelic to *sd-1*), which was derived from cv *Fujiminori* through  $\gamma$  irradiation, was released in 1966 (Futsuhara et al., 1967). *Reimei* was grown over wide areas of the northern part of Japan. Furthermore, it has been used as a parent strain for cross-breeding, taking advantage of its lodging resistance and high yields. We demonstrated that the introduction of the truncated *SLR1* gene results in a range of plant heights (Figure 8C). DNA gel blot analysis revealed that these plants possess a single gene introduction and that the diversity in height is due mainly to the positional effect of the transgene insertion (data not shown), suggesting the potential for a plant height-regulating system. Because the *SLR1* gene is distinct from *sd-1* (the rice green revolution gene), it presents an alternative means to produce dwarf rice plants. Further investigation is needed to evaluate this hypothesis.

## METHODS

### Isolation of *slender* Rice

A recessive mutant, *slender1-1*, was isolated from the progeny of rice seeds (*Oryza sativa* cv Nipponbare, a Japonica rice) irradiated with 200 Gy of  $\gamma$ -rays (dose rate 2 Gy/min). Surface-sterilized seeds of wild-type and mutant plants were soaked in water for 3 days and then placed in artificial soil for 11 days and grown in a greenhouse. Mutant segregants were distinguished from normal segregants by the extremely extended phenotype. The other *slender* alleles (*slr1-2*, *1-3*, and *1-4*) were independently isolated.

### Measurement of Shoot Elongation

Shoot elongation was quantified by a modification of the method described by Matsukura et al. (1998). Seeds of wild-type and mutant plants ( $n = 25$ ) were surface sterilized for 30 min with a 3% NaClO solution, washed three times with sterile distilled water, soaked in the distilled water for 24 hr in the presence or absence of 6.9  $\mu$ M uniconazole, and then placed in sterile distilled water for an additional 24 hr. The seeds were then placed on a 1% agar plate and grown under fluorescent lamps at 30°C. For the surface anatomy, the second leaf sheath was dissected from third leaf-stage plants and stained with Safranin (Wako Chemical, Osaka, Japan).

### Agar Plate Assay, Immunoblotting, and Zymography of $\alpha$ -Amylase

The agar plate assay was performed essentially as described by Lanahan and Ho (1988). Seeds were cut transversely, and the half-seed containing the embryos were planted to determine their phenotypes. The embryoless half-seeds were surface sterilized with 3% NaClO for 15 min and washed six times with sterile water. These half-seeds were then placed on 2% agar plates containing 10 mM sodium acetate and 2 mM CaCl<sub>2</sub> at pH 5.3. GA plates were made by adding 1  $\mu$ M gibberellic acid (GA<sub>3</sub>) to the cooled agar. To detect secreted  $\alpha$ -amylase activity, we added soluble potato starch (0.2%) to the agar before autoclaving. Agar plates were developed by incubating the plates in I<sub>2</sub> gas. Half-seeds that synthesized and secreted  $\alpha$ -amylase had transparent halos around them resulting from the digestion of the starch by amylases.

Proteins to be examined were separated using SDS-PAGE or isoelectric focusing, transferred to a nitrocellulose membrane using the Novablot Protein Transfer Kit, and analyzed using the anti- $\alpha$ -amylase antibody (Mitsunaga and Yamaguchi, 1993). An alkaline phosphatase-labeled secondary antibody was used to detect the immunoreactive band.

The crude extract was examined by isoelectric focusing using Pharmacia broad range (pH 3.5 to 9.5) Ampholine Pageplates. Isoelectric focusing was performed for 1.5 hr according to the manufacturer's instructions. Samples (15  $\mu$ L of crude extract) were applied to application paper placed 3 cm distant from the cathode. The application paper was removed after 45 min of electrophoresis, and the run was continued for an additional 45 min. To visualize the bands of amylolytic activities, the gel was incubated for 1 hr in 50 mM sodium acetate buffer, pH 5.2, containing 10 mM CaCl<sub>2</sub> and 1% boiled solu-

ble starch. After washing with distilled water, the gel was stained with 0.6% I<sub>2</sub> and 6% KI solution (Perata et al., 1992).

### Quantitative Analyses of Endogenous GAs

GAs from each sample were extracted and purified according to the method of Hisamatsu et al. (1998). As internal standards, 70 ng of <sup>17,17,2</sup>H<sub>2</sub>-GA<sub>1</sub>, 100 ng of <sup>17,17,2</sup>H<sub>2</sub>-GA<sub>19</sub>, and 50 ng of <sup>17,17,2</sup>H<sub>2</sub>-GA<sub>20</sub> were added to each sample. After several HPLC purification steps, fractions that had GA activity on a rice assay (Nishijima et al., 1992) were methylated and trimethylsilylated in glass tubes. The derived samples were analyzed using gas chromatography-mass spectrometry. The concentrations of GA<sub>1</sub>, GA<sub>19</sub>, and GA<sub>20</sub> in the tubes were calculated from the ratios of peak areas at mass-to-charge ratios of 506:508, 434:436, and 418:420, respectively, according to the method of Gaskin and MacMillan (1991).

### Screening of the *SLR1* Gene from Rice Genomic Libraries

To isolate the *SLR1* gene, we used the rice *GAI* homologous expressed sequence tag clone (accession number D39460). Genomic DNA libraries were constructed from rice. Nuclear genomic DNA was isolated from 2-week-old seedlings. The DNA was partially digested with Sau3AI and enriched for fragments of ~20 kb on a sucrose gradient. The fragments were cloned into the BamHI site of EMBL3 (Stratagene, La Jolla, CA).

Screening by hybridization was performed in 50% formamide, 6  $\times$  SSC (1  $\times$  SSC is 0.15 M NaCl and 0.015 M sodium citrate), 5  $\times$  Denhardt's solution (1  $\times$  Denhardt's solution is 0.02% Ficoll, 0.02% polyvinylpyrrolidone, and 0.02% BSA), 0.5% SDS, and 0.1 mg/mL salmon sperm DNA at 42°C for 14 hr using the *GAI* homologous expressed sequence tag clone as a probe.

### Mapping of the *SLR1* Gene in Rice Recombinant Inbred Lines

To map the *SLR1* gene, 71 recombinant inbred lines from a cross between two rice cultivars, Asominori (a Japonica rice) and IR24 (an Indica rice), were used. Restriction fragment length polymorphism analysis was performed with a probe specific for the rice *GAI* homologous expressed sequence tag clone. DNA gel blot hybridization was performed as described by Church and Gilbert (1984) except that membranes were hybridized at high stringency (68°C). The linkage analysis was performed using the MAPMAKER program (Whitehead Institute for Biomedical Research/Massachusetts Institute of Technology Center for Genome Research, Wilmington, MA).

### Genomic DNA Isolation and Direct Genome Sequencing

Rice genomic DNA was isolated with the ISOPLANT DNA isolation kit (Nippon Gene Co., Tokyo, Japan). The coding region of the *SLR1* gene was amplified by polymerase chain reaction (PCR) using specific primers S6 (5'-TCGTCGCTCCTCATCGTCGTC-3') and A6 (5'-GCAGCGGTGCGACTCGAAC-3') from 300 ng of rice genomic DNA. PCR products were gel purified, and the nucleotide sequences were determined using the dideoxynucleotide chain termination method with an automated sequencing system (ABI373A; Applied Biosystem, Inc., Foster City, CA). Analysis of the nucleotide sequences was

performed using DANSIS computer software (Hitachi Software Engineering, Tokyo, Japan).

#### Rice Transformation and Complementation of the *slr1-1* Mutant

A genomic clone, *pAI1*, including the entire coding region and the 5' and 3' flanking regions, was inserted as a 6-kb restriction fragment between the SmaI sites of the hygromycin resistance binary vector pBI101-Hm3 (Sato et al., 1999). The *pAI1* gene construct was introduced into the *slr1-1* mutant as described above. Control plants were transformed using the vector *pBI-cont*.

*pSLR1tr* was made as follows. A 123-bp fragment was amplified by PCR from 6 kb of genomic DNA containing the SLR ORF using a 5' primer containing an additional XbaI site (5'-TCTAGAATGAAGCGC-GAGTA-3') and a 3' primer containing an additional Sall site (5'-GTCGACGTCTCTCTCTCC-3'). A 1722-bp fragment was amplified by PCR from 6 kb of genomic DNA containing the SLR ORF using a 5' primer containing an additional Sall site (5'-GTCGAC-GTCGCGCAGAAGCT-3') and a 3' primer containing an additional SmaI site (5'-CCCGGGTCACGCCGCGGCGA-3'). These PCR products were cloned into a *pBIAct1nos* vector (Yamamuro et al., 2000). *pBIAct1nos* containing the actin (*Act1*) promoter (Zhang et al., 1991) and the nopaline synthase terminator was prepared by substituting the *Act1* promoter in the hygromycin resistance binary vector between the XbaI and SmaI sites. *pBI-cont*, which contains no insert, was used as a control vector. *pSLR1tr* and the control construct were introduced into rice cv Nipponbare by *Agrobacterium tumefaciens*-mediated transformation, as described by Hiei et al. (1994).

#### ACKNOWLEDGMENTS

We are grateful to Dr. Pierdomenico Perata for critically reviewing the manuscript and Professor Kenzo Nakamura (Nagoya University) for providing us with the pBI101-Hm3 vector. This work was supported by Grants-in-Aid for Scientific Research (Grant Nos. 09660004 and 10460003) from the Ministry of Education, Science, Sports, and Culture, Japan. A.I. acknowledges a Research Fellowship of the Japan Society for the Promotion of Science for Young Scientists (2000–2001). We thank Dr. T. Sasaki of the Japanese Rice Genome Program for the expressed sequence tag clones.

Received October 23, 2000; accepted March 7, 2001.

#### REFERENCES

- Chandler, P.M.** (1988). Hormonal regulation of gene expression in the "slender" mutant of barley (*Hordeum vulgare* L.). *Planta* **175**, 115–120.
- Church, G., and Gilbert, W.** (1984). Genomic sequencing. *Proc. Natl. Acad. Sci. USA* **81**, 1991–1995.
- Crocker, S.J., Hedden, P., Lenton, J.R., and Stoddart, J.L.** (1990). Comparison of gibberellins in normal and slender barley seedlings. *Plant Physiol.* **94**, 194–200.
- Darnell, J.E., Jr.** (1997). STATs and gene regulation. *Science* **277**, 1630–1635.
- Favret, E.A., and Malvarez, E.M.** (1975). Genetic regulatory mechanisms for seedling growth in barley. *Barley Genet.* **3**, 37–43.
- Foster, C.A.** (1977). Slender: An accelerated extension growth mutant of barley. *Barley Genet. Newsl.* **7**, 24–27.
- Futsuhara, Y., Toriyama, K., and Tsunoda, K.** (1967). Breeding of the new rice variety, "Reimei," through gamma-ray irradiation. *Jpn. J. Breed.* **17**, 85–90.
- Gaskin, P., and MacMillan, J.** (1991). Quantitative analysis. In *GC-MS of the Gibberellins and Related Compounds: Methodology and a Library of Spectra*, P. Gaskin and J. MacMillan, eds (Bristol, UK: Cantock's Enterprises), pp. 117–124.
- Gubler, F., Kalla, R., Roberts, J.K., and Jacobsen, J.V.** (1995). Gibberellin-regulated expression of a *myb* gene in barley aleurone cells: Evidence for Myb transactivation of a high-pI  $\alpha$ -amylase gene promoter. *Plant Cell* **7**, 1879–1891.
- Hargrove, T.R., Coffman, W.R., and Cabanilla, V.L.** (1980). Ancestry of improved cultivars of Asian rice. *Crop Sci.* **20**, 721–727.
- Hiei, Y., Ohta, S., Komari, T., and Kumashiro, T.** (1994). Efficient transformation of rice (*Oryza sativa* L.) mediated by *Agrobacterium* and sequence analysis of the boundaries of the T-DNA. *Plant J.* **6**, 271–282.
- Hisamatsu, T., Koshioka, M., Kubota, S., King, R.W., and Mander, L.N.** (1998). Isolation and identification of GA12 (12 $\beta$ -hydroxy-GA12) in *Matthiola incana*. *Phytochemistry* **47**, 3–6.
- Hooley, R.** (1994). Gibberellins: Perception, transduction and responses. *Plant Mol. Biol.* **26**, 1529–1555.
- Itoh, K., Yamaguchi, J., Huang, N., Rodriguez, R.L., Akazawa, T., and Shimamoto, K.** (1995). Developmental and hormonal regulation of rice  $\alpha$ -amylase (*RAmy1A*)-*gusA* fusion genes in transgenic rice seeds. *Plant Physiol.* **107**, 25–31.
- Izumi, K., Yamaguchi, I., Wada, A., Oshio, H., and Takahashi, N.** (1984). Effects of a new plant growth retardant (E)-1-(4-chlorophenyl)-4,4-dimethyl-2-(1,2,4-triazol-1-yl)-1-penten-3-ol (S-3307) on the growth and gibberellin content of rice plants. *Plant Cell Physiol.* **25**, 611–617.
- Jacobsen, S.E., and Olszewski, N.E.** (1993). Mutations at the *SPINDLY* locus of *Arabidopsis* alter gibberellin signal transduction. *Plant Cell* **5**, 887–896.
- Jones, M.G.** (1987). Gibberellins and the *procera* mutant of tomato. *Planta* **172**, 280–284.
- Jones, R.L., and Jacobsen, J.V.** (1991). Regulation of synthesis and transport of secreted proteins in cereal aleurone. *Int. Rev. Cytol.* **126**, 49–88.
- Lanahan, M.B., and Ho, T.-H.D.** (1988). Slender barley: A constitutive gibberellin-response mutant. *Planta* **175**, 107–114.
- Matsukura, C., Itoh, S., Nemoto, K., Tanimoto, E., and Yamaguchi, J.** (1998). Promotion of leaf sheath growth by gibberellic acid in a dwarf mutant of rice. *Planta* **205**, 145–152.
- Mitsui, T., Yamaguchi, J., and Akazawa, T.** (1996). Physicochemical and serological characterization of rice  $\alpha$ -amylase isoforms and identification of their corresponding genes. *Plant Physiol.* **110**, 1395–1404.

- Mitsunaga, S., and Yamaguchi, J.** (1993). Production of  $\alpha$ -amylase is repressed by uniconazole, an inhibitor of the biosynthesis of gibberellin, in a dwarf mutant of rice, Waito-C. *Plant Cell Physiol.* **34**, 243–249.
- Mitsunaga, S., Tashiro, T., and Yamaguchi, J.** (1994). Identification and characterization of gibberellin-insensitive mutants selected from among dwarf mutants of rice. *Theor. Appl. Genet.* **87**, 705–712.
- Murakami, H.** (1968). A new rice seedling test for gibberellins, 'microdrop method' and its use for testing extracts of rice and morning glory. *Bot. Mag. Tokyo* **81**, 33–43.
- Nakamura, I.** (1992). A rice mutant showing accelerated internode overgrowth. *Rice Genet. Newsl.* **9**, 61–62.
- Nishijima, T., Koshioka, M., and Yamaji, H.** (1992). A non-dwarf rice seedling bioassay for gibberellins. *Plant Physiol.* **98**, 962–965.
- Ogawa, M., Kusano, T., Katsumi, M., and Sano, H.** (2000). Rice gibberellin-insensitive gene homolog, *OsGAI*, encodes a nuclear-localized protein capable of gene activation at the transcriptional level. *Gene* **245**, 21–29.
- Peng, J., and Harberd, N.P.** (1993). Derivative alleles of the *Arabidopsis* gibberellin-insensitive (*gai*) mutation confer a wild-type phenotype. *Plant Cell* **5**, 351–360.
- Peng, J., Carol, P., Richards, D.E., King, K.E., Cowling, R.J., Murphy, G.P., and Harberd, N.P.** (1997). The *Arabidopsis* *GAI* gene defines a signaling pathway that negatively regulates gibberellin responses. *Genes Dev.* **11**, 3194–3205.
- Peng, J., et al.** (1999). 'Green revolution' genes encode mutant gibberellin response modulators. *Nature* **400**, 256–261.
- Penson, S.P., Schuurink, R.C., Fath, A., Gubler, F., Jacobsen, J.V., and Jones, R.L.** (1996). cGMP is required for gibberellic acid-induced gene expression in barley aleurone. *Plant Cell* **8**, 2325–2333.
- Perata, P., Pozueta-Romero, J., Akazawa, T., and Yamaguchi, J.** (1992). Effect of anoxia on starch breakdown in rice and wheat seeds. *Planta* **188**, 611–618.
- Perata, P., Matsukura, C., Vernieri, P., and Yamaguchi, J.** (1997). Sugar repression of a gibberellin-dependent signaling pathway in barley embryos. *Plant Cell* **9**, 2197–2208.
- Potts, W.C., Reid, J.B., and Murfet, I.C.** (1985). Internode length in *Pisum*: Gibberellins and the slender phenotype. *Physiol. Plant.* **63**, 357–364.
- Sato, Y., Sentoku, N., Miura, Y., Hirochika, H., Kitano, H., and Matsuoka, M.** (1999). Loss of function mutations in the rice homeobox gene *OSH15* affect the architecture of internodes resulting in dwarf plants. *EMBO J.* **18**, 992–1002.
- Schuurink, R.C., Chan, P.V., and Jones, R.L.** (1996). Modulation of calmodulin mRNA and protein levels in barley aleurone. *Plant Physiol.* **111**, 371–380.
- Silverstone, A.L., and Sun, T.** (2000). Gibberellins and the green revolution. *Trends Plant Sci.* **5**, 1–2.
- Silverstone, A.L., Mak, P.Y.A., Casamitjana-Martinez, E., and Sun, T.-p.** (1997). The new *RGA* locus encodes a negative regulator of gibberellin response in *Arabidopsis thaliana*. *Genetics* **146**, 1087–1099.
- Silverstone, A.L., Ciampaglio, C.N., and Sun, T.-p.** (1998). The *Arabidopsis* *RGA* gene encodes a transcriptional regulator repressing the gibberellin signal transduction pathway. *Plant Cell* **10**, 155–159.
- Sugimoto, N., Takeda, G., Nagato, Y., and Yamaguchi, J.** (1998). Temporal and spatial expression pattern of  $\alpha$ -amylase gene during germination in rice and barley. *Plant Cell Physiol.* **39**, 323–333.
- Swain, S.M., and Olszewski, N.E.** (1996). Genetic analysis of gibberellin signal transduction. *Plant Physiol.* **112**, 11–17.
- Taylor, C.B.** (1998). GA signaling: Genes and GTPases. *Plant Cell* **10**, 131–133.
- Thornton, T.M., Swain, S.M., and Olszewski, N.E.** (1999). Gibberellin signal transduction presents . . . the SPY who O-GlcNAc'd me. *Trends Plant Sci.* **4**, 424–428.
- Toyofuku, K., Loreti, E., Vernieri, P., Alpi, A., Perata, P., and Yamaguchi, J.** (2000). Glucose modulates the abscisic acid-inducible *Rab16A* gene in cereal embryos. *Plant Mol. Biol.* **42**, 451–460.
- Yamaguchi, J.** (1998). Analysis of embryo-specific  $\alpha$ -amylase using isolated mature rice embryos. *Breeding Sci.* **48**, 365–370.
- Yamamoto, C., Ihara, Y., Wu, X., Noguchi, T., Fujioka, S., Takatsuto, S., Ashikari, M., Kitano, H., and Matsuoka, M.** (2000). Loss of function of a rice *brassinosteroid insensitive1* homolog prevents internode elongation and bending of the lamina joint. *Plant Cell* **12**, 1591–1606.
- Zhang, W., McElroy, D., and Wu, R.** (1991). Analysis of rice *Act1* 5' region activity in transgenic rice plants. *Plant Cell* **3**, 1155–1165.

***slender Rice, a Constitutive Gibberellin Response Mutant, Is Caused by a Null Mutation of the SLR1 Gene, an Ortholog of the Height-Regulating Gene GAI/RGA/RHT/D8***

Akira Ikeda, Miyako Ueguchi-Tanaka, Yutaka Sonoda, Hidemi Kitano, Masaji Koshioka, Yuzo Futsuhara, Makoto Matsuoka and Junji Yamaguchi

*Plant Cell* 2001;13:999-1010

DOI 10.1105/tpc.13.5.999

This information is current as of November 16, 2019

<b>References</b>	This article cites 45 articles, 19 of which can be accessed free at: <a href="/content/13/5/999.full.html#ref-list-1">/content/13/5/999.full.html#ref-list-1</a>
<b>Permissions</b>	<a href="https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&amp;issn=1532298X&amp;WT.mc_id=pd_hw1532298X">https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&amp;issn=1532298X&amp;WT.mc_id=pd_hw1532298X</a>
<b>eTOCs</b>	Sign up for eTOCs at: <a href="http://www.plantcell.org/cgi/alerts/ctmain">http://www.plantcell.org/cgi/alerts/ctmain</a>
<b>CiteTrack Alerts</b>	Sign up for CiteTrack Alerts at: <a href="http://www.plantcell.org/cgi/alerts/ctmain">http://www.plantcell.org/cgi/alerts/ctmain</a>
<b>Subscription Information</b>	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: <a href="http://www.aspb.org/publications/subscriptions.cfm">http://www.aspb.org/publications/subscriptions.cfm</a>