

The Molecular and Genetic Bases of S-RNase-Based Self-Incompatibility

Teh-hui Kao¹ and Tatsuya Tsukamoto

Department of Biochemistry and Molecular Biology, Pennsylvania State University, University Park, Pennsylvania 16802-4500

INTRODUCTION

The majority of flowering plants produce perfect flowers that contain both the male and female reproductive organs in close proximity; consequently, they would have a strong tendency to self-fertilize if there were no mechanisms to prevent them from doing so. Because inbreeding can result in reduced fitness in the progeny, hermaphroditic plants have adopted a variety of reproductive strategies, including self-incompatibility (SI), by which inbreeding is prevented and outcrosses are promoted. SI allows the pistil of a flower to distinguish between genetically related (self) and unrelated (non-self) pollen. This self/non-self recognition results in the inhibition of germination of self-pollen on the stigmatic surface or the inhibition of growth of self-pollen tubes in the style. Thus, SI is a prezygotic reproductive barrier by which incompatible pollen/pollen tubes are prevented from delivering the sperm cells to the ovary to effect double fertilization.

SI can be classified into homomorphic and heteromorphic types based on whether it is associated with floral polymorphism. In species that exhibit homomorphic SI, all individuals produce the same type of flower and the outcome of pollination depends only on the genetic identity of the male and female partners. In contrast, species that exhibit heteromorphic SI produce two or three different flower morphologies (e.g., a flower with short anthers and long style or a flower with long anthers and short style). For successful pollination, pollen must come from genetically unrelated individuals whose anthers are of the same height as the style of the flower being pollinated. To date, much of what we know about the molecular basis of SI has been deduced from studies of homomorphic SI, which will be the focus of this review. A monograph by de Nettancourt (2001) provides a comprehensive treatise on SI, including a discussion of the heteromorphic type.

For homomorphic SI (hereafter referred to as SI), self/non-self discrimination between pollen and pistil is determined by one or more polymorphic loci, and this type of SI is further classified into gametophytic and sporophytic types based on the genetic control of pollen behavior. To date, four of the families that exhibit gametophytic SI (GSI), Solanaceae, Rosaceae, Scrophulariaceae, and Papaveraceae, and one of the families that exhibit

sporophytic SI (SSI), Brassicaceae, have been studied extensively at the molecular level (Table 1). A single polymorphic locus, termed the *S*-locus, controls the SI response in all five of these families. As described below, other loci often are required for the full manifestation of the SI response, but by definition, the *S*-locus determines the specificity of the response. It is now known that two separate genes at the *S*-locus control male and female specificities. Thus, the term “haplotypes” is used to describe variants of the *S*-locus, whereas the term “alleles” is used to describe variants of an *S*-locus gene.

For the four GSI families, SI occurs when the *S*-haplotype of the pollen matches either of the two *S*-haplotypes carried by the pistil. That is, the SI phenotype of the pollen (gametophyte) is determined by its own *S*-genotype. For the SSI family, in the simplest case, SI occurs when the pollen-producing parent shares one or both *S*-haplotypes with the pistil. That is, the SI phenotype of the pollen is determined by the *S*-genotype of its diploid parent. For SSI, complex relationships often exist between the different *S*-haplotypes of the pollen and pistil parents. One *S*-haplotype could be dominant over or recessive to another, or it could interact with another to result in mutual weakening or in an entirely new *S*-haplotype specificity (Thompson and Taylor, 1966).

THREE TYPES OF SI MECHANISMS

During the past two decades, much progress has been made in identifying and characterizing the *S*-locus genes that control the specificity of the SI interaction in the five families mentioned above. Comparisons of the *S*-locus genes expressed in the pistil among the different families have revealed three biochemically distinct mechanisms (Table 1). The Solanaceae, Rosaceae, and Scrophulariaceae use the same mechanism, the Papaveraceae uses another, and the Brassicaceae uses a third. For the Solanaceae and Papaveraceae mechanisms, the gene that controls female specificity has been identified; these genes were named the *S*-RNase gene and the *S*-gene, respectively. Our understanding of the Solanaceae mechanism has progressed further, with the recent identification of a promising candidate for the male specificity gene. The Solanaceae mechanism involves *S*-RNase-mediated degradation of RNA in self-pollen tubes. The Papaveraceae mechanism is mediated by a signal transduction cascade in pollen that involves a number of known components of signal transduction (e.g., Ca²⁺, phosphoinositides, protein kinases, and phosphatases). For the SSI mechanism found in the Brassicaceae, both the gene that controls male specificity, *S*-locus cysteine-rich protein (*SCR*)/*S*-locus protein-11 (*SP11*),

¹ To whom correspondence should be addressed. E-mail txk3@psu.edu; fax 814-863-9416.

Article, publication date, and citation information can be found at www.plantcell.org/cgi/doi/10.1105/tpc.016154.

Table 1. Summary of Three Types of SI Mechanisms

Plant Family	Type of SI	Genetic Locus	Female Determinant	Male Determinant	Mechanism
Solanaceae, Rosaceae, Scrophulariaceae	GSI	S-locus	<i>S-RNase</i>	<i>SLF/SFB?</i>	S-RNase-mediated degradation of pollen tube RNA
Papaveraceae	GSI	S-locus	S-gene	Unknown	S-protein-mediated signaling cascade in pollen
Brassicaceae	SSI	S-locus	<i>SRK</i>	<i>SCR/SP11</i>	Receptor-kinase-mediated signaling in stigma

and the gene that controls female specificity, S-locus receptor kinase (*SRK*), have been identified. The SI response is mediated via a signal transduction cascade in the stigmatic papilla, which is elicited by the interaction of a pollen-borne ligand, *SCR/SP11*, and *SRK*, a receptor kinase in the stigmatic papilla.

The discussion below focuses on the Solanaceae type of SI. For a recent review of the Brassicaceae type of SI, see Kachroo et al. (2002); for a recent review of the Papaveraceae type of SI, see Thomas et al. (2003).

The Solanaceae type of SI was first discovered in *Nicotiana glauca* (East and Mangelsdorf, 1925), and to date, this type of SI has been studied at the molecular level in four genera of the Solanaceae (*Lycopersicon*, *Nicotiana*, *Petunia*, and *Solanum*), three genera of the Rosaceae (*Malus*, *Prunus*, and *Pyrus*), and one genus of the Scrophulariaceae (*Antirrhinum*). The rejection of self-pollen occurs during pollen tube growth in the style, and the timing of the rejection coincides with the transition of pollen tube growth from the slow ("autotrophic") growth phase to the accelerated ("heterotrophic") growth phase (Herrero and Hormaza, 1996). The increase in the growth rate is presumed to result from the increased acquisition of nutrients provided by the pistil tissue. Interestingly, several critical cellular events also occur around the time of this transition (e.g., mitotic division of the generative cell to give rise to two sperm nuclei).

FEMALE SPECIFICITY DETERMINANT: THE S-RNASE GENE

The search for the female determinant of SI was based on the prediction that the gene encoding it must exhibit allele-specific sequence differences and must be expressed in the pistil. Pistil-specific proteins that showed allele-specific differences in molecular mass and/or isoelectric point were first identified in *Nicotiana glauca* (Bredemeijer and Blass, 1981), and the first sequence of such a protein was deduced from the cloning and sequencing of the corresponding cDNA (Anderson et al., 1986). These proteins were initially named S-allele-associated proteins or S-proteins, and the gene was named the pistil S-gene. Similar approaches were used to identify S-proteins and to isolate their cDNAs from other solanaceous species (Ai et al., 1990; Clark et al., 1990) and several rosaceous species (Sassa et al., 1993; Ishimizu et al., 1996).

Sequence comparisons of solanaceous S-proteins have revealed five conserved regions and two hypervariable regions (Ioerger et al., 1991; Tsai et al., 1992). The presence of distinct conserved regions has made it possible to clone genomic and cDNA fragments for S-proteins by PCR. This is particularly useful

for population and evolutionary studies of SI (Richman et al., 1995; Wang et al., 2001) and to genotype plants (Tao et al., 1999). For rosaceous species, the ability to identify S-genotypes by this rapid approach is particularly useful because it circumvents the time-consuming and labor-intensive pollination tests. Most notably, this approach led Xue et al. (1996) to clone cDNAs for the homologs of S-proteins in *Antirrhinum hispanicum* and to discover that this species exhibits the same type of SI mechanism as the solanaceous and rosaceous species. Phylogenetic analysis of S-proteins has suggested that the Solanaceae-type mechanisms used by the species in these three distantly related dicot families share a common origin, and that this mechanism might be exhibited by the ancestor of ~75% of all dicots (Igic and Kohn, 2001). At present, it is not known whether the Solanaceae mechanism is exhibited by any other dicot families. This question can be addressed by using PCR to ascertain whether orthologs of S-proteins are present in any other dicot families.

Biochemical and Structural Characteristics of S-Proteins/S-RNases

The biochemical nature of S-proteins was revealed when the sequence of RNase T₂ of *Aspergillus oryzae* (Kawata et al., 1988) was determined and found, unexpectedly, to share sequence similarity with S-proteins (McClure et al., 1989). This finding led to the subsequent confirmation that S-proteins have RNase activity in vitro (McClure et al., 1989; Singh et al., 1991). Significantly, two regions of RNase T₂, each of which contains a catalytic His, are now known to be present in all S-proteins and to correspond to two of the five conserved regions identified by Ioerger et al. (1991). Thus, S-proteins have been renamed S-RNases and the gene has been renamed the *S-RNase* gene. S-RNases do not appear to have any substrate specificity in vitro (Singh et al., 1991). RNase T₂ and S-RNases have been placed in a large family of RNases, named the T₂/S-RNase family, which also includes S-like RNases and relic S-RNases (Green, 1994; Golz et al., 1998). S-like RNases do not exhibit allelic sequence diversity, and they have been identified from both self-incompatible and self-compatible species of the Solanaceae, Rosaceae, and Scrophulariaceae as well as from self-compatible species of several other families. Relic S-RNases have been identified from both self-incompatible and self-compatible species of the Solanaceae, Rosaceae, and Scrophulariaceae, and they are more similar to S-RNases than to S-like RNases and, like S-RNases, are specific to the pistil. However, relic S-RNases, like S-like RNases, do not exhibit allelic sequence

polymorphism. Relic *S-RNase* genes could have been derived from the *S-RNase* gene as a result of gene duplication, followed by translocation to other loci. Despite their sequence similarity to *S-RNases*, relic *S-RNases* and *S-like RNases* are unlikely to play a role in SI, and to date, the physiological function of most of them remains unknown.

The role of the *S-RNase* gene in the SI interaction has been established via transgenic experiments (Lee et al., 1994; Murfett et al., 1994). These experiments showed that introduction of a new allele of the *S-RNase* gene into transgenic plants was sufficient to confer on the plants the ability to reject pollen carrying the same allele as the introduced *S-RNase* gene. Conversely, suppression of the expression of an endogenous allele of the *S-RNase* gene by the antisense RNA approach abolished the ability of the transgenic plants to reject the pollen carrying the affected allele.

To understand the biochemical mechanism of *S-RNase*-mediated self-rejection, it is imperative to know whether the *RNase* activity of *S-RNases* is an integral part of their function. Site-directed mutagenesis was used to replace the codon for one of the two catalytic His residues of S_3 -*RNase* of *Petunia inflata* with an Asn codon, and transgenic plants that produced S_3 -*RNase* without *RNase* activity failed to reject S_3 pollen (Huang et al., 1994). Consistent with this result is the finding that a self-compatible accession of *Lycopersicon peruvianum* produced a catalytically inactive *S-RNase* with one of the active-site His residues replaced with Arg (Kowyama et al., 1994; Royo et al., 1994). Because the *RNase* activity of *S-RNases* is required for pollen rejection, it is reasonable to infer that the degradation of RNA by *S-RNases* inside self-pollen tubes results in growth inhibition. McClure et al. (1990) obtained results consistent with this prediction. They showed that pollen tube rRNA was degraded after self-pollination but not after cross-pollination. However, the design of the experiments leaves open the possibility that the observed degradation of RNA is a consequence, but not the cause, of growth inhibition of self-pollen tubes.

S-Allele-Specificity Determinant of S-RNases

Attempts have been made to identify the region (or regions) of *S-RNases* that determines *S*-allele specificity (i.e., regions that are involved in the interaction with the pollen *S*-allele products). Because *S-RNases* are glycoproteins that vary in the number and position of N-linked glycan chains, the *S*-allele-specificity determinant could reside, a priori, in the carbohydrate moiety and/or the protein backbone. To address this question, a nonglycosylated S_3 -*RNase* of *P. inflata* (with the Asn residue for the only N-glycosylation site replaced with Asp) was produced in transgenic plants, and the SI behavior of the pistil was examined. The nonglycosylated S_3 -*RNase* was found to have *RNase* activity similar to that of the wild-type S_3 -*RNase* and to function as well as the wild-type S_3 -*RNase* in rejecting S_3 pollen (Karunanandaa et al., 1994). Thus, the *S*-allele-specificity determinant of *S-RNases* resides in their protein backbone.

One notable feature of *S-RNases* is their high degree of allelic sequence diversity. For example, the two most divergent solanaceous *S-RNases* share only 38% sequence identity (Tsai et al., 1992). There are a large number of variable sites scattered

throughout the protein; however, the most highly variable sites are clustered in two regions, named HVa and HVb. These two hypervariable regions were identified initially from comparison of solanaceous *S-RNases* (Ioerger et al., 1991; Tsai et al., 1992). They were found subsequently to be the hypervariable regions of *Antirrhinum* *S-RNases* as well (Xue et al., 1996) and to correspond to two of the four regions of rosaceous *S-RNases* for which evidence of positive selection has been found (Ishimizu et al., 1998). The crystal structures of a solanaceous *S-RNase* and a rosaceous *S-RNase* show that both HVa and HVb regions are exposed on the surface of the protein and accessible to solvent (Ida et al., 2001; Matsuura et al., 2001). Together, HVa and HVb are considered the most likely candidates for the determinant of *S*-allele specificity.

Domain-swapping experiments have been performed to ascertain the role, if any, of HVa, HVb, and other regions of *S-RNases* in *S*-allele specificity (Kao and McCubbin, 1996; Matton et al., 1997; Zurek et al., 1997). For each chimeric *S-RNase* gene, the bulk of the sequence was from one allele of the *S-RNase* gene, with the sequence of the region to be examined contributed by another allele. Transgenic plants that produce each chimeric *S-RNase* then were examined for their ability to reject pollen of the two alleles used in the chimeric construct. When pairs of *S-RNases* with a high degree of sequence diversity (e.g., 74.1% amino acid identity between S_1 - and S_3 -*RNases* of *P. inflata*) were used in the domain swapping, none of these domains alone could bestow on the chimeric protein its unique *S*-allele specificity. Moreover, the specificity of the allele used as the backbone of the chimeric *S-RNase* gene was abolished (Kao and McCubbin, 1996; Zurek et al., 1997). Because all of these chimeric *S-RNases* retained normal *RNase* activity, their failure to reject self-pollen was attributed to the loss of the recognition function.

When two *S-RNases* of *Solanum chacoense*, S_{11} -*RNase* and S_{13} -*RNase*, that share 92% amino acid sequence identity (with only 10 dissimilar amino acids) were used for the construction of chimeric *S-RNase* genes, it was found that the HVa and HVb regions together were sufficient to confer on the chimeric *S-RNases* the new allelic specificity (Matton et al., 1997). That is, when the amino acids of HVa and HVb of S_{11} -*RNase* were changed to those of S_{13} -*RNase*, transgenic plants that produced this chimeric *S-RNase* rejected S_{13} pollen but not S_{11} pollen. It should be noted, however, that domain-swapping experiments can only address the role of those amino acids exchanged that differ between the two proteins. Thus, the results of Matton et al. (1997) cannot rule out the involvement of amino acids outside of HVa and HVb that are conserved between S_{11} -*RNase* and S_{13} -*RNase* (Verica et al., 1998). The finding that two *P. inflata* *S-RNases*, S_6 -*RNase* and S_9 -*RNase*, have identical sequence in HVa and differ by only two amino acids in HVb (Wang et al., 2001) suggests that amino acids outside of these two hypervariable regions likely are involved in the determination of *S*-allele specificity.

MALE-SPECIFICITY DETERMINANT

Classic genetic studies showed that the pollen and pistil functions in SI could mutate independently to result in either

pollen-part or pistil-part self-compatible mutants (de Nettancourt, 2001). This finding strongly suggests that separate genes control these two functions. In all of the transgenic experiments performed to ascertain the function of the *S-RNase* gene described above, the pistil function, but not the pollen function, was affected by the manipulation of the *S-RNase* gene, consistent with the notion that the S-RNase gene does not control male specificity. Most directly, Sassa et al. (1997) showed that the *S₄-RNase* gene was deleted in a self-compatible cultivar of *Pyrus serotina* (Japanese pear) and that this deletion affected the pistil function but not the pollen function. During the past few years, a flurry of research activities have been directed toward the identification of the gene that controls male specificity, the pollen S-gene.

S-Locus Linked Genes

One approach to identifying the pollen S-gene is to search for pollen-expressed genes that exhibit S-haplotype-specific restriction fragment length polymorphism, because the pollen S-gene is expected to show a significant degree of allelic sequence diversity. A number of such genes have been identified in *N. alata* and *P. inflata* by RNA differential display and subtractive hybridization (Dowd et al., 2000; Li et al., 2000; McCubbin et al., 2000). Recombination analysis has been performed to determine if each of these genes is tightly linked to the *S-RNase* gene. Because recombination at the S-locus is suppressed as a result of its centromeric location (Entani et al., 1999), a large number of plants segregating for S-haplotypes are needed to accurately assess the linkage to the *S-RNase* gene. Nine of the pollen-expressed genes of *P. inflata* (Wang et al., 2004) and one of the pollen-expressed genes of *N. alata* (Li et al., 2000) were found to be tightly linked to the *S-RNase* gene. The nucleotide sequences of different alleles of each of these genes were obtained to assess the allelic sequence diversity. The deduced amino acid sequences of all of these genes exhibit very low allelic sequence diversity. Also, analysis of the nucleotide sequence variation of 48A of *N. alata* has revealed no evidence of positive selection, which is expected of the S-locus genes involved in SI (Takebayashi et al., 2003). In the case of *P. inflata*, chromosome walking through the *S₂-locus* region has shown that the nine genes that are tightly linked to the *S-RNase* gene are located at least ~180 kb upstream or at least ~700 kb downstream from the *S-RNase* gene and could be as far as 4 Mb away (T. Tsukamoto, Y. Wang, K.-W. Yi, A.G. McCubbin, and T.-h. Kao, unpublished data). Thus, none of the pollen-expressed genes identified by this approach is likely to be the pollen S-gene, and their allelic sequence polymorphism may result simply from their tight genetic linkage to the highly polymorphic S-locus.

The S-Locus F-Box Gene: A Candidate for the Male-Specificity Gene

Another approach to identifying the pollen S-gene is based on the prediction that the pollen S-gene must be very tightly linked to the *S-RNase* gene. Recombination in the chromosomal region between these two genes would inevitably result in the breakdown of SI by generating different S-haplotype specificities

for pollen and pistil, but such recombinant genotypes have never been observed (de Nettancourt, 2001). Lai et al. (2002) sequenced a 63-kb region containing the *S₂-RNase* gene of *A. hispanicum* and identified 10 additional open reading frames (ORFs); 4 encode retrotransposons and only 1 of the remaining ORFs is expressed in the anther (tapetum and pollen). This gene, located ~9 kb downstream of the *S₂-RNase* gene, encodes an F-box-containing protein and was named *AhSLF* (*A. hispanicum* S-locus F-box gene). cDNA encoding a homolog of *AhSLF-S₂* was isolated from a line of *S₁S₅* genotype, and its deduced amino acid sequence is 97.9% identical to that of *AhSLF-S₂*. However, it is not clear from that report whether *AhSLF-S₂* and its homolog, named *AhSLF-S₂L*, are allelic.

Entani et al. (2003) and Ushijima et al. (2003) also attempted to identify the pollen S-gene by sequencing the S-loci of *Prunus mume* (Japanese apricot) and *Prunus dulcis* (almond), respectively. An ~70 kb chromosomal region of *P. dulcis* that contains the *S^c-RNase* gene is considered the functional region of the S-locus (Ushijima et al., 2001) based on the following findings. First, genomic blot analysis showed that the sequence of this region is highly divergent between different S-haplotypes, whereas the sequences flanking this region are similar between different S-haplotypes. Second, a self-compatible mutant with a chromosomal deletion spanning this region was defective in both pollen and pistil functions. Sequencing of this ~70-kb region revealed 10 ORFs in addition to the *S^c-RNase* gene. As with the *Antirrhinum S₂-locus*, some of the ORFs encode retrotransposons. Only two of the other ORFs are expressed in pollen, and interestingly, both encode F-box proteins. One of the F-box genes, named *SFB* (S-haplotype-specific F-box gene), exhibits a similarly high level of allelic sequence diversity as the *S-RNase* gene. The pairwise amino acid sequence identities of *S^a*-, *S^b*-, *S^c*-, and *S^d*-alleles range from 68.4 to 76.4%, and those of the same four alleles of the *S-RNase* gene range from 55.6 to 77.1% (Yamane et al., 2003). For the other F-box gene, named *PdSLF* (*P. dulcis* S-locus F-box gene), the deduced amino acid sequences of *S^c*- and *S^d*-alleles are 95.1% identical. *SFB* is a good candidate for the pollen S-gene, because (1) it is linked physically to the *S-RNase* gene, located within 30 kb of the *S-RNase* gene in the four S-haplotypes studied; (2) it is expressed specifically in pollen; and (3) it shows a high level of allelic sequence diversity. By contrast, *PdSLF*, despite its physical linkage to the *S-RNase* gene and its pollen expression, is considered unlikely to be the pollen S-gene because of the low degree of allelic sequence diversity. It should be noted that *PdSLF* was so named to emphasize that it shows approximately the same low level of allelic sequence diversity as *AhSLF* identified in *Antirrhinum* (Lai et al., 2002) rather than to indicate that *PdSLF* is an ortholog of *AhSLF*. In fact, both *SFB* and *PdSLF* are <25% identical to *AhSLF* in their deduced amino acid sequences.

Entani et al. (2003) identified four F-box genes in a 62.5-kb region of the S-locus of *P. mume* that contains the *S₇-RNase* gene; three of them also are located in a 64-kb region of the S-locus containing the *S₇-RNase* gene. The F-box gene closest to the *S-RNase* gene was named *SLF*; it is expressed in pollen and shows a high level of allelic sequence diversity. The amino acid sequence identities of the *S₇*-, *S₇*-, and *S₉*-alleles of *SLF*

range from 77.8 to 81.3%. Thus, *SLF* of *P. mume* is most likely the ortholog of *SFB* of *P. dulcis*. Two alleles of *SLF* in another rosaceous species, *P. avium* (sweet cherry), have been cloned by PCR, and their deduced amino acid sequences are 79.5% identical (Yamane et al., 2003). Hereafter, *SLF/SFB* will be used to indicate the F-box gene that is the prime candidate for the pollen *S*-gene. The other three F-box genes identified in *P. mume* were named *SLFL1*, *SLFL2*, and *SLFL3*, all of which, like *PdSLF* of *P. dulcis*, show much lower degrees of allelic sequence diversity than *SLF/SFB*. For example, the deduced amino acid sequences of the *S*₁- and *S*₂-alleles of *SLFL1* are 92.5% identical.

An alignment of the amino acid sequences of all nine *SLF/SFB* of the three *Prunus* species is shown in Figure 1. The F-box motif is located at the N terminus, and it is relatively conserved among these proteins. To identify regions of variability, the Normed Variability Index (NVI; as defined by Ioerger et al., 1991) was calculated for each site of the alignment, and the NVI of each

site was averaged with that of its neighbors in a sliding window of size 7. A window-averaged plot of NVI over the length of *SLF/SFB* identified two very prominent variable regions, named *HVa* and *HVb*, which correspond approximately to the variable regions A and B, respectively, identified from the sequence comparison of four *P. dulcis* *SFBs* (Ushijima et al., 2003). These two regions together contain 8 of the 13 most variable sites, and the *HVa* region in particular shows the highest peak of window-averaged NVI. The presence of these hypervariable regions is consistent with the potential role of *SLF/SFB* in determining male specificity.

Most F-box proteins are involved in ubiquitin-mediated protein degradation. This system uses E1 (ubiquitin-activating enzyme), E2 (ubiquitin-conjugating enzyme), and E3 (ubiquitin ligase) to catalyze the formation of polyubiquitin chains on specific substrates for degradation by the 26S proteasome (Bai et al., 1996). An F-box protein is a component of one class of E3, called SCF, which also consists of Skp1, cullin (Cul1), and a RING-HC




Figure 1. Alignment of Deduced Amino Acid Sequences of *SLF/SFBs* of Three *Prunus* Species.

The sequences were aligned by DNASIS version 3.5 (Hitachi Software Engineering Co. Ltd., Tokyo, Japan). The two variable regions, *HVa* and *HVb*, were identified as described in the text. The 13 most variable residues (with NVI scores > 0.1016) are indicated with asterisks. For each site, the most frequent residue is shown on a dark background in each sequence. The names of the proteins are indicated by the two-letter abbreviation of the species name (Pa, *P. avium*; Pd, *P. dulcis*; Pm, *P. mume*) followed by the protein name (*SFB* or *SLF*) and the identity of the *S*-allele. Amino acid residue numbers of the first and last amino acids in each row of sequence are indicated at left and right, respectively.

plumbaginifolia. Homologs of *HT-B* also have been identified in two other genera of the Solanaceae, *Lycopersicon* and *Solanum* (Kondo et al., 2002; O'Brien et al., 2002). Suppression of the expression of *HT-B* by antisense RNA and/or RNA interference led to the loss of S-haplotype-specific rejection of pollen in both *N. alata* and *S. chacoense* (McClure et al., 1999; O'Brien et al., 2002). Also, a cultivar of tomato (*Lycopersicon esculentum*) was found to carry a defective *HT-B* in addition to not producing any S-RNase (Kondo et al., 2002). *HT-B* encodes a protein that contains a stretch of 20 Asn and Asp residues (the ND domain) near the C terminus (Cruz-Garcia et al., 2003). Database searches have not yielded any insight into the possible function of *HT-B*. Because the transcript or protein level of the S-RNase gene was not affected in the antisense and RNA interference transgenic plants, *HT-B* is not required for the expression of the S-RNase gene. No direct interaction between *HT-B* and S-RNases has been detected. A current hypothesis is that *HT-B*, perhaps working in conjunction with other pistil proteins, is required for the uptake of S-RNases into pollen tubes (Cruz-Garcia et al., 2003). *HT-B* is the first modifier gene of GSI to be cloned, and the elucidation of its role in SI will contribute to a better understanding of the mechanism of S-RNase-based SI.

Another approach to identifying the modifier genes is to isolate pistil and pollen proteins that interact with S-RNases. A pollen protein of *Petunia hybrida* that interacts with the N-terminal part of S-RNases containing the HVa and HVb regions has been identified by the yeast two-hybrid protein-protein interaction assay (Sims and Ordanic, 2001). This protein, named PhSBP1, is a novel protein and contains a RING-HC motif at its C terminus. RING-HC proteins also are involved in ubiquitin-mediated protein degradation. If *PhSBP1* plays a role in SI, it is more likely to be a general one and not as the S-allele-specificity determinant, because *PhSBP1* is expressed in other tissues, it does not show any allelic sequence polymorphism, and the interaction between its protein product and S-RNases is not allele specific.

Affinity chromatography has been used to identify pistil proteins that interact with S-RNases of *N. alata* (McClure et al., 2000; Cruz-Garcia et al., 2003). Interestingly, three of the five proteins identified to date are pistil glycoproteins that have previously been implicated in pollen tube growth. A 120-kD protein, first identified in *N. alata*, has been shown to be taken up into the cytoplasm of pollen tubes (Lind et al., 1996). TTS, first identified in *N. tabacum*, has been shown to be associated with the pollen tube membrane (Cheung et al., 1993; Wu et al., 1995). PELP1, first identified in *N. tabacum*, has been shown to be associated with the callose wall and callose plugs of pollen tubes (de Graaf et al., 2003). Because these proteins are located in the extracellular space of the transmitting tract tissue and their interactions with S-RNases are not allele specific (at least in vitro), they are thought to play a role in facilitating the uptake of S-RNases into pollen tubes (Cruz-Garcia et al., 2003). In addition to these three glycoproteins, an 11-kD protein with sequence similarity to a class of copper binding proteins called phytochelatins, as well as S-RNase itself, also were found to interact with S-RNases. It is not clear what role the 11-kD protein might have, nor is it known whether S-RNases form dimers or higher multimers in vivo.

It should be noted that because none of these proteins that interact with the S-RNase is specific to SI, they are likely to be required for some other physiological processes as well. Thus, mutations in their genes could be lethal and not recoverable from genetic screens of self-compatible mutants.

MODELS FOR S-RNASE-MEDIATED SI RESPONSE

Because the RNase activity of S-RNases is essential for their function in SI, it is generally accepted, though not yet demonstrated definitively, that the degradation of pollen tube RNAs by the self S-RNase results in the growth inhibition of self-pollen tubes in the style. Two different models (and their modified versions) have been put forward to explain how S-RNases might mediate the specific degradation of self-pollen tube RNA (Thompson and Kirch, 1992; Kao and McCubbin, 1996). The receptor model predicts that the specificity lies in the uptake of S-RNases into a pollen tube: the self S-RNase, but not any non-self S-RNase, would be taken up into the cytoplasm of a pollen tube. Thus, this model predicts that the products of pollen S-alleles are membrane-bound or cell wall-bound receptors that serve as gatekeepers so that only the matching (self) S-RNase is allowed to enter a pollen tube. The inhibitor model, in its simplest form, predicts that the products of pollen S-alleles are cytosolic RNase inhibitors, with each allelic product specifically inhibiting the RNase activity of all non-self S-RNases but not that of the self S-RNase.

One approach to assess the validity of these two models is to determine if the uptake of S-RNases into a pollen tube is S-haplotype specific, as predicted by the receptor model, or if both self and non-self S-RNases are taken up by a pollen tube, as predicted by the inhibitor model. Luu et al. (2000) used immunocytochemistry to localize S₁₁-RNase of *S. chacoense* in pistils (of S₁₁S₁₃ genotype) that had been pollinated with incompatible or compatible pollen. They found that S₁₁-RNase was present in the cytoplasm of both self-pollen tubes (S₁₁ genotype) and non-self pollen tubes of S₁₂ and S₁₃ genotypes. This finding supports the inhibitor model and suggests that if the uptake of S-RNases by pollen tubes requires a receptor (or a receptor complex), it most likely recognizes some common features of S-RNases.

Simple Inhibitor Model

One way that S-haplotype-specific inhibition of S-RNases could be accomplished is depicted in Figure 2A. It is reasonable to predict that S-RNases contain two separate functional domains, an S-allele-specificity domain, which is unique to each S-RNase, and a catalytic domain, which is common to all S-RNases. As described above, the chimeric S-RNases of *P. inflata* and *N. alata* produced in the domain-swapping experiments (Kao and McCubbin, 1996; Zurek et al., 1997) all retained RNase activity, even though they did not retain the original S-allele specificity or gain a new one, suggesting the existence of two separate domains. Similarly, the RNase inhibitors produced by pollen S-alleles also could contain two separate domains, an RNase-inhibitor domain and an S-allele-specificity domain.




Figure 2. Models for S-Haplotype-Specific Inhibition of Pollen Tube Growth and Competitive Interaction.

(A) Simple inhibitor model. Both S_1 - and S_2 -RNases are taken up by an S_1 pollen tube, but only S_1 -RNase is active in degrading pollen RNA because the pollen S_1 -allele product specifically inhibits the RNase activity of S_2 -RNase. Binding of S_1 -RNase to the pollen S_1 -allele product through their matching S-allele-specificity domains (SD) blocks binding of the inhibitor domain (ID) of the pollen S_1 -allele product to the catalytic domain (CD) of S_1 -RNase. Interaction between S_2 -RNase and the pollen S_1 -allele product is through the CD of the former and the ID of the latter, thus rendering S_2 -RNase inactive.

(B) Modified inhibitor model. This model differs from the simple inhibitor model in several respects: pollen S-allele products are homotetramers and they only contain the S-allele-specificity domain; and a general RNase inhibitor is responsible for the inhibition of the RNase activity of S-RNases. The S-haplotype-specific inhibition of the RNase activity of S-RNases is achieved in a manner similar to that described for **(A)**.

(C) Competitive interaction based on the modified inhibitor model. Pollen S_1 -allele and S_2 -allele products mainly form heterotetramers, which cannot bind the S-allele-specificity domain of either S_1 -RNase or S_2 -RNase. As a result, the RNase activities of both S_1 - and S_2 -RNases are inhibited by the general RNase inhibitor.

In all of these models, additional pistil proteins that are required for the SI response are not depicted.

As depicted in Figure 2A, when a pollen tube of S_1 genotype is growing in a pistil of S_1S_2 genotype, the pollen S_1 -allele product interacts with S_1 - and S_2 -RNases differently. In the case of self-interaction, the S-allele-specificity domain of the pollen S_1 -allele product would interact with the S-allele-specificity domain of S_1 -RNase by virtue of the match between the paired domains; the RNase activity of S_1 -RNase would not be affected in such an interaction. However, in the case of non-self interaction, the RNase-inhibitor domain of the pollen S_1 -allele product would interact with the catalytic domain of S_2 -RNase in the absence of the match between their S-allele-specificity domains; such an interaction would inhibit the RNase activity of S_2 -RNase. This model is predicated on the assumption that the interaction between the S-allele-specificity domains of a pollen S-allele product and its cognate S-RNase is stronger than the interaction between the RNase-inhibitor domain of the pollen S-allele product and the catalytic domain of its cognate S-RNase. Moreover, binding to one domain of the S-RNase precludes binding to the other domain. Thus, binding of a pollen S-allele

product to the S-allele-specificity domain locks the S-RNase into its active form.

This inhibitor model can explain a well-known phenomenon termed competitive interaction, which refers to a breakdown of pollen function in SI caused by the presence of two S-loci of different haplotypes in the pollen. This occurs when the entire S-locus, or a critical part of it, is duplicated in diploid plants that carry two different S-haplotypes or when diploid SI plants (carrying two different S-haplotypes) become tetraploid (de Nettancourt, 2001). S-locus duplication can be generated by γ -ray irradiation, with the duplicated region existing as a free centric fragment or as a translocated chromosomal segment. Among the pollen grains produced by such pollen-part self-compatible mutants, only those whose resident S-locus is of a different S-haplotype from that of the extra S-locus fail to function in SI. For example, if an S_1S_2 plant carries an extra S_1 -locus, the only pollen grains that will not be rejected by the pistil upon self-pollination are S_2 pollen grains that carry the duplicated S_1 -locus. Golz et al. (1999, 2001) found that in some cases the duplicated region lacked the S-RNase gene but contained markers that flank the S-locus, consistent with the belief that the duplicated pollen S-allele causes competitive interaction. Pollen carrying two different pollen S-alleles may be referred to as heteroallelic pollen (Luu et al., 2001). The inhibitor model predicts that when two different pollen S-alleles are expressed in the same pollen grain, their products together inhibit the RNase activity of all S-RNases, thus resulting in the breakdown of SI.

Modified Inhibitor Model

A modified inhibitor model was proposed by Luu et al. (2001) to explain the SI behavior of a dual-specificity chimeric S-RNase they had generated (Matton et al., 1999). This S-RNase, named $S_{11/13}$ -RNase, is a chimeric protein between S_{11} - and S_{13} -RNases of *S. chacoense* that exhibits both S_{11} - and S_{13} -allele specificities. Plants that produce the $S_{11/13}$ -RNase rejected both S_{11} and S_{13} pollen. Interestingly, pistils of transgenic plants that produce the $S_{11/13}$ -RNase were completely incompatible with pollen from tetraploid plants of $S_{11}S_{11}S_{13}S_{13}$ genotype, suggesting that the diploid pollen of $S_{11}S_{13}$ genotype produced by the tetraploids was rejected by the $S_{11/13}$ -RNase. Luu et al. (2001) further showed that diploid pollen of $S_{11}S_{13}$ genotype was compatible with pistils producing monospecific S_{11} - and S_{13} -RNases, as would be expected from competitive interaction. Thus, the dual-specificity $S_{11/13}$ -RNase behaves differently from its two corresponding monospecific S-RNases in its ability to recognize and reject diploid pollen carrying two different S-alleles. This finding also suggests that both pollen S-alleles are expressed in diploid pollen and rules out the possibility that the breakdown of SI caused by competitive interaction is attributable to silencing of the expression of the two pollen S-alleles.

The modified inhibitor model predicts that (1) the active form of pollen S-allele products is a homotetramer; (2) the pollen S-allele products contain only the S-allele-specificity domain; and (3) a general RNase inhibitor is responsible for the inhibition of S-RNases. According to this model (Figure 2B), the general inhibitor would bind and inactivate all S-RNases, unless an

S-RNase were bound to its cognate pollen S-allele product through their matching S-allele-specificity domains. When a pollen grain expresses two different alleles of the pollen S-gene, the products would mainly form heterotetramers, which could not efficiently bind either cognate S-RNase; as a result, the general inhibitor would inhibit the RNase activity of both cognate S-RNases (Figure 2C). In fact, this model also predicts that the heterotetramers could not bind any other S-RNases. Thus, pollen tubes carrying two different pollen S-alleles would be compatible with pistils of any S-genotype. In the case of the dual-specificity S-RNase, because it could still bind the heterotetramers formed by the two pollen S-allele products, binding of the general inhibitors to the S-RNase would be blocked. This would explain the SI behavior of the dual-specificity S-RNase.

Potential Role of SLF/SFB in SI

If *SLF/SFB* indeed encodes the male-specificity determinant, each allelic product could interact with its cognate and non-self S-RNases differently, so that only non-self S-RNases are degraded by the 26S proteasome. This would be consistent with the inhibitor model, except that it is the stability, and not the RNase activity, of S-RNases that is regulated by the SI interaction. The S-allele-specific degradation could be accomplished if the interaction between the S-allele-specificity domains of an SLF/SFB and its cognate S-RNase would some-

how prevent ubiquitination of the self S-RNase and the lack of interaction between the S-allele-specificity domains of an SLF/SFB and non-self S-RNases would result in ubiquitination of the latter.

One could take advantage of the competitive interaction phenomenon to examine the function of *SLF/SFB*, as illustrated in Figure 3. Here, the S_1 -allele of *SLF/SFB* (S_1 -*SLF/SFB*) is used to transform S_1S_2 plants and the pollen of the resulting transgenic plants is used to pollinate an S_1S_2 tester plant. If *SLF/SFB* is the pollen S-gene, S_2 pollen that carries the S_1 -*SLF/SFB* transgene will be compatible with S_1S_2 pistils because of competitive interaction. Thus, this pollination will be compatible and result in two different genotypes in the progeny, $S_1S_2+S_1$ -*SLF/SFB* and $S_2S_2+S_1$ -*SLF/SFB*. That is, all of the progeny should inherit the transgene and no S_1S_1 genotype will be obtained in the progeny. Importantly, the latter result will serve as an internal control for the breakdown of SI attributable to competitive interaction. Moreover, the $S_1S_2+S_1$ -*SLF/SFB* progeny will be compatible with S_1S_2 pistils and the $S_2S_2+S_1$ -*SLF/SFB* progeny will be compatible with S_2S_2 pistils.

If *SLF/SFB* is confirmed to be the pollen S-gene, it would be of interest to determine what effect the suppression of its expression has on SI behavior. If *SLF/SFB* contains both the S-allele-specificity domain and the inhibitor domain, as predicted by the simple inhibitor model, the absence of this protein would render the pollen unable to inhibit any S-RNases. For




Figure 3. Scheme of a Transgenic Approach to Ascertain the Function of *SLF/SFB* in SI.

(A) Pollination of an S_1S_2 pistil with pollen from an S_1S_2 transgenic plant that carries an S_1 -*SLF/SFB* transgene. The pollination will be compatible if *SLF/SFB* is the pollen S-gene. Among the pollen produced by the transgenic plant, S_2 pollen carrying the S_1 -*SLF/SFB* transgene will be compatible with the S_1S_2 pistil because of competitive interaction, whereas S_1 pollen, S_2 pollen, and S_1 pollen carrying the S_1 -*SLF/SFB* transgene will be rejected by the pistil. Two S-genotypes will be produced in the progeny, $S_1S_2+S_1$ -*SLF/SFB* transgene and $S_2S_2+S_1$ -*SLF/SFB* transgene.

(B) SI behavior of the progeny with the genotype of $S_1S_2+S_1$ -*SLF/SFB* transgene. Such progeny should be compatible with an S_1S_2 pistil because S_2 pollen carrying the transgene will not be rejected by the pistil as a result of competitive interaction.

(C) SI behavior of the progeny with the genotype of $S_2S_2+S_1$ -*SLF/SFB* transgene. For the same reason given in **(B)**, such progeny should be compatible with an S_2S_2 pistil.

Dark circles indicate pollen grains that carry the transgene.

example, if the antisense S_1 -allele of *SLF/SFB* were introduced into S_1S_1 plants, half of the pollen would express the transgene and the other half would not. The former should be incompatible with pistils of any S -genotype, whereas the latter should behave normally in SI. Thus, if pollen from the transgenic plants were used to pollinate S_1S_1 and S_2S_2 plants, only the S_2S_2 plants would set seed and none of the progeny would carry the transgene. Moreover, the transgene should transmit normally through pollen in crosses with plants that do not produce functional S-RNase. However, if *SLF/SFB* contains only the S -allele-specificity domain, as predicted by the modified inhibitor model, suppression of the expression of *SLF/SFB* would render the pollen unable to block the action of the general inhibitor on any S-RNases. As a result, the pollen not producing *SLF/SFB* would be compatible with pistils of any S -genotypes. Therefore, in the crosses described above, both S_1S_1 and S_2S_2 plants would set seed, with all of the S_1S_1 progeny carrying the antisense transgene and half of the S_2S_2 progeny carrying the antisense transgene.

FUTURE PERSPECTIVES

Since the discovery of the *S-RNase* gene almost two decades ago, much of what we have learned about the Solanaceae type of SI is limited to this female determinant of the SI interaction. The recent identification of the *SLF/SFB* gene very likely will change the landscape of research in this type of SI. The most urgent task, in the short run, is to determine, by in vivo approaches, whether *SLF/SFB* encodes the male-specificity determinant of SI. If *SLF/SFB* is confirmed to be the pollen S -gene, this will open new avenues of research and bring us closer to an understanding of the mechanism of S -haplotype-specific inhibition of pollen tube growth. Questions can be asked regarding whether *SLF/SFB* functions as a conventional F-box protein in mediating the specific degradation of all non-self S-RNases or whether it functions in some unexpected manner. With the genes that encode both the male and female determinants in hand, we also could address one of the most perplexing questions about any type of SI systems: how did the male and female specificity genes coevolve to maintain SI? The fact that multiple F-box genes are linked to the *S-RNase* gene in all three families that exhibit the Solanaceae-type SI also raises questions about the physiological functions of the non-SI-related F-box genes that are linked to the S -locus and about the evolutionary relationships among the various S -linked F-box genes.

Although it is important to focus on how S -haplotype specificity is determined, we also should keep in mind that additional proteins are required for the full manifestation of the SI response. Because most of the candidate proteins identified to date do not appear to be specific to the SI system, understanding how they function in SI will likely have implications for other developmental processes.

Accession Numbers

The GenBank accession numbers for the sequences shown in Figure 1 are AB096857 (PaSFB3), AB096858 (PaSFB6), AB092966 (PdSFBa), AB092967 (PdSFBb), AB079776

(PdSFBc), AB081648 (PdSFBd), AB092621 (PmSLF1), AB092622 (PmSLF7), and AB092645 (PmSLF9).

ACKNOWLEDGMENTS

We thank Ming Tien for assistance in the analysis of *SLF/SFB* sequences, Ryutaro Tao, Marcy Uyenoyama, and Yongbiao Xue for communicating unpublished results, and Bruce A. McClure, Andrew G. McCubbin, and Anu Singh-Cundy for valuable comments. Research in our laboratory was supported by grants from the National Science Foundation.

Received August 8, 2003; accepted December 26, 2003.

REFERENCES

- Ai, Y., Kron, E., and Kao, T.H. (1991). S -alleles are retained and expressed in a self-compatible cultivar of *Petunia hybrida*. *Mol. Gen. Genet.* **230**, 353–358.
- Ai, Y., Singh, A., Coleman, C.E., Ioerger, T.R., Kheyr-Pour, A., and Kao, T.H. (1990). Self-incompatibility in *Petunia inflata*: Isolation and characterization of cDNAs encoding three S -allele-associated proteins. *Sex. Plant Reprod.* **3**, 130–138.
- Anderson, M.A., et al. (1986). Cloning of cDNA for a stylar glycoprotein associated with expression of self-incompatibility in *Nicotiana glauca*. *Nature* **321**, 38–44.
- Bai, C., Sen, P., Hofman, K., Ma, L., Goebel, M., Harper, W., and Elledge, S. (1996). Skp1 connects cell cycle regulation to the ubiquitin proteolysis machinery through a novel motif, the F-box. *Cell* **86**, 263–274.
- Bernatzky, R., Glaven, R.H., and Rivers, B.A. (1995). S -related protein can be recombinant with self-compatibility in interspecific derivatives of *Lycopersicon*. *Biochem. Genet.* **33**, 215–225.
- Bredemeijer, G.M., and Blass, J. (1981). S -specific proteins in styles of self-incompatible *Nicotiana glauca*. *Theor. Appl. Genet.* **59**, 185–190.
- Cenciarelli, C., Chiaur, D.S., Guardavaccaro, D., Parks, W., Vidal, M., and Pagano, M. (1999). Identification of a family of human F-box proteins. *Curr. Biol.* **9**, 1177–1179.
- Cheung, A.Y., May, B., Kawata, E., Gu, Q., and Wu, H.M. (1993). Characterization of cDNAs for stylar transmitting tissue-specific proline-rich proteins in tobacco. *Plant J.* **3**, 151–160.
- Chung, I.K., Lee, S.Y., Lim, P.O., Oh, S.A., Kim, Y.S., and Nam, H.G. (1999). An S RNase gene of *Lycopersicon peruvianum* L. is highly expressed in transgenic tobacco but does not affect self-incompatibility. *J. Plant Physiol.* **154**, 63–70.
- Clark, K.R., Okuley, J.J., Collins, P.D., and Sims, T.L. (1990). Sequence variability and developmental expression of S -alleles in self-incompatible and pseudo-self-compatible petunia. *Plant Cell* **2**, 815–826.
- Cruz-Garcia, F., Hancock, C.N., and McClure, B. (2003). S-RNase complexes and pollen rejection. *J. Exp. Bot.* **54**, 123–130.
- de Graaf, B.H.J., Knuiman, B.A., Derksen, J., and Mariani, C. (2003). Characterization and localization of the transmitting tissue-specific PELP III proteins of *Nicotiana glauca*. *J. Exp. Bot.* **54**, 55–63.
- de Nettancourt, D. (2001). Incompatibility and Incongruity in Wild and Cultivated Plants. (Berlin: Springer-Verlag).
- Dowd, P.E., McCubbin, A.G., Wang, X., Verica, J.A., Tsukamoto, T., Ando, T., and Kao, T.H. (2000). Use of *Petunia* as a model for the study of solanaceous type self-incompatibility. *Ann. Bot. Suppl. A* **85**, 87–93.

- East, E.M., and Mangelsdorf, A.J.** (1925). A new interpretation of the hereditary behavior of self-sterile plants. *Proc. Natl. Acad. Sci. USA* **11**, 166–171.
- Entani, T., Iwano, M., Shiba, H., Che, F.-S., Isogai, A., and Takayama, S.** (2003). Comparative analysis of the self-incompatibility (S-) locus region of *Prunus mume*: Identification of a pollen-expressed F-box gene with allelic diversity. *Genes Cells* **8**, 203–213.
- Entani, T., Iwano, M., Shiba, H., Takayama, S., Fukui, K., and Isogai, A.** (1999). Centromeric localization of an S-RNase gene in *Petunia hybrida* Vilm. *Theor. Appl. Genet.* **99**, 391–397.
- Golz, J.F., Clarke, A.E., Newbigin, E., and Anderson, M.A.** (1998). A relic S-RNase is expressed in the styles of self-compatible *Nicotiana sylvestris*. *Plant J.* **16**, 591–596.
- Golz, J.F., Oh, H.-Y., Su, V., Kusaba, M., and Newbigin, E.** (2001). Genetic analysis of *Nicotiana* pollen-part mutants is consistent with the presence of an S-ribonuclease inhibitor at the S locus. *Proc. Natl. Acad. Sci. USA* **98**, 15372–15376.
- Golz, J.F., Su, V., Clarke, A.E., and Newbigin, E.** (1999). A molecular description of mutations affecting the pollen component of the *Nicotiana alata* S locus. *Genetics* **152**, 1123–1135.
- Green, P.J.** (1994). The ribonucleases of higher plants. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **45**, 421–445.
- Herrero, M., and Hormaza, J.I.** (1996). Pistil strategies controlling pollen tube growth. *Sex. Plant Reprod.* **9**, 343–347.
- Huang, S., Lee, H.-S., Karunanandaa, B., and Kao, T.H.** (1994). Ribonuclease activity of *Petunia inflata* S proteins is essential for rejection of self-pollen. *Plant Cell* **6**, 1021–1028.
- Ida, K., Norioka, S., Yamamoto, M., Kumasaka, T., Yamashita, E., Newbigin, E., Clarke, A.E., Sakiyama, F., and Sato, M.** (2001). The 1.55 Å resolution structure of *Nicotiana alata* S_{F11}-RNase associated with gametophytic self-incompatibility. *J. Mol. Biol.* **314**, 103–112.
- Igic, B., and Kohn, J.R.** (2001). Evolutionary relationships among self-incompatibility RNases. *Proc. Natl. Acad. Sci. USA* **98**, 13167–13171.
- Ioerger, T.R., Gohlke, J.R., Xu, B., and Kao, T.H.** (1991). Primary structural features of the self-incompatibility protein in Solanaceae. *Sex. Plant Reprod.* **4**, 81–87.
- Ishimizu, T., Endo, T., Yamaguchi-Kabata, Y., Nakamura, K.T., Sakiyama, F., and Norioka, S.** (1998). Identification of regions in which positive selection may operate in S-RNase of Rosaceae: Implication for S-allele-specific recognition sites in S-RNase. *FEBS Lett.* **440**, 337–342.
- Ishimizu, T., Sato, Y., Saito, T., Yoshimura, Y., Norioka, S., Nakanishi, T., and Sakiyama, F.** (1996). Identification and partial amino acid sequences of seven S-RNases associated with self-incompatibility of the Japanese pear, *Pyrus pyrifolia* Nakai. *J. Biochem.* **120**, 326–334.
- Kachroo, A., Nasrallah, M.E., and Nasrallah, J.B.** (2002). Self-incompatibility in the Brassicaceae: Receptor-ligand signaling and cell-to-cell communication. *Plant Cell* **14** (Suppl.), S227–S238.
- Kao, T.H., and McCubbin, A.G.** (1996). How flowering plants discriminate between self and non-self pollen to prevent inbreeding. *Proc. Natl. Acad. Sci. USA* **93**, 12059–12065.
- Karunanandaa, B., Huang, S., and Kao, T.H.** (1994). Carbohydrate moiety of the *Petunia inflata* S₃ protein is not required for self-incompatibility interactions between pollen and pistil. *Plant Cell* **6**, 1933–1940.
- Kawata, Y., Sakiyama, F., and Tamaoki, H.** (1988). Amino-acid sequence of ribonuclease T₂ from *Aspergillus oryzae*. *Eur. J. Biochem.* **176**, 683–697.
- Kondo, K., Yamamoto, M., Matton, D.P., Sato, T., Hirai, M., Norioka, S., Hattori, T., and Kowiyama, Y.** (2002). Cultivated tomato has defects in both S-RNase and HT genes required for stylar function of self-incompatibility. *Plant J.* **29**, 627–636.
- Kowiyama, Y., Kunz, C., Lewis, I., Newbigin, E., Clarke, A.E., and Anderson, M.A.** (1994). Self-compatibility in a *Lycopersicon peruvianum* variant (LA2157) is associated with a lack of style S-RNase activity. *Theor. Appl. Genet.* **88**, 859–864.
- Lai, Z., Ma, W., Han, B., Liang, L., Zhang, Y., Hong, G., and Xue, Y.** (2002). An F-box gene linked to the self-incompatibility (S) locus of *Antirrhinum* is expressed specifically in pollen and tapetum. *Plant Mol. Biol.* **50**, 29–42.
- Lee, H.-S., Huang, S., and Kao, T.H.** (1994). S proteins control rejection of incompatible pollen in *Petunia inflata*. *Nature* **367**, 560–563.
- Li, J.-H., Nass, N., Kusaba, M., Dodds, P.N., Treloar, N., Clarke, A.E., and Newbigin, E.** (2000). A genetic map of the *Nicotiana alata* S locus that includes three pollen-expressed genes. *Theor. Appl. Genet.* **100**, 956–964.
- Lind, J.L., Bönig, I., Clarke, A.E., and Anderson, M.A.** (1996). A style-specific 120 kDa glycoprotein enters pollen tubes of *Nicotiana alata* in vivo. *Sex. Plant Reprod.* **9**, 75–86.
- Luu, D.-T., Qin, K., Laublin, G., Yang, Q., Morse, D., and Cappadocia, M.** (2001). Rejection of S-heteroallelic pollen by a dual-specific S-RNase in *Solanum chacoense* predicts a multimeric SI pollen component. *Genetics* **159**, 329–335.
- Luu, D.-T., Qin, X., Morse, D., and Cappadocia, M.** (2000). S-RNase uptake by compatible pollen tubes in gametophytic self-incompatibility. *Nature* **407**, 649–651.
- Martin, F.W.** (1961). The inheritance of self-incompatibility in hybrids of *Lycopersicon esculentum* Mill. × *L. chilense* Dun. *Genetics* **46**, 1443–1454.
- Matsuura, T., Sakai, H., Unno, M., Ida, K., Sato, M., Sakiyama, F., and Norioka, S.** (2001). Crystal structure at 1.5-Å resolution of *Pyrus pyrifolia* pistil ribonuclease responsible for gametophytic self-incompatibility. *J. Biol. Chem.* **276**, 45261–45269.
- Matton, D.P., Luu, D.T., Xike, Q., Laublin, G., O'Brien, M., Maes, O., Morse, D., and Cappadocia, M.** (1999). Production of an S-RNase with dual specificity suggests a novel hypothesis for the generation of new S alleles. *Plant Cell* **11**, 2087–2097.
- Matton, D.P., Maes, O., Laublin, G., Xike, Q., Bertrand, C., Morse, D., and Cappadocia, M.** (1997). Hypervariable domains of self-incompatibility RNases mediate allele-specific pollen recognition. *Plant Cell* **9**, 1757–1766.
- McClure, B.A., Cruz-Garcia, F., Beecher, B.S., and Sulaman, W.** (2000). Factors effecting inter- and intra-specific pollen rejection in *Nicotiana*. *Ann. Bot. Suppl. A* **85**, 113–123.
- McClure, B.A., Gray, J.E., Anderson, M.A., and Clarke, A.E.** (1990). Self-incompatibility in *Nicotiana alata* involves degradation of pollen rRNA. *Nature* **347**, 757–760.
- McClure, B.A., Haring, V., Ebert, P.R., Anderson, M.A., Simpson, R.J., Sakiyama, F., and Clarke, A.E.** (1989). Style self-incompatibility gene products of *Nicotiana alata* are ribonucleases. *Nature* **342**, 955–957.
- McClure, B.A., Mou, B., Canevascini, S., and Bernatzky, R.** (1999). A small asparagine-rich protein required for S-allele-specific pollen rejection in *Nicotiana*. *Proc. Natl. Acad. Sci. USA* **96**, 13548–13553.
- McCubbin, A.G., Wang, X., and Kao, T.H.** (2000). Identification of self-incompatibility (S-) locus linked pollen cDNA markers in *Petunia inflata*. *Genome* **43**, 619–627.
- Murfett, J., Atherton, T.L., Mou, B., Gasser, C.S., and McClure, B.A.** (1994). S-RNase expressed in transgenic *Nicotiana* causes S-allele-specific pollen rejection. *Nature* **367**, 563–566.
- Murfett, J.M., Strabala, T.J., Zurek, D.M., Mou, B., Beecher, B., and McClure, B.A.** (1996). S-RNase and interspecific pollen rejection in the genus *Nicotiana*: Multiple pollen-rejection pathways contribute to unilateral incompatibility between self-incompatible and self-compatible species. *Plant Cell* **8**, 943–958.

- O'Brien, M., Kapfer, C., Major, G., Laurin, M., Bertrand, C., Kondo, K., Kowiyama, Y., and Matton, D.P. (2002). Molecular analysis of the stilar-expressed *Solanum chacoense* small asparagine-rich protein family related to the HT modifier of gametophytic self-incompatibility in *Nicotiana*. *Plant J.* **32**, 985–996.
- Richman, A.D., Kao, T.H., Schaeffer, S.W., and Uyenoyama, M.K. (1995). S-allele sequence diversity in natural populations of *Solanum carolinense* (horsenettle). *Heredity* **75**, 405–415.
- Royo, J., Kunz, C., Kowiyama, Y., Anderson, M.A., Clarke, A.E., and Newbigin, E. (1994). Loss of a histidine residue at the active site of S-locus ribonuclease is associated with self-compatibility in *Lycopersicon peruvianum*. *Proc. Natl. Acad. Sci. USA* **91**, 6511–6514.
- Sassa, H., Hirano, H., and Ikehashi, H. (1993). Identification and characterization of stilar glycoproteins associated with self-incompatibility genes of Japanese pear, *Pyrus serotina* Rehd. *Mol. Gen. Genet.* **241**, 17–25.
- Sassa, H., Hirano, H., Nishio, T., and Koba, T. (1997). Style-specific self-incompatibility mutation caused by deletion of the S-RNase gene in Japanese pear (*Pyrus serotina*). *Plant J.* **12**, 223–227.
- Sims, T.L., and Ordanic, M. (2001). Identification of a S-ribonuclease-binding protein in *Petunia hybrida*. *Plant Mol. Biol.* **47**, 771–783.
- Singh, A., Ai, Y., and Kao, T.H. (1991). Characterization of ribonuclease activity of three S-allele-associated proteins of *Petunia inflata*. *Plant Physiol.* **96**, 61–68.
- Takebayashi, N., Brewer, P.B., Newbigin, E., and Uyenoyama, M.K. (2003). Patterns of variation within self-incompatibility loci. *Mol. Biol. Evol.* **20**, 1778–1794.
- Tao, R., Yamane, H., Sugiura, A., Murayama, H., Sassa, H., and Mori, H. (1999). Molecular typing of S-alleles through identification, characterization and cDNA cloning for S-RNases in sweet cherry. *J. Am. Soc. Hortic. Sci.* **124**, 224–233.
- Thomas, S., Osman, K., de Graaf, B.H.J., Shevchenko, G., Wheeler, M.J., Franklin, F.C.H., and Franklin-Tong, V.E. (2003). Investigating mechanisms involved in the self-incompatibility response in *Papaver rhoeas*. *Philos. Trans. R. Soc. Lond. B* **358**, 1033–1036.
- Thompson, K.F., and Taylor, J.P. (1966). Non-linear dominance relationships between S-alleles. *Heredity* **21**, 345–362.
- Thompson, R., and Kirch, H. (1992). The S-locus of flowering plants: When self-rejection is self-interest. *Trends Genet.* **8**, 381–387.
- Tsai, D.-S., Lee, H.-S., Post, L.C., Kreiling, K.M., and Kao, T.H. (1992). Sequence of an S-protein of *Lycopersicon peruvianum* and comparison with other solanaceous S-proteins. *Sex. Plant Reprod.* **5**, 256–263.
- Tsukamoto, T., Ando, T., Kokubun, H., Watanabe, H., Masada, M., Zhu, X., Marchesi, E., and Kao, T.H. (1999). Breakdown of self-incompatibility in a natural population of *Petunia axillaris* (Solanaceae) in Uruguay containing both self-incompatible and self-compatible plants. *Sex. Plant Reprod.* **12**, 6–13.
- Tsukamoto, T., Ando, T., Kokubun, H., Watanabe, H., Sato, T., Masada, M., Marchesi, E., and Kao, T.H. (2003). Breakdown of self-incompatibility in a natural population of *Petunia axillaris* caused by a modifier locus that suppresses the expression of an S-RNase gene. *Sex. Plant Reprod.* **15**, 255–266.
- Tyers, M., and Jorgensen, P. (2000). Proteolysis and the cell cycle: With this RING I do thee destroy. *Curr. Opin. Genet. Dev.* **10**, 54–64.
- Ushijima, K., Sassa, H., Dandekar, A.M., Gradziel, T.M., Tao, R., and Hirano, H. (2003). Structural and transcriptional analysis of the self-incompatibility locus of almond: Identification of a pollen-expressed F-box gene with haplotype-specific polymorphism. *Plant Cell* **15**, 771–781.
- Ushijima, K., Sassa, H., Tamura, M., Kusaba, M., Tao, R., Gradziel, T.M., Dandekar, A.M., and Hirano, H. (2001). Characterization of the S-locus region of almond (*Prunus dulcis*): Analysis of a somaclonal mutant and a cosmid contig for an S haplotype. *Genetics* **158**, 379–386.
- Ushijima, K., Sassa, H., Tao, R., Yamane, H., Dandekar, A.M., Gradziel, T.M., and Hirano, H. (1998). Cloning and characterization of cDNAs encoding S-RNases from almond (*Prunus dulcis*): Primary structural features and sequence diversity of the S-RNases in Rosaceae. *Mol. Gen. Genet.* **260**, 261–268.
- Verica, J.A., McCubbin, A.G., and Kao, T.H. (1998). Are the hyper-variable regions of S-RNases sufficient for allele-specific recognition of pollen? *Plant Cell* **10**, 314–316.
- Wang, X., Hughes, A.L., Tsukamoto, T., Ando, T., and Kao, T.H. (2001). Evidence that intragenic recombination contributes to allelic diversity of the S-RNase gene at the self-incompatibility (S) locus in *Petunia inflata*. *Plant Physiol.* **125**, 1012–1022.
- Wang, Y., Wang, X., McCubbin, A.G., and Kao, T.H. (2004). Genetic mapping and molecular characterization of the self-incompatibility (S-) locus in *Petunia inflata*. *Plant Mol. Biol.*, in press.
- Wu, H.M., Wang, H., and Cheung, A.Y. (1995). A pollen tube growth stimulatory glycoprotein is deglycosylated by pollen tubes and displays a glycosylation gradient in the flower. *Cell* **83**, 395–403.
- Xue, Y., Carpenter, R., Dickinson, H.G., and Coen, E.S. (1996). Origin of allelic diversity in *Antirrhinum* S locus RNases. *Plant Cell* **8**, 805–814.
- Yamane, H., Ikeda, K., Ushijima, K., Sassa, H., and Tao, R. (2003). A pollen-expressed gene for a novel protein with an F-box motif that is very tightly linked to a gene for S-RNase in two species of cherry, *Prunus cerasus* and *P. avium*. *Plant Cell Physiol.* **44**, 764–769.
- Zhou, J., Wang, F., Ma, W., Zhang, Y., Han, B., and Xue, Y. (2003). Structural and transcriptional analysis of S-locus F-box (SLF) genes in *Antirrhinum*. *Sex. Plant Reprod.* **16**, 165–177.
- Zurek, D.M., Mou, B., Beecher, B., and McClure, B. (1997). Exchanging sequence domains between S-RNases from *Nicotiana glauca* disrupts pollen recognition. *Plant J.* **11**, 797–808.

The Molecular and Genetic Bases of S-RNase-Based Self-Incompatibility

Teh-hui Kao and Tatsuya Tsukamoto

Plant Cell 2004;16;S72-S83; originally published online March 9, 2004;

DOI 10.1105/tpc.016154

This information is current as of February 23, 2019

References

This article cites 76 articles, 23 of which can be accessed free at:
/content/16/suppl_1/S72.full.html#ref-list-1

Permissions

https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&issn=1532298X&WT.mc_id=pd_hw1532298X

eTOCs

Sign up for eTOCs at:
<http://www.plantcell.org/cgi/alerts/ctmain>

CiteTrack Alerts

Sign up for CiteTrack Alerts at:
<http://www.plantcell.org/cgi/alerts/ctmain>

Subscription Information

Subscription Information for *The Plant Cell* and *Plant Physiology* is available at:
<http://www.aspb.org/publications/subscriptions.cfm>