

The Exoribonuclease XRN4 Is a Component of the Ethylene Response Pathway in *Arabidopsis*

Thomas Potuschak,^a Amérin Vansiri,^a Brad M. Binder,^b Esther Lechner,^a Richard D. Vierstra,^c and Pascal Genschik^{a,1}

^a Institut de Biologie Moléculaire des Plantes, Laboratoire Propre du Centre National de la Recherche Scientifique, Unité Propre de Recherche 2357, Conventioonné avec l'Université Louis Pasteur, 67084 Strasbourg, France

^b Department of Botany, University of Wisconsin, Madison, Wisconsin 53706

^c Department of Genetics, University of Wisconsin, Madison, Wisconsin 53706

EXORIBONUCLEASE4 (XRN4), the *Arabidopsis thaliana* homolog of yeast XRN1, is involved in the degradation of several unstable mRNAs. Although a role for XRN4 in RNA silencing of certain transgenes has been reported, *xrn4* mutant plants were found to lack any apparent visible phenotype. Here, we show that XRN4 is allelic to the unidentified components of the ethylene response pathway *ETHYLENE-INSENSITIVE5/ACC-INSENSITIVE1 (EIN5/AIN1)* and *EIN7*. *xrn4* mutant seedlings are ethylene-insensitive as a consequence of the upregulation of *EIN3 BINDING F-BOX PROTEIN1 (EBF1)* and *EBF2* mRNA levels, which encode related F-box proteins involved in the turnover of EIN3 protein, a crucial transcriptional regulator of the ethylene response pathway. Epistasis analysis placed *XRN4/EIN5/AIN1* downstream of *CTR1* and upstream of *EBF1/2*. XRN4 does not appear to regulate ethylene signaling via an RNA-INDUCED SILENCING COMPLEX–based RNA silencing mechanism but acts by independent means. The identification of XRN4 as an integral new component in ethylene signaling adds RNA degradation as another posttranscriptional process that modulates the perception of this plant hormone.

INTRODUCTION

The gaseous hormone ethylene regulates a wide range of developmental processes in plants and their response to stress and pathogens (Johnson and Ecker, 1998). Moreover, manipulation of the ethylene response pathway is of agronomic importance given its role in fruit ripening and floral abscission. Current knowledge of this signaling pathway is based on the extensive characterization of *Arabidopsis thaliana* mutants with altered ethylene responses (Alonso and Stepanova, 2004) (see Supplemental Figure 1 online). Ethylene signal transduction begins with ethylene binding to and inactivating a family of ethylene receptors. In the absence of ethylene, these receptors activate CONSTITUTIVE TRIPLE RESPONSE1 (CTR1), a mitogen-activating protein kinase kinase kinase (MAPKKK) that negatively regulates the pathway (Kieber et al., 1993). After CTR1 inactivation, ETHYLENE-INSENSITIVE2 (EIN2), a positive regulator of the signaling cascade, which shares homology with the N-Ramp family of metal transporters (Alonso et al., 1999), promotes ethylene responses via the downstream transcription factor EIN3 and most likely also via EIN3-like1 (EIL1) and other EILs. EIN3 and EILs then upregulate primary target genes of the ethylene transcriptional cascade.

EIN3 (and possibly EILs) is regulated at the posttranslational level by the two closely related F-box proteins EIN3 BINDING F-BOX PROTEIN1 (EBF1) and EBF2 (for EIN3 binding F-BOX PROTEIN1 and -2), which are the substrate binding subunits of ubiquitin-ligating SCF complexes (Guo and Ecker, 2003; Potuschak et al., 2003; Gagne et al., 2004). In the absence of ethylene, the EIN3 protein is constitutively ubiquitinated and degraded in an EBF1/2-dependent manner. However, in the presence of ethylene, or in *ebf1 ebf2* double mutant plants, EIN3 is stabilized and accumulates (Guo and Ecker, 2003; Potuschak et al., 2003; Yanagisawa et al., 2003; Gagne et al., 2004). Conversely, plants that ectopically overexpress EBF1 or EBF2 are less sensitive to ethylene because of increased degradation of EIN3 protein (Guo and Ecker, 2003; Potuschak et al., 2003). We previously showed that the accumulation of EIN3 regulates *EBF1* and *EBF2* mRNA levels, which presumably provides a feedback mechanism to limit the accumulation of EIN3 (Potuschak et al., 2003). Consistent with this mechanism, the transcript levels of *EBF2* and, to a lesser extent, *EBF1*, show a pattern of ethylene-dependent accumulation that is disrupted in an *ein3* mutant background (Potuschak et al., 2003).

At the protein level, ethylene-dependent accumulation of EIN3 protein is completely abolished in *ein2* mutant plants and reduced in *ein5* and *ein6* mutant plants, suggesting that these genes act upstream of EIN3 in the ethylene response pathway (Guo and Ecker, 2003). *EIN5*, also known as *AIN1* (for ACC-INSENSITIVE1), and *EIN6* (Van Der Straeten et al., 1993; Roman et al., 1995; Smalle et al., 1997) are recessive ethylene-insensitive mutants whose corresponding genes still remain unknown. Because these loci have been tentatively placed upstream of *EIN3* in the ethylene response pathway, we sought to investigate

¹To whom correspondence should be addressed. E-mail pascal.genschik@ibmp-ulp.u-strasbg.fr; fax 33-03-88-61-4442.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Pascal Genschik (pascal.genschik@ibmp-ulp.u-strasbg.fr).

 Online version contains Web-only data.

 Open Access articles can be viewed online without a subscription. www.plantcell.org/cgi/doi/10.1105/tpc.106.046508

their role in the EBF1/2-dependent turnover of EIN3 protein. Here, we show that *EIN5* is allelic to the cytoplasmic *EXORIBONUCLEASE4* (*XRN4*), encoding a homolog of XRN1 of budding yeast. Epistasis analysis places *EIN5* downstream of *CTR1* and upstream of *EBF1/2*. Unlike other ethylene-insensitive mutants, which accumulate fewer *EBF1/2* mRNAs, presumably because of less EIN3 protein, *ein5-1* mutant plants accumulate higher levels of *EBF1* and *EBF2* mRNAs. *ein5-1* seedlings have concomitantly reduced levels of EIN3 protein and are less ethylene-sensitive, which are both reminiscent of EBF1-overexpressing plants. XRN4 was recently implicated in the microRNA (miRNA)-mediated turnover of some *Arabidopsis* transcripts (Souret et al., 2004) and in the small interfering RNA (siRNA)-mediated turnover of transgene-derived transcripts (Gazzani et al., 2004). With respect to ethylene perception, XRN4 does not affect RNA-INDUCED SILENCING COMPLEX (RISC)-based silencing pathways but likely acts by independent means.

RESULTS

EBF1 and EBF2 Act Downstream of EIN5, and Their Transcript Levels Are Upregulated in an *ein5* Mutant Background

Our previous analysis of the transcriptional accumulation of *EBF1/2* in ethylene mutants, which led to the establishment of the EIN3-EBF1/2 feedback loop (Potuschak et al., 2003), did not include the *ein5* mutant. Similar to other ethylene-insensitive mutants (Potuschak et al., 2003), we expected to find a downregulation of *EBF1/2* transcripts in *ein5-1* seedlings. Instead, we found that *EBF1* and *EBF2* transcript levels were moderately increased in the *ein5-1* background compared with wild-type *Arabidopsis* (Figures 1A and 1B). Coupled with the reduced and delayed accumulation of EIN3 protein in the *ein5-1* background, as described by Guo and Ecker (2003), this observation led us to speculate that EIN5 decreases EBF1/2 levels, which in turn promotes ethylene response by decreasing EIN3 protein degradation. In support of this notion, ectopic overexpression of EBF1 in wild-type *Arabidopsis* plants was shown previously to reduce ethylene sensitivity (Guo and Ecker, 2003; Potuschak et al., 2003). Under this scenario, EIN5 should act upstream of EBF1/2 in the ethylene response pathway. To test this possibility, we generated the triple mutant combination *ein5-1 ebf1-1 ebf2-1* (Figures 1C and 1D). We used here the hypomorphic *ebf2-1* allele (Potuschak et al., 2003), as complete loss-of-function mutants of both *EBF1* and *EBF2* exhibit a substantial growth arrest (Gagne et al., 2004). The *ein5-1 ebf1-1 ebf2-1* plants displayed a constitutive ethylene response, also named a triple response, because it consists of three features: a short and thick hypocotyl, an exaggerated apical hook, and a short root. This phenotype (Figures 1C and 1D) is nearly identical to that of the *ebf1-1 ebf2-1* double mutants when germinated in the dark, indicating that *EIN5* acts upstream of *EBF1* and *EBF2*. Adult *ein5-1 ebf1-1 ebf2-1* plants also showed a severe constitutive ethylene response phenotype like that of the *ebf1-1 ebf2-1* double mutant (Potuschak et al., 2003) when grown under normal light condi-

tions (Figure 1E). In addition, *ein5-1 ebf1-1 ebf2-1* plants show delayed bolting compared with the already late-flowering *ebf1-1 ebf2-1*. It is worth noting that *ACC-insensitive1* (*ain1*)/*ein5* plants were found to have a mild late-flowering phenotype (Van Der Straeten et al., 1993). The additive effect of *ein5-1* and *ebf1 ebf2* on this aspect of plant development suggests that some roles of EIN5 appear to be independent of EBF1/2.

Roman et al. (1995) previously reported that *EIN5* acts downstream of *CTR1* in the ethylene signaling pathway. We confirmed this position by recreating *ctr1-1 ein5-1* mutant plants (Figures 1F and 1G). Indeed, *EIN5* is epistatic to *CTR1*, as the *ein5-1* mutation suppresses the constitutive triple response of *ctr1* seedlings. In conclusion, *EIN5* acts downstream of *CTR1* but upstream of *EBF1/2*.

EIN5 Is Allelic to the Exoribonuclease *XRN4*

A recent microarray analysis by Souret et al. (2004) identified the *EBF1* mRNA (designated FBL6) among a small set of transcripts that are possible targets of the *Arabidopsis* XRN4 exoribonuclease. *xrn4* knockout mutants were shown to accumulate slightly higher levels of *EBF1/FBL6* mRNA as well as a shorter RNA fragment that might be a cleavage product. In striking contrast with comparable exoribonuclease mutants in other model organisms, no obvious phenotype was found for *xrn4*-deficient plants (Souret et al., 2004). Because *ein5* mutant plants also upregulate *EBF1* transcript levels and are phenotypically normal in the absence of ethylene, we compared *EBF1* mRNA accumulation in *ein5* and *xrn4* plants (Figures 1A and 1B). Both lines showed a similar upregulation of the *EBF1* transcript as well as an upregulation of the *EBF2* transcript. To investigate whether this upregulation of *EBF1/2* mRNAs in an *xrn4-3* background might affect ethylene signaling, we germinated the *xrn4-3* mutant in the dark on medium containing the ethylene precursor 1-aminocyclopropane-1-carboxylic acid (ACC) (Figures 1F and 1G). Indeed, *xrn4-3* plants were ethylene-insensitive to a similar level as *ein5-1* plants. Moreover, like *ein5-1 ctr1* plants, *xrn4-3 ctr1-1* double mutant plants were ethylene-insensitive.

Because *EIN5* was mapped like *XRN4* to chromosome 1 in *Arabidopsis*, we speculated that *EIN5* and *XRN4* are allelic. This was indeed the case (Figures 2A to 2C), as F1 plants derived from a cross between the two homozygous recessive parents *ein5-1* and *xrn4-3* were also ethylene-insensitive. Sequencing of the *XRN4* open reading frame (ORF) from the *ein5-1* background revealed a 1-bp deletion at nucleotide position 1658 causing a frame-shift mutation (Figure 2D). This mutant should generate a truncated protein of 552 residues plus 13 additional residues before termination of the protein if translated. It is noteworthy that *XRN4* transcripts are downregulated in *ein5-1* and truncated and downregulated in *xrn4-3* alleles (Figure 1A), with the latter suggesting that nonsense-mediated mRNA decay subsequently breaks down the *XRN4* transcript.

ein7 is a poorly characterized ethylene-insensitive mutant that has been proposed to be allelic to *ein5* (Roman et al., 1995). However, to date, allelism between *EIN5* and *EIN7* has not been described, presumably because of the reported semidominant nature of the *ein7* locus. As with *ein5-1* and *xrn4-3*, *ein7* seedlings




Figure 1. *EBF1* and *EBF2* Transcript Levels Are Upregulated in *ein5* Mutant Plants, and Epistasis Analysis of *ein5*.

(A) RNA gel analysis of *EBF1* and *EBF2* transcript accumulation in ecotype Columbia (Col-0), *ein5-1*, and *xrn4-3*. EtBr, ethidium bromide.
 (B) Relative *EBF1* (hatched bars) and *EBF2* (dotted bars) transcript accumulation in Col-0, *ein5-1*, and *xrn4-3* by quantification using a phosphor imager.
 (C) Phenotypes of 3-d-old etiolated seedlings of the indicated genotypes grown without ACC.
 (D) Hypocotyl (light gray) and root (dark gray) length measurements of 3-d-old dark-grown Col-0, *ebf1-1 ebf2-1*, *ein5-1*, and *ein5-1 ebf1-1 ebf2-1* seedlings germinated in the absence of ACC. Values shown are average lengths (means \pm SE) of >10 hypocotyls or roots.
 (E) Phenotypes of mature *ebf1-1 ebf2-1*, *ein5-1 ebf1-1 ebf2-1*, and wild-type (Col-0) plants grown on soil. *ein5-1* plants are similar to wild-type plants on soil and therefore are not shown.
 (F) Phenotypes of 3-d-old etiolated seedlings of the indicated genotypes grown on Murashige and Skoog (MS) medium supplemented with 10 μ M ACC.
 (G) Hypocotyl (light gray) and root (dark gray) length measurements of 3-d-old dark-grown Col-0, *ein5-1*, *xrn4-3*, *ctr1-1*, *ctr1-1 ein5-1*, and *ctr1-1 xrn4-3* seedlings in the presence of 10 μ M ACC. Values shown are average lengths (means \pm SE) of >10 hypocotyls or roots.

have a moderately increased level of *EBF1* transcript levels compared with the wild type that was maintained after ACC treatment (Figure 2E). Consistent with these results, the accumulation of EIN3 protein after ACC treatment was also attenuated in the *ein7* background (Figure 2F). We obtained comparable data with *ein5-1* and *xrn4-3* (see Supplemental Figure 2 online). Thus, *ein7* seedlings, like *ein5-1* (Guo and Ecker, 2003), were able to accumulate EIN3 after ethylene stimulation, but to a lower level

than wild-type seedlings. We sequenced the *XRN4* ORF from the *ein7* background and identified a 1-bp deletion causing a frame shift in the fifth exon at nucleotide 671. This change should lead to a truncation of the *XRN4* peptide after residue 223 and create a short tail of 13 residues before termination of the protein (Figure 2D). As a result, we conclude that *ein7*, like *ein5-1*, disrupts production of the exoribonuclease *XRN4*. Because *XRN4* acts downstream of *CTR1*, we tested whether *XRN4* is




Figure 2. *ein5* and *ein7* Are Mutants in the Exoribonuclease *XRN4*.

(A) Phenotypes of 4-d-old etiolated seedlings of the indicated genotypes grown on MS medium.

(B) Hypocotyl (light gray) and root (dark gray) length measurements of 4-d-old dark-grown Col-0, *xrn4-3*, and *ein5-1* seedlings germinated in the absence of ACC. Values shown are average lengths (means \pm SE) of >10 hypocotyls or roots.

(C) Hypocotyl (light gray) and root (dark gray) length measurements of 4-d-old dark-grown Col-0, *xrn4-3*, *ein5-1*, and F1 seedlings of a *xrn4-3* \times *ein5-1* cross in the presence of 10 μ M ACC. Values shown are average lengths (means \pm SE) of >10 hypocotyls or roots.

(D) Gene structure of *XRN4*. Introns are indicated by lines. Shaded and hatched boxes indicate coding regions and 5' and 3' untranslated regions, respectively. The positions of *ein5-1* and *ein7* mutations are indicated and correspond to cytosine deletions creating truncated proteins in both cases. The position of the T-DNA of the *xrn4-3* allele is also indicated.

(E) *EBF1* transcript levels in Col-0 and *ein7* at different time points during treatment with 50 μ M ACC. RNA was extracted from 3-week-old Col-0 and *ein7* seedlings at different time points of the ACC treatment, subjected to RNA gel blot analysis, and hybridized with the indicated probes. EtBr, ethidium bromide.

(F) EIN3 protein accumulation in Col-0 and *ein7* at different time points during 50 μ M ACC treatment. Total protein extracts from the same sample as indicated in (E) were subjected to immunoblot assays.

(G) RNA gel analysis of *XRN4* transcript accumulation in different mutant backgrounds as indicated.

transcriptionally misregulated in *ctr1-1* or other mutants in the ethylene response pathway. However, *XRN4* mRNA accumulated to similar levels in several ethylene signaling mutants, indicating that if *XRN4* is regulated in an ethylene-dependent manner, this must occur at the posttranscriptional level (Figure 2G).

xrn4/ein5 Plants Are Deficient in EIN3/EIL1-Dependent Growth Inhibition by Ethylene

Earlier studies of etiolated *Arabidopsis* seedlings revealed two phases of ethylene-induced growth inhibition for wild-type Col-0 hypocotyls (Binder et al., 2004a, 2004b). The first phase of inhibition begins ~10 min after ethylene is applied. Thirty minutes after the addition of ethylene, a new steady state growth rate is reached, which is maintained for ~20 min before a second phase of growth inhibition begins, which reduces the growth rate to a lower steady state. At saturating concentrations of the hormone, this lower growth rate is maintained for as long as the hormone is present. If ethylene is removed, hypocotyl elongation recovers by ~90 min to pretreatment growth rates by a dampening oscillation. EIN2 is essential for both phases. Whereas the first phase of growth inhibition is independent of EIN3 and EIL1, the second phase response requires both of these transcription factors (Binder et al., 2004b).

Because both the *xrn4-3* and *ein5-1* mutants increase *EBF1/2* mRNA levels and accumulate lower levels of EIN3 protein in response to ethylene or its precursor ACC, we examined the ethylene growth response kinetics of these mutants. As shown in Figure 3, wild-type Col-0 hypocotyls have a growth inhibition response similar to that described in previous reports (Binder et al., 2004a, 2004b). Recovery after the removal of ethylene was slightly slower in wild-type seedlings in this study, taking ~105 min to recover to the initial, pretreatment growth rates. Both the *xrn4-3* and *ein5-1* mutants showed onset kinetics indistinguishable from those of the wild type after ethylene addition, with both phases of growth inhibition being evident (Figure 3). However, both mutants recovered markedly faster than the wild type after ethylene removal and appeared to begin growth recovery even before the ethylene was removed (Figures 3A and 3B). Experiments with *ein7* generated similar kinetics (Figure 3C). These data were consistent with our observed upregulation of *EBF1/2* RNA levels in *xrn4/ein5/ein7*-deficient plants and the subsequent reduced accumulation of EIN3 protein in these lines.

The *xrn4* Mutation Does Not Affect *EBF1/2* mRNA Turnover

Because *XRN4* functions in mRNA decay, one obvious scenario is that the upregulation of *EBF1* in the *xrn4-3* background is the consequence of increased *EBF1* transcript stability. To test this possibility, we monitored the stability of the *EBF1* and *-2* mRNAs, using the adenosine analogue cordycepin (3'-deoxyadenosine) to block transcription. Under these conditions, both mRNAs were equally short-lived in wild-type, *ein5-1*, and *ein7* plants, suggesting that the mode of action of *XRN4* must be more complex (Figure 4A; data not shown). This finding is in agreement with that described by Souret et al. (2004), who also failed to find a stabilization of *FBL6/EBF1* transcripts in *xrn4*




Figure 3. Growth Kinetics of Etiolated *Arabidopsis* Hypocotyls in Response to Ethylene.

Growth rates were recorded for 1 h in air followed by a 2-h exposure to 10 μ L/L ethylene. This was followed by 5 h in air. The responses of wild-type Col-0 hypocotyls are shown in both panels (open squares) for comparison with the homozygous mutants (closed symbols) *xrn4-3* (A), *ein5-1* (B), and *ein7* (C). All data represent averages of at least five seedlings \pm SD.

mutants. We further investigated the stability of ectopically expressed *EBF1* mRNA under the control of the cauliflower mosaic virus 35S promoter and lacking the endogenous 5' and 3' untranslated regions of the transcript. The transgenic line selected showed a higher *EBF1* mRNA level compared with the *ein5* mutant (Figure 4A) and consistently exhibited a stronger ethylene-insensitive phenotype than the latter (Figures 4B and




Figure 4. *EBF1* and *EBF2* mRNAs Remain Short-Lived in *xrn4/ein5* Mutants, and Their Turnover Is Ethylene Independent.

(A) RNA gel analysis of a time-course experiment for *EBF1* and *EBF2* transcript half-life determination in Col-0 and the *ein5-1* mutant. The blot was hybridized with either the 3' untranslated region (UTR; detecting endogenous transcripts) or ORF probes (also detecting ectopically expressed *EBF1*) as indicated, and *EBF1/2* transcript half-lives were determined relative to EF1- α , which is a stable transcript. The asterisk indicates the possible *EBF1* mRNA cleavage product as indicated by Souret et al. (2004), and the arrowhead indicates the shorter *EBF1* transcript originating from the 35S:*EBF1* transgene. EtBr, ethidium bromide.

(B) Strong *EBF1* overexpression exacerbates the ethylene-insensitive phenotype of *ein5*. An *Arabidopsis* T-DNA-transformed line that expresses a high level of *EBF1* mRNA (see [A]) exhibits strong ethylene insensitivity when germinated in the dark in the presence of 10 μ M ACC. This phenotype is maintained when the transgene is introduced into the *ein5-1* mutant.

(C) Hypocotyl (light gray) and root (dark gray) length measurements of 3-d-old dark grown Col-0, *ein5-1*, Col-0::EBF1ox, and *ein5-1*::EBF1ox seedlings in the presence of 10 μ M ACC. Values shown are average lengths (means \pm SE) of >10 hypocotyls or roots.

(D) RNA gel analysis of a time-course experiment for *EBF1* and *EBF2* endogenous transcript half-life determination in Col-0 and in *ein5-1* and *ein7* mutants pretreated with or without 50 μ M ACC. *EBF1/2* transcript half-lives were determined relative to EF1- α .

4C). Moreover, when the transgene was expressed in the *ein5* mutant background, the ethylene-insensitive phenotype of *ein5* was much stronger, further demonstrating the correlation between the *EBF1* expression levels and ethylene insensitivity. Surprisingly, the transcript produced from the synthetic construct was also short-lived in both wild-type and *ein5* plants (Figure 4A). Furthermore, when *EBF1* was overexpressed ectopically, wild-type transcript levels of *EBF1* and *EBF2* were greatly reduced, as shown previously (Potuschak et al., 2003), and were apparently less affected by the cordycepin treatment

(Figure 4A). Nevertheless, this change in the turnover is not affected by the *ein5-1* mutation.

We next asked whether ethylene might change the half-life of *EBF1* and/or *EBF2* mRNAs in *ein5/ein7* mutant backgrounds. However, *EBF1* and *EBF2* mRNAs remained short-lived in both *ein5* and *ein7* with and without preincubation with the ethylene precursor ACC (Figure 4D). From these experiments, we conclude that the higher accumulation of *EBF1* and *EBF2* mRNAs in the exoribonuclease mutants most likely is not the consequence of an altered turnover of these transcripts.

XRN4 Function in Ethylene Signaling Is Independent of Known Mutants in miRNA and siRNA Pathways

Although we cannot totally exclude a direct role of XRN4 in the turnover of *EBF1/2* mRNAs, our results suggest that XRN4 modulates EBF1/2 indirectly, either by positively regulating a repressor or by negatively regulating an activator of *EBF1/2* transcription. Such a regulation of *EBF1/2* mRNA abundance by XRN4 could occur via a miRNA- or siRNA-dependent mechanism, as several known miRNA substrates of XRN4 are miRNA targets (Souret et al., 2004) and because the suppression of SHOOT MERISTEMLESS and WUSCHEL overexpression in an *xrn4* mutant background requires *RDR6/SDE1* (for RNA-DEPENDENT RNA POLYMERASE6/SILENCING DEFECTIVE1), the RNA-dependent RNA polymerase involved in siRNA-dependent transgene silencing (Gazzani et al., 2004). Therefore, we tested mutants that are impaired in miRNA and siRNA pathways, such as *hen1*, *rdr2*, *dcl2*, *dcl3*, *sde1*, and the recently described *dcl4-1* (for *hua enhancer1*, RNA-dependent RNA polymerase2, and *dicer-like 2/3/4*) (Park et al., 2002; Boutet et al., 2003; Xie et al., 2004; Gascioli et al., 2005) for their response toward ethylene (see Supplemental Figure 3 online). None showed a significant degree of ethylene insensitivity (data not shown) or higher EBF1/2 transcript levels under the conditions tested (see Supplemental Figure 3A online). We generated *ein5 sde1* and *ein5 rdr2* double mutant plants and found that they were indistinguishable from *ein5* when germinated in the presence of ACC (see Supplemental Figures 3B and 3C online). Finally, we created an *xrn4-3 hen1* double mutant that was also ethylene-insensitive (see Supplemental Figures 3D and 3E online). The slight increase in hypocotyl length observed for this double mutant compared with the *xrn4* single mutant might be attributed to the mix of two different ecotypes (Col and Landsberg *erecta*). We conclude from these experiments that the ethylene-insensitive phenotype of *xrn4* does not require functional *HEN1*, *RDR6/SDE1*, or *RDR2*.

DISCUSSION

The control of RNA turnover is a crucial aspect of gene expression in all organisms (Parker and Song, 2004; Wilusz and Wilusz, 2004). In animal cells, XRN1, a 5' → 3' processive exoribonuclease, plays an important role in the ARE (for AU-rich element in the 3' untranslated region) mRNA degradation decay pathway (Stoecklin et al., 2006) and in nonsense-mediated mRNA decay, a pathway required to destroy aberrant RNAs harboring premature translation termination codons (Lejeune et al., 2003; Gatfield and Izaurralde, 2004). It appears to function by processively degrading RNAs from the 5' end after removal of the 5' cap, thus requiring a first cleavage event by another ribonuclease before substrate recognition. In *Arabidopsis*, the cytoplasmic XRN4 appears to represent the plant ortholog of XRN1 (Kastenmayer and Green, 2000). *Arabidopsis* also possesses two homologs of XRN4, called XRN2 and XRN3, that localize to the nucleus and therefore act on a different population of RNA substrates than XRN4 (Kastenmayer and Green, 2000). Although the function(s) of XRN4 is not completely resolved, it was recently implicated in

the degradation of several, but not all, miRNA-directed cleavage products (Souret et al., 2004). A role for XRN4 in transgene silencing via RNA interference in *Arabidopsis* has also been proposed (Gazzani et al., 2004). However, in contrast with other model organisms (Tishkoff et al., 1995; Newbury and Woollard, 2004), *Arabidopsis xrn4* null mutants are viable, with no apparent deleterious phenotype (Souret et al., 2004).

The work presented here confers a physiological function to a member of the *XRN1/XRN4* exoribonuclease family in hormone signaling. Based on our genetic analysis, *XRN4* is an integral component of the ethylene response pathway in *Arabidopsis*. It likely acts downstream of the MAPKKK *CTR1* and upstream of the F-box proteins *EBF1/2*, given the ability of *xrn4* mutants to suppress the constitutive ethylene response seen in *ctr1* but not the constitutive ethylene response phenotype of the *ebf1-1 ebf2-1* double mutants (Figure 5). We also present evidence that *XRN4* loss of function results in increased levels of EBF1 and EBF2 transcripts, which encode two F-box proteins that target EIN3 for ethylene-dependent degradation. This provides a rationale for the reduced accumulation of EIN3 protein observed in *ein5* mutant plants (Guo and Ecker, 2003). This reduced




Figure 5. Representation of the Ethylene Signaling Cascade Including XRN4/EIN5.

Ethylene is perceived by the ETHYLENE RESPONSE (ETR/ERS) receptors located in the endoplasmic reticulum membrane. Binding of ethylene to the receptors results in the inactivation of both the receptors and CTR1, thereby causing the derepression of positive regulatory factors, such as EIN2 (data not shown) and XRN4/EIN5. How CTR1 regulates EIN2 and XRN4/EIN5 and the relationship of EIN2 with XRN4/EIN5 in the ethylene signaling cascade are unknown. Strikingly, our model proposes the existence of two degradation pathways in ethylene signaling: one RNA decay pathway into the cytosol that indirectly controls the steady state levels of EBF1/2 mRNAs, which are components of a protein degradation pathway that controls EIN3 stability in the nucleus.

accumulation of EIN3 also explains the faster recovery in growth after the removal of ethylene that we observed. It is worth noting that *ain1/ein5* mutant plants have no obvious signaling defects for other hormones (Van Der Straeten et al., 1993), suggesting that XRN4/EIN5 has a specific role in ethylene signaling.

The molecular functions of several components in the ethylene signaling pathway remain rather mysterious (Alonso and Stepanova, 2004). Thus, after the ethylene signal has been transmitted from the receptors to the Raf-like protein kinase CTR1, there are few insights into the molecular mechanism that leads to the nuclear accumulation of the EIN3 transcription factor. In particular, the MAPK module predicted to be downstream of CTR1 still remains to be characterized. Moreover, neither the function nor the subcellular location of EIN2, which is known to act downstream of CTR1 but upstream of EBF1/2, has been solved. This work adds another level of complexity to this pathway with the identification of a component, XRN4, that is involved in RNA decay. XRN4 is known to act in the cytosol to regulate the transcript levels of EBF1 and EBF2. These two F-box proteins operate in a proteolytic degradation pathway in the nucleus (Figure 5). Although both EIN2 and XRN4/EIN5 are epistatic to CTR1, it is unknown whether they act in the same or independent pathways. Interestingly, it was recently found that EIN2, but not EIN5, is required for insect resistance; by contrast, EIN5, but not EIN2, was required for the growth enhancement induced by bacterial harpin proteins (Dong et al., 2004). Thus, EIN2 and XRN4/EIN5 might act independently in at least some pathways, although they are both necessary for an efficient EIN3 accumulation after ethylene perception (Guo and Ecker, 2003).

One attractive hypothesis was that EBF1/2 mRNAs are directly targeted by a miRNA or a siRNA and that this mechanism requires XRN4 for the initial cleavage and the turnover of the cleavage products. However, several lines of evidence do not support such a model. First, EBF1/2 remain short-lived mRNA species in *xrn4/ein5* mutants (Souret et al., 2004; this work). Second, exhaustive sequencing of small RNAs in *Arabidopsis* failed to identify sequences matching EBF1 or EBF2 mRNAs (Lu et al., 2005); in particular, none was found to fit the predicted EBF1 cleavage site (Souret et al., 2004). Third, previous extensive screens failed to detect ethylene-insensitive mutants that fall into the miRNA, siRNA, and transacting small interfering RNA classes of mutants. Indeed, the mutants that we tested (*hen1*, *rdr2*, *sde1*, *dcl2*, and *dcl3*) and that affect these different RNA silencing pathways (Brodersen and Voinnet, 2006) show a normal accumulation of EBF1/2 transcripts. Nevertheless, even if these mutants do not result in ethylene insensitivity, it is still possible that one of them is required for the ethylene-insensitive phenotype seen in *xrn4/ein5* mutant plants. Such a scenario would be reminiscent of the suppressor phenotype of *xrn4* mutations, which acts through siRNA-dependent silencing of a reporter gene and requires the RNA-dependent RNA polymerase SDE1 (Gazzani et al., 2004). A similar pathway could act on an upstream regulator of EBF1/2, but not directly on EBF1/2 (or cauliflower mosaic virus 35S promoter-driven EBF1), as they are not silenced in an *xrn4/ein5* background. Nevertheless, this also seems unlikely, as the *hen1 xrn4-3*, *ein5-1 sde1*, and *ein5-1 rdr2* mutants generated were found to have an *ein5*-like ethylene

response. Thus, we conclude that XRN4 indirectly regulates EBF1/2 mRNA levels by a mechanism that does not seem to require RISC-based RNA silencing pathways.

Interestingly, proteins involved in RNA metabolism are increasingly being shown to be important in plant hormone regulation (Fedoroff, 2002). The *Arabidopsis* HYPONASTIC LEAVES1 gene encodes a nuclear double-stranded RNA binding protein, and its loss of function increases plant sensitivity to abscisic acid but reduces the sensitivity to auxin and cytokinin (Lu and Fedoroff, 2000). Other RNA binding proteins that affect abscisic acid signaling are the Sm-like protein SUPERSENSITIVE TO ABA AND DRAUGHT (Xiong et al., 2001) and the mRNA cap binding protein ABA HYPERSENSITIVE1 (Hugouvieux et al., 2001). The cloning of the abscisic acid-hypersensitive mutant *ABA-hypersensitive germination2* revealed a gene encoding a poly(A)-specific ribonuclease supposed to function in mRNA decay (Nishimura et al., 2005). Future analyses will certainly reveal the mechanisms for how these RNA binding and processing enzymes control plant hormonal pathways and also identify their target RNAs.

METHODS

Arabidopsis Lines and Plant Crosses

All *Arabidopsis thaliana* lines used have been described elsewhere: *ago4-1* (Zilberman et al., 2003); *ctr1-1* (Kieber et al., 1993); *ein5-1* and *ein7* (Roman et al., 1995); *hen1* (Chen et al., 2002); *rdr2*, *dcl2-1*, and *dcl3-1* (Xie et al., 2004); *dcl4-1* (Gascioli et al., 2005); *ebf1-1* and *ebf2-1* (Potuschak et al., 2003); *sde1* (Dalmay et al., 2000); *xrn4-3* (SALK_014209) (Gazzani et al., 2004). Note that EIN5 has been shown to be allelic to AIN1 and that *ein5-1* has been renamed *ain1-10* (Smalle et al., 1997). However, the bulk of the current literature still uses the name EIN5; therefore, we continue to use this name to avoid confusion.

For the generation of *ein5-1 ebf1-1 ebf2-1* triple mutant plants, *ebf1-1 ebf2-1* plants were pollinated with *ein5-1*-derived pollen. In the F2 generation, *ein5-1* homozygote lines were selected based on their ethylene-insensitive phenotype and tested by PCR for the *ebf1-1* and *ebf2-1* T-DNA insertions. No plants showing an *ein5-1*-like ethylene-insensitive phenotype were found to be also double homozygous for *ebf1-1* and *ebf2-1*. Plants that were *ein5-1* homozygous and double heterozygous for the *ebf1-1* and *ebf2-1* insertion were allowed to self, and F3 seeds were tested for their seedling phenotype. *ein5-1 ebf1-1 ebf2-1* triple mutant plants were segregating at the expected ratio and confirmed by PCR testing. For the generation of *ein5-1 ctr1-1* and *xrn4-3 ctr1* double mutant plants, *ctr1* plants were pollinated with pollen from *ein5-1* or *xrn4-3* donors; *ctr1* homozygotes were then selected in the F2 generation based on their phenotype and genotyped for the *ein5-1* mutation and *xrn4-3* T-DNA insertions. No *ctr1 ein5* or *ctr1 xrn4-3* homozygotes were identified that way, but several *ctr1* plants were found to be heterozygote for the *ein5-1* mutation or the *xrn4-3* T-DNA insertion. The plants were allowed to set seeds, and they segregated ethylene-insensitive plants in the next generation that were found to be double homozygotes. For the generation of *ein5-1 sde1*, *ein5-1 rdr2-1*, and *xrn4-3 hen1* plants, F2 seedlings were genotyped directly and double homozygotes were allowed to set seeds. Ethylene responses of double homozygotes were measured with F3 and F4 seeds.

Sequencing of the XRN4 ORF from *ein5-1* and *ein7*

Sequencing of RT-PCR amplification products of the XRN4 ORF from *ein5-1* and *ein7* material identified a 1-bp deletion at position 1658

and 671 for *ein5-1* and *ein7*, respectively. In the case of *ein5-1*, the deletion causes the truncation of the XRN4 protein sequence after amino acid 552 and leads to the addition of 13 amino acids (RYLNFTLLLSLT) before a premature stop codon. In the case of *ein7*, the deletion leads to the termination of the XRN4 protein sequence after amino acid 223 and the addition of the short peptide QIHGIVYMWVWQI before a premature stop codon. To confirm the sequencing results, PCR markers for *ein5-1* and *ein7* were designed and tested with genomic DNA. For detection of the *ein5-1* mutation, a derived cleaved amplified polymorphic sequence marker was designed using the primers *ein5F* (5'-GTTGATGACTGATCCCTCATCCT-3') and *ein5R* (5'-GAGTGCAACTATCCAGCATGAA-3'). *ein5-1*-specific PCR products were cleaved by Taq1, whereas wild-type specific PCR products were not cleaved. For detection of the *ein7* mutation, a cleaved-amplified polymorphic sequence marker was designed with the primers *ein7F* (5'-TTCAAATGTTCCGGGAGAAG-3') and *ein7R* (5'-GACGAAGCACCAACACCTTA-3'). *ein7*-derived PCR products were cleavable with the restriction enzyme *BclI*. PCR products derived from wild-type DNA were not cleaved.

Seedling Responses to Ethylene, and Growth Rate Measurements on Hypocotyls

Seedling responses to ethylene were measured exactly as described previously (Potuschak et al., 2003).

The effect of ethylene on the growth rate of hypocotyls was measured using etiolated *Arabidopsis* seedlings as described previously (Binder et al., 2004a, 2004b). Mutant and wild-type seeds were surface-sterilized by treatment with 70% ethanol for ~30 s, placed on sterile filter paper to dry, and plated on half-strength MS medium, pH 5.7, containing 0.8% agar and B5 vitamins, consisting of inositol (100 mg/mL), nicotinic acid (1 mg/mL), pyridoxin HCl (1 mg/mL), and thiamine HCl (10 mg/mL) with no added sugar. Five micromolar L- α -(2-aminoethoxyvinyl)-glycine was included to inhibit the biosynthesis of ethylene.

Seeds were cold-treated for 2 to 4 d at 4°C and exposed to light for 2 to 8 h before being grown vertically in the dark for 2 d at 22°C. Growth rate measurements were performed as described (Binder et al., 2004b). After 1 h of treatment with air to establish a basal growth rate, ethylene was introduced at a flow rate of 10 mL/min, giving a final concentration of 10 μ L/L. Ethylene was removed 2 h later. Gas flow was maintained at 100 mL/min using Side-Trak mass flow meters and controller (Sierra Instruments). Hypocotyl growth was measured from digital images that were captured every 5 min for 7 h with either an EDC-1000N CCD (Electrim) or an Infinity 2-1M camera (Luminera) and light provided by an infrared light-emitting diode. Growth rates were calculated using custom software generated by Edgar Spalding in LabVIEW 5.0 (National Instruments) as described previously (Parks and Spalding, 1999; Folta and Spalding, 2001). All data presented represent averages of at least five seedlings from a minimum of three separate experiments.

Plasmid Construction and Transformation

For the chimeric EBF1 construct, the ORF of EBF1 was modified by PCR and inserted via *NcoI* into the multiple cloning site of a modified pPily (Farras et al., 2001), which carried an additional six hemagglutinin epitope tags to facilitate protein detection. The expression cassette containing the 2 \times cauliflower mosaic virus 35S promoter, the epitope-tagged EBF1 ORF, and the nopaline synthase terminator was excised via *NotI* cleavage and inserted in a *XmaI*-digested pCambia1380 binary vector (www.cambia.org). *NotI* and *XmaI* restriction sites were partially filled in before ligation. After transformation of Col-0 by flower dip (Clough and Bent, 1998), homozygote lines containing single inserts were selected. One such line was crossed into *ein5-1*. Again, plants homozygous for *ein5-1* and the T-DNA insertion were selected.

RNA Gel Analyses and Transcript Half-Life Measurements

Unless stated otherwise, RNA was extracted from 3-week-old light-grown seedlings. RNA preparation, RNA gel blotting, and hybridization were performed using standard protocols. Quantification of RNA gel blots were performed with a Fuji BAS 1000 Imager and MacBAS software version 2.1. For the estimation of half-life measurements with cordycepin, the protocol of Seeley et al. (1992) with modifications for *Arabidopsis* (Gutierrez et al., 2002) was followed. To test the influence of ACC on the stability of mRNAs, 50 μ M ACC was added to the plant material submerged in the cordycepin incubation buffer (Seeley et al., 1992) 30 min before the addition of cordycepin. To test the influence of ACC on EIN3 protein accumulation or EBF1/2 transcript levels, 3-week-old light-grown plants were submerged into half-strength MS buffer (1% sucrose) and treated with 50 μ M ACC for the indicated period of time.

Protein Gel Blots

Samples of 15 μ g of proteins were separated on SDS gels and blotted onto Immobilon-P membranes (Millipore). The probing procedure and antibodies against EIN3 have been described (Guo and Ecker, 2003; Yanagisawa et al., 2003).

Supplemental Data

The following materials are available in the online version of this article

Supplemental Figure 1. Flow Chart of the Ethylene Response Pathway.

Supplemental Figure 2. Reduced EIN3 Protein Accumulation in *ein5-1* and *xrn4-3* Plants in Response to ACC Treatment.

Supplemental Figure 3. Mutants Deficient in miRNA, siRNA, and Transacting Small Interfering RNA Pathways Do Not Exhibit Higher *EBF1* and *EBF2* mRNA Levels and Do Not Significantly Alter the Ethylene-Insensitive Phenotype of *ein5*.

ACKNOWLEDGMENTS

We thank Olivier Voinnet and members of his laboratory for insightful discussions and the gift of *Arabidopsis* strains and Patrick Achard for discussions and critically reading the manuscript. We thank Hongwei Guo and Joseph R. Ecker for an antibody against EIN3. This work was supported by the Centre National de la Recherche Scientifique and a grant from the French Ministry of Research (Grant ACI 2004 No. BCMS 167) to P.G. and by a grant from the U.S. National Science Foundation *Arabidopsis* 2010 Program (Grant MCB-0115870) to R.D.V.

Received August 8, 2006; revised September 5, 2006; accepted October 13, 2006; published November 3, 2006.

REFERENCES

- Alonso, J.M., Hirayama, T., Roman, G., Nourizadeh, S., and Ecker, J.R. (1999). EIN2, a bifunctional transducer of ethylene and stress responses in *Arabidopsis*. *Science* **284**, 2148–2152.
- Alonso, J.M., and Stepanova, A.N. (2004). The ethylene signaling pathway. *Science* **306**, 1513–1515.
- Binder, B.M., Mortimore, L.A., Stepanova, A.N., Ecker, J.R., and Bleecker, A.B. (2004a). Short term growth responses to ethylene in *Arabidopsis* seedlings are EIN3/EIL1 independent. *Plant Physiol.* **136**, 2921–2927.

- Binder, B.M., O'Malley, R.C., Wang, W., Moore, J.M., Parks, B.M., Spalding, E.P., and Bleeker, A.B.** (2004b). Arabidopsis seedling growth response and recovery to ethylene: A kinetic analysis. *Plant Physiol.* **136**, 2913–2920.
- Boutet, S., Vazquez, F., Liu, J., Beclin, C., Fagard, M., Gratiyas, A., Morel, J.B., Crete, P., Chen, X., and Vaucheret, H.** (2003). Arabidopsis HEN1: A genetic link between endogenous miRNA controlling development and siRNA controlling transgene silencing and virus resistance. *Curr. Biol.* **13**, 843–848.
- Brodersen, P., and Voinnet, O.** (2006). The diversity of RNA silencing pathways in plants. *Trends Genet.* **22**, 268–280.
- Chen, X., Liu, J., Cheng, Y., and Jia, D.** (2002). HEN1 functions pleiotropically in Arabidopsis development and acts in C function in the flower. *Development* **129**, 1085–1094.
- Clough, S.J., and Bent, A.F.** (1998). Floral dip: A simplified method for *Agrobacterium*-mediated transformation of *Arabidopsis thaliana*. *Plant J.* **16**, 735–743.
- Dalmay, T., Hamilton, A., Rudd, S., Angell, S., and Baulcombe, D.C.** (2000). An RNA-dependent RNA polymerase gene in Arabidopsis is required for posttranscriptional gene silencing mediated by a transgene but not by a virus. *Cell* **101**, 543–553.
- Dong, H.P., Peng, J., Bao, Z., Meng, X., Bonasera, J.M., Chen, G., Beer, S.V., and Dong, H.** (2004). Downstream divergence of the ethylene signaling pathway for harpin-stimulated Arabidopsis growth and insect defense. *Plant Physiol.* **136**, 3628–3638.
- Farras, R., Ferrando, A., Jasik, J., Kleinow, T., Okresz, L., Tiburcio, A., Salchert, K., del Pozo, C., Schell, J., and Koncz, C.** (2001). SKP1-SnRK protein kinase interactions mediate proteasomal binding of a plant SCF ubiquitin ligase. *EMBO J.* **20**, 2742–2756.
- Fedoroff, N.V.** (2002). RNA-binding proteins in plants: The tip of an iceberg? *Curr. Opin. Plant Biol.* **5**, 452–459.
- Folta, K.M., and Spalding, E.P.** (2001). Unexpected roles for cryptochrome 2 and phototropin revealed by high-resolution analysis of blue light-mediated hypocotyl growth inhibition. *Plant J.* **26**, 471–478.
- Gagne, J.M., Smalle, J., Gingerich, D.J., Walker, J.M., Yoo, S.D., Yanagisawa, S., and Vierstra, R.D.** (2004). Arabidopsis EIN3-binding F-box 1 and 2 form ubiquitin-protein ligases that repress ethylene action and promote growth by directing EIN3 degradation. *Proc. Natl. Acad. Sci. USA* **101**, 6803–6808.
- Gascioli, V., Mallory, A.C., Bartel, D.P., and Vaucheret, H.** (2005). Partially redundant functions of Arabidopsis DICER-like enzymes and a role for DCL4 in producing trans-acting siRNAs. *Curr. Biol.* **15**, 1494–1500.
- Gatfield, D., and Izaurralde, E.** (2004). Nonsense-mediated messenger RNA decay is initiated by endonucleolytic cleavage in *Drosophila*. *Nature* **429**, 575–578.
- Gazzani, S., Lawrenson, T., Woodward, C., Headon, D., and Sablowski, R.** (2004). A link between mRNA turnover and RNA interference in Arabidopsis. *Science* **306**, 1046–1048.
- Guo, H., and Ecker, J.R.** (2003). Plant responses to ethylene gas are mediated by SCF(EBF1/EBF2)-dependent proteolysis of EIN3 transcription factor. *Cell* **115**, 667–677.
- Gutierrez, R.A., Ewing, R.M., Cherry, J.M., and Green, P.J.** (2002). Identification of unstable transcripts in Arabidopsis by cDNA microarray analysis: Rapid decay is associated with a group of touch- and specific clock-controlled genes. *Proc. Natl. Acad. Sci. USA* **99**, 11513–11518.
- Hugouvieux, V., Kwak, J.M., and Schroeder, J.I.** (2001). An mRNA cap binding protein, ABH1, modulates early abscisic acid signal transduction in Arabidopsis. *Cell* **106**, 477–487.
- Johnson, P.R., and Ecker, J.R.** (1998). The ethylene gas signal transduction pathway: a molecular perspective. *Annu. Rev. Genet.* **32**, 227–254.
- Kastenmayer, J.P., and Green, P.J.** (2000). Novel features of the XRN-family in Arabidopsis: Evidence that AtXRN4, one of several orthologs of nuclear Xrn2p/Rat1p, functions in the cytoplasm. *Proc. Natl. Acad. Sci. USA* **97**, 13985–13990.
- Kieber, J.J., Rothenberg, M., Roman, G., Feldmann, K.A., and Ecker, J.R.** (1993). CTR1, a negative regulator of the ethylene response pathway in Arabidopsis, encodes a member of the raf family of protein kinases. *Cell* **72**, 427–441.
- Lejeune, F., Li, X., and Maquat, L.E.** (2003). Nonsense-mediated mRNA decay in mammalian cells involves decapping, deadenylation, and exonucleolytic activities. *Mol. Cell* **12**, 675–687.
- Lu, C., and Fedoroff, N.** (2000). A mutation in the Arabidopsis HYL1 gene encoding a dsRNA binding protein affects responses to abscisic acid, auxin, and cytokinin. *Plant Cell* **12**, 2351–2366.
- Lu, C., Tej, S.S., Luo, S., Haudenschild, C.D., Meyers, B.C., and Green, P.J.** (2005). Elucidation of the small RNA component of the transcriptome. *Science* **309**, 1567–1569.
- Newbury, S., and Woollard, A.** (2004). The 5'-3' exoribonuclease xrn-1 is essential for ventral epithelial enclosure during *C. elegans* embryogenesis. *RNA* **10**, 59–65.
- Nishimura, N., Kitahata, N., Seki, M., Narusaka, Y., Narusaka, M., Kuromori, T., Asami, T., Shinozaki, K., and Hirayama, T.** (2005). Analysis of ABA hypersensitive germination2 revealed the pivotal functions of PARN in stress response in Arabidopsis. *Plant J.* **44**, 972–984.
- Park, W., Li, J., Song, R., Messing, J., and Chen, X.** (2002). CARPEL FACTORY, a Dicer homolog, and HEN1, a novel protein, act in microRNA metabolism in Arabidopsis thaliana. *Curr. Biol.* **12**, 1484–1495.
- Parker, R., and Song, H.** (2004). The enzymes and control of eukaryotic mRNA turnover. *Nat. Struct. Mol. Biol.* **11**, 121–127.
- Parks, B.M., and Spalding, E.P.** (1999). Sequential and coordinated action of phytochromes A and B during Arabidopsis stem growth revealed by kinetic analysis. *Proc. Natl. Acad. Sci. USA* **96**, 14142–14146.
- Potuschak, T., Lechner, E., Parmentier, Y., Yanagisawa, S., Grava, S., Koncz, C., and Genschik, P.** (2003). EIN3-dependent regulation of plant ethylene hormone signaling by two Arabidopsis F box proteins: EBF1 and EBF2. *Cell* **115**, 679–689.
- Roman, G., Lubarsky, B., Kieber, J.J., Rothenberg, M., and Ecker, J.R.** (1995). Genetic analysis of ethylene signal transduction in Arabidopsis thaliana: Five novel mutant loci integrated into a stress response pathway. *Genetics* **139**, 1393–1409.
- Seeley, K.A., Byrne, D.H., and Colbert, J.T.** (1992). Red light-independent instability of oat phytochrome mRNA in vivo. *Plant Cell* **4**, 29–38.
- Smalle, J., Haegman, M., Kurepa, J., Van Montagu, M., and Straeten, D.V.** (1997). Ethylene can stimulate Arabidopsis hypocotyl elongation in the light. *Proc. Natl. Acad. Sci. USA* **94**, 2756–2761.
- Souret, F.F., Kastenmayer, J.P., and Green, P.J.** (2004). AtXRN4 degrades mRNA in Arabidopsis and its substrates include selected miRNA targets. *Mol. Cell* **15**, 173–183.
- Stoecklin, G., Mayo, T., and Anderson, P.** (2006). ARE-mRNA degradation requires the 5'-3' decay pathway. *EMBO Rep.* **7**, 72–77.
- Tishkoff, D.X., Rockmill, B., Roeder, G.S., and Kolodner, R.D.** (1995). The sep1 mutant of *Saccharomyces cerevisiae* arrests in pachytene and is deficient in meiotic recombination. *Genetics* **139**, 495–509.
- Van Der Straeten, D., Djudzman, A., Van Caeneghem, W., Smalle, J., and Van Montagu, M.** (1993). Genetic and physiological analysis of a new locus in Arabidopsis that confers resistance to 1-aminocyclopropane-1-carboxylic acid and ethylene and specifically affects the ethylene signal transduction pathway. *Plant Physiol.* **102**, 401–408.

- Wilusz, C.J., and Wilusz, J.** (2004). Bringing the role of mRNA decay in the control of gene expression into focus. *Trends Genet.* **20**, 491–497.
- Xie, Z., Johansen, L.K., Gustafson, A.M., Kasschau, K.D., Lellis, A.D., Zilberman, D., Jacobsen, S.E., and Carrington, J.C.** (2004). Genetic and functional diversification of small RNA pathways in plants. *PLoS Biol.* **2**, E104.
- Xiong, L., Gong, Z., Rock, C.D., Subramanian, S., Guo, Y., Xu, W., Galbraith, D., and Zhu, J.K.** (2001). Modulation of abscisic acid signal transduction and biosynthesis by an Sm-like protein in *Arabidopsis*. *Dev. Cell* **1**, 771–781.
- Yanagisawa, S., Yoo, S.D., and Sheen, J.** (2003). Differential regulation of EIN3 stability by glucose and ethylene signalling in plants. *Nature* **425**, 521–525.
- Zilberman, D., Cao, X., and Jacobsen, S.E.** (2003). ARGONAUTE4 control of locus-specific siRNA accumulation and DNA and histone methylation. *Science* **299**, 716–719.

NOTE ADDED IN PROOF

While this manuscript was under review, a report by Olmeda et al. (2006) identified EIN5 as XRN4.

Olmedo, G., Guo, H., Gregory, B.D., Nourizadeh, S.D., Aguilar-Henonin, L., Li, H., An, F., Guzman, P., and Ecker, J.R. (2006). ETHYLENE INSENSITIVE5 encodes a 5' → 3' exoribonuclease required for regulation of the EIN3-targeting F-box proteins EBF1/2. *Proc. Natl. Acad. Sci. USA* **103**, 13286–13293.

The Exoribonuclease XRN4 Is a Component of the Ethylene Response Pathway in *Arabidopsis*
Thomas Potuschak, Amérin Vansiri, Brad M. Binder, Esther Lechner, Richard D. Vierstra and Pascal Genschik

Plant Cell 2006;18;3047-3057; originally published online November 3, 2006;
DOI 10.1105/tpc.106.046508

This information is current as of January 19, 2018

Supplemental Data	/content/suppl/2006/11/03/tpc.106.046508.DC1.html
References	This article cites 45 articles, 22 of which can be accessed free at: /content/18/11/3047.full.html#ref-list-1
Permissions	https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&issn=1532298X&WT.mc_id=pd_hw1532298X
eTOCs	Sign up for eTOCs at: http://www.plantcell.org/cgi/alerts/ctmain
CiteTrack Alerts	Sign up for CiteTrack Alerts at: http://www.plantcell.org/cgi/alerts/ctmain
Subscription Information	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: http://www.aspb.org/publications/subscriptions.cfm