

Seed Germination of GA-Insensitive *sleepy1* Mutants Does Not Require RGL2 Protein Disappearance in *Arabidopsis* ^W

Tohru Ariizumi^a and Camille M. Steber^{a,b,1}

^aDepartment of Crop and Soil Science, Washington State University, Pullman, Washington 99164-6420

^bU.S. Department of Agriculture–Agricultural Research Service, Washington State University, Pullman, Washington 99164-6420

We explore the roles of gibberellin (GA) signaling genes *SLEEPY1* (*SLY1*) and *RGA-LIKE2* (*RGL2*) in regulation of seed germination in *Arabidopsis thaliana*, a plant in which the hormone GA is required for seed germination. Seed germination failure in the GA biosynthesis mutant *ga1-3* is rescued by GA and by mutations in the DELLA gene *RGL2*, suggesting that RGL2 represses seed germination. RGL2 protein disappears before wild-type seed germination, consistent with the model that GA stimulates germination by causing the SCF^{SLY1} E3 ubiquitin ligase complex to trigger ubiquitination and destruction of RGL2. Unlike *ga1-3*, the GA-insensitive *sly1* mutants show variable seed dormancy. Seed lots with high seed dormancy after-ripened slowly, with stronger alleles requiring more time. We expected that if RGL2 negatively controls seed germination, *sly1* mutant seeds that germinate well should accumulate lower RGL2 levels than those failing to germinate. Surprisingly, RGL2 accumulated at high levels even in after-ripened *sly1* mutant seeds with 100% germination, suggesting that RGL2 disappearance is not a prerequisite for seed germination in the *sly1* background. Without GA, several GA-induced genes show increased accumulation in *sly1* seeds compared with *ga1-3*. It is possible that the RGL2 repressor of seed germination is inactivated by after-ripening of *sly1* mutant seeds.

INTRODUCTION

The seed is the main mechanism of plant propagation, and the proper regulation of seed dormancy and germination is of critical importance to plant survival. Seeds that fail to germinate under favorable environmental conditions are said to be dormant (reviewed in Bewley and Black, 1994). Dormant seeds acquire the ability to germinate following a period of dry storage called after-ripening. Dormancy helps to ensure the survival of a species by allowing some ungerminated seeds to survive natural catastrophes. Seed dormancy is also important from the agronomic viewpoint because the absence of dormancy in several cereal grains causes preharvest sprouting when cool humid conditions persist close to harvest. There are two classes of seed dormancy: coat-imposed and embryo dormancy. In coat-imposed dormancy, radicle emergence is blocked by the physical barrier of the testa, aleurone, or endosperm layer. Seed coat-imposed dormancy, which is commonly found in *Arabidopsis thaliana*, can be overcome by cutting the seed coat (Debeaujon and Koornneef, 2000). Conversely, seeds with embryo dormancy will not germinate when the seed coat is cut (Bewley and Black, 1994). Germination of dormant seeds can also be stimulated by

a period of cold imbibition called stratification prior to imbibition under conditions that stimulate germination.

Genetic analysis of seed dormancy is complicated by the fact that the degree of dormancy is dependent on both environmental and genetic factors. However, some genes influencing the degree of seed dormancy have been cloned. For example, quantitative trait loci analysis of a cross between highly dormant *Arabidopsis* ecotype Cvi (Cape Verde Islands) and the less dormant ecotype Landsberg *erecta* (*Ler*) has identified seven quantitative trait loci associated with seed dormancy (Alonso-Blanco et al., 2003). Loss of the *COMATOSE* gene encoding a putative ATP binding cassette transporter results in strong seed dormancy and reduced germination potential without affecting seed or plant morphology (Russell et al., 2000; Footitt et al., 2002). Loss of the *DOF AFFECTING GERMINATION1* gene encoding a putative zinc finger transcription factor results in reduced seed dormancy as a maternal effect without affecting seed coat structure (Papi et al., 2000, 2002). Other examples of genes involved in the establishment and maintenance of seed dormancy include *ABSCISIC ACID INSENSITIVE3*, *LEAFY COTYLEDON1* (*LEC1*), *LEC2*, and *FUSCA3* (Ooms et al., 1993; Keith et al., 1994; Nambara et al., 1995; Parcy and Giraudat, 1997).

Seed dormancy is subject to hormonal control. The plant hormone abscisic acid (ABA) is needed to induce seed dormancy during embryo maturation, whereas the hormone gibberellin (GA) is needed to stimulate seed germination. Mutants with decreased ABA biosynthesis or sensitivity show decreased seed dormancy, whereas mutants with decreased ABA catabolism or increased ABA sensitivity show increased seed dormancy (reviewed in Finkelstein and Rock, 2002; Kushiro et al., 2004; Okamoto et al., 2006).

¹ To whom correspondence should be addressed. E-mail csteber@wsu.edu; fax 509-335-2553.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Camille M. Steber (csteber@wsu.edu).

^W Online version contains Web-only data.

www.plantcell.org/cgi/doi/10.1105/tpc.106.048009

Several lines of evidence demonstrate the importance of GA biosynthesis and signaling for stimulating seed germination in *Arabidopsis*. The role of GA in promoting various stages in *Arabidopsis* growth and development is demonstrated by the phenotypes of GA biosynthesis mutants like *ga1-3*, including severe dwarfism, male sterility, and strong coat-imposed seed dormancy (reviewed in Olszewski et al., 2002). GA application rescues all of these defects. GA does not appear to directly control dormancy per se but acts as an antagonist to ABA by stimulating radicle emergence (reviewed in Bewley, 1997). Many GA biosynthetic genes are upregulated in seed tissues during seed germination and in the cold (Yamaguchi et al., 2001; Penfield et al., 2006). GA signaling is also required for efficient seed germination in *Arabidopsis*. Mutations in the *SLEEPY1* (*SLY1*) GA signaling gene lead to increased seed dormancy, and a triple knockout of the GA receptors *GID1A*, *GID1B*, and *GID1C* leads to failure to germinate (Steber et al., 1998; Griffiths et al., 2006).

Our current understanding of the control of *Arabidopsis* seed germination by GA signaling is derived mostly from studies of the GA signaling genes *SLY1* and the DELLA gene *RGA-LIKE2* (*RGL2*) (Steber et al., 1998; Lee et al., 2002; Bassel et al., 2004; Tyler et al., 2004). DELLA proteins are a subfamily of the GRAS (for GAI, RGA, and Scarecrow) family of putative transcription factors that are nuclear-localized and negatively regulate GA signaling (Peng et al., 1999; Pysh et al., 1999). There are five DELLA proteins in *Arabidopsis* with partially overlapping functions in GA signaling. Consistent with the negative role of DELLA proteins, loss-of-function mutations in DELLA genes result in a decreased requirement for GA in plant growth and development. Based on phenotypic analysis, the main negative regulator of stem elongation is the DELLA protein RGA, although GAI and RGL1 also function in stem elongation. DELLA proteins RGA, RGL2, and RGL1 are the main regulators of flowering (reviewed in Thomas et al., 2005). No function has yet been attributed to RGL3. Finally, RGL2 is the main negative regulator of seed germination, whereas RGA, GAI, and RGL1 play lesser roles (Cao et al., 2005).

GA relieves DELLA inhibition of seed germination, stem elongation, and transition to flowering by triggering DELLA protein destruction via the ubiquitin-proteasome pathway. It has been demonstrated that the GA signal in rice (*Oryza sativa*) is perceived by a GA receptor with homology to human hormone-sensitive lipase, *GID1* (for GA-INSENSITIVE DWARF1) (Ueguchi-Tanaka et al., 2005). Recently, three homologs of the rice *GID1* gene (*GID1a*, *GID1b*, and *GID1c*) have been shown to function in *Arabidopsis* GA signaling (Nakajima et al., 2006). The fact that *GID1* shows GA-dependent interaction with the five DELLA proteins suggests that *Arabidopsis* *GID1* controls DELLA function. On binding GA, *GID1* binds to DELLA protein and appears to stimulate ubiquitination of DELLA proteins by promoting binding to the SCF^{SLY1} E3 ubiquitin ligase (Griffiths et al., 2006; Hartweck and Olszewski, 2006). DELLA proteins have been shown to directly interact with *SLY1* and are believed to disappear rapidly from the nucleus once ubiquitinated due to proteolytic degradation by the 26S proteasome pathway. The *SLY1* gene is a positive regulator of GA signaling encoding the F-box subunit that determines the substrate specificity of an SCF (for Skp1,

Cullin, F-box) E3 ubiquitin ligase complex (McGinnis et al., 2003; Dill et al., 2004; Fu et al., 2004). Mutations in *SLY1* were isolated in screens for decreased germination capacity (Steber et al., 1998; Steber and McCourt, 2001). The *sly1* mutants show increased seed dormancy and increased sensitivity to inhibition of seed germination by ABA (Strader et al., 2004). By contrast, DELLA genes are negative regulators of GA signaling, and mutations in the DELLA gene *RGL2* result in increased capacity to germinate (Lee et al., 2002; Tyler et al., 2004). Thus, it appears that SCF^{SLY1} may positively regulate GA signaling by causing the destruction of the DELLA protein RGL2 during seed germination.

Several lines of evidence indicate that RGL2 is a critical negative regulator of seed germination. The strongest evidence is that mutations in *RGL2* restore seed germination in the *ga1-3* background (Lee et al., 2002; Tyler et al., 2004). The pattern of *RGL2* mRNA expression is consistent with a regulatory role in seed germination. In wild-type seeds, the *RGL2* transcript level rises during a stratification treatment of seed imbibition at low temperature (4°C) and rapidly declines after the seeds are transferred to 23°C, a temperature that favors seed germination (Lee et al., 2002). In the *ga1-3* mutant, however, *RGL2* mRNA levels increase during imbibition at low temperature and remain high even after the seeds are transferred to 23°C. If GA is added, the *RGL2* transcript level rapidly declines and germination capacity increases. Thus, *RGL2* transcript levels negatively correlate with seed germination, suggesting that downregulation of *RGL2* transcript is necessary for seed germination in *Arabidopsis* (Lee et al., 2002; Tyler et al., 2004). Subsequent work by Bassel et al. (2004) showed that the *RGL2* mRNA does not completely disappear prior to radicle emergence, indicating that disappearance of *RGL2* mRNA is not absolutely required for seed germination. Interestingly, protein gel blot analysis showed that RGL2 protein disappears within 5 h of addition of GA to incubating *ga1-3* seeds (Tyler et al., 2004). RGL2 protein did not disappear, however, within 5 h of adding GA to imbibing *sly1-10* seeds. Thus, the current model is that disappearance of RGL2 protein is required for seed germination and that rapid disappearance of RGL2 protein requires the SCF^{SLY1} E3 ubiquitin ligase (Tyler et al., 2004). Based on genetic evidence, it is clear that RGL2 negatively regulates seed germination. The question that remains is whether the decreases in *RGL2* transcript and protein levels are the sole mechanisms controlling GA-stimulated seed germination.

To address this question, this study examines whether the pattern of RGL2 protein accumulation in *sly1* mutants is consistent with the current model for control of seed germination by RGL2. This study shows that the dormant seed phenotype of *sly1* mutants is variable between different seed lots. Those seed lots showing high seed dormancy eventually after-ripen. Stronger alleles show a stronger seed germination phenotype and require more time to after-ripen. For example, the *sly1-2* allele can require up to 2 years to fully after-ripen. Based on the current model, we expected that the level of RGL2 protein accumulation should be higher in those *sly1* lines that show lower germination capacity. Surprisingly, we found that RGL2 protein accumulated at high levels in *sly1* mutants regardless of the degree of seed dormancy. RGL2 protein accumulated to the same level in after-ripened and dormant *sly1-2* seeds prior to radicle emergence.

Indeed, after-ripened *sly1-2* seeds that were capable of 91% germination expressed high levels of RGL2 protein. This suggests that SCF^{SLY1} is the main E3 ubiquitin ligase controlling RGL2 protein accumulation and that in the *sly1* mutant background, RGL2 protein disappearance is not a prerequisite for seed germination. The possibility that the RGL2 protein accumulating in after-ripened *sly1* mutant seeds is either inactive or controlled by other factors is considered.

RESULTS

Isolation of a New *SLY1* Allele

Here, we compare two new mutant alleles to the two recessive alleles of *sly1* in the *Ler* background previously described (Figure 1A; Steber et al., 1998; Steber and McCourt, 2001; McGinnis et al., 2003). The *sly1-2* allele is a 2-bp deletion causing a frame shift and the predicted loss of the last 40 out of 151 amino acids. The *sly1-10* allele contains a 23-bp deletion followed by an ~8-kb insertion, causing the predicted loss of the last eight amino acids and the addition of 46 random amino acids (Figure 1A). The new allele *sly1-DS* in the *Ler* background resembles *sly1-10* and contains a transposon insertion resulting in loss of the last 12 amino acids (Sundaresan et al., 1995). The new *sly1-t2* allele in the Wassilewskija (*Ws*) background was identified in the Sussman lines (Krysan et al., 1999). It contains a T-DNA insertion within the F-box motif (at +219 bp) and is predicted to result in loss of the last 78 amino acids. This is the most severe lesion identified in the *SLY1* gene thus far. The *sly1-t2* allele showed

severe defects in plant development, such as dwarfism and reduced fertility (see Supplemental Table 1 online). The effect of each allele on *SLY1* mRNA levels was examined by RT-PCR using *SLY1* gene-specific primers (Figure 1B). When the *SLY1*-specific primers were employed, no significant change in mRNA accumulation was observed in *sly1-10* and *sly1-2*, whereas a marked decrease was seen in *sly1-t2*. These results indicate that *SLY1* transcript accumulation is reduced by the T-DNA insertion in *sly1-t2*, whereas the other two alleles, *sly1-10* and *sly1-2*, do not alter transcript levels compared with the wild type.

The *sly1* Germination Phenotype Is Variable but Dependent on the Allele and the Age of the Seed

Because GA synthesis is required for seed germination, it was expected that GA signaling would also be required for seed germination. Consistent with this, mutants in the GA signaling gene *SLY1* cause increased seed dormancy and increased sensitivity to ABA in seed germination (Strader et al., 2004). Subsequently, anecdotal evidence showed that the *sly1* seed dormancy phenotype was variable. To characterize the *sly1* seed dormancy phenotype, the percentage of germination of many independent *sly1* seed lots of different ages were compared after 7 d of incubation at 22°C (Figure 2). While the germination capacity of independent seed lots varied significantly, it did correlate with allele and with the period of seed dry storage.

In general, *sly1-2* seeds showed stronger dormancy than *sly1-10* seeds. The *sly1-t2* allele showed the highest degree of seed dormancy. However, the germination capacity of *sly1-t2* seeds cannot be directly compared with *sly1-2* and *sly1-10* because it is in a different genetic background. In *sly1-10*, the germination rate of the 0.5-month-old seed lots ranged from 8.0 to 98.4% (Figure 2A). Less variation in germination rate was seen after seeds were subjected to 1 to 5 months (77.1 to 100%) and 12 to 31 months (96.3 to 100%) of room temperature dry storage. The *sly1-2* allele showed more severe seed dormancy (Figure 2B). In *sly1-2*, the germination rate of the 0.5-month-old seed lots ranged from 2.0 to 43.0%. The germination rate increased following longer periods of dry storage for 1 to 5 months (22.0 to 56.1%), 12 to 20 months (64.2 to 75.4%), and 26 to 28 months (73.4 to 91.1%). Unlike *sly1-10*, *sly1-2* seeds never reached 100% germination. The *sly1-t2* seeds showed the most severe dormancy phenotype. No seed germination was observed in 16 independent seed lots stored for 8 months or less, and seeds stored for 11 to 18 months showed between 0 and 9.6% seed germination (Figure 2C; data not shown). It is not known whether *sly1-t2* seeds ever fully after-ripen. These results indicate that the degree of seed dormancy is dependent on the severity of the *sly1* allele. However, other factors, possibly environmental, may cause the variation in dormancy.

The data in Figures 2A and 2B show that both *sly1-2* and *sly1-10* seeds lose their capacity to germinate following long periods of dry storage. In *sly1-10*, seed lots stored for 49 to 54 months showed 32.9 to 72.3% seed germination, whereas *sly1-2* seeds stored for 56 to 66 months showed 9.2 to 25.2% germination. It is possible that this loss of germination capacity is due to seed deterioration and that this loss of viability is more severe in the stronger allele.




Figure 1. Structure of the *SLY1* Gene and the *SLY1* mRNA Expression Detected by RT-PCR in Three Mutant Alleles.

(A) The mutation sites are shown in three alleles. The *sly1-10* mutant has an 8-kb insertion at +429 nucleotides from translational start. The *sly1-2* mutant has a 2-bp deletion at the +333-nucleotide position. The T-DNA is inserted at the +219-nucleotide position in *sly1-t2*.

(B) The top panel shows the *SLY1* expression in three mutant alleles and the wild type using the *sly1-2f* and *sly1-2r* primers (pair A) that flank the insertion in *sly1-t2* giving no product. The middle panel shows the expression using the 2-63f and *sly1-r* primers (pair B) that flank the *sly1-10* rearrangement giving no product. The bottom panel shows the *ACTIN2* control.




Figure 2. Germination Rates of *sly1* Mutant Seeds Varied but Were Dependent on Allele and Age of Seeds.

Seeds plated on MS-agar were incubated 3 d at 4°C followed by 7 d at 22°C. Percentage of germination was calculated based on cotyledon emergence for independent seed lots of: *sly1-10* (A), *sly1-2* (B), and *sly1-t2* (C). Duration of dry storage or age of seed is indicated above the bars (months). N.G., none of the seeds germinated. Error bars show SE ($n = 3$). Seed lot numbers (#) are indicated on the x axis.

To clarify if dormancy or poor viability caused seeds to show low percentage of germination, intact seeds were stained with tetrazolium solution, a vital stain (Debeaujon et al., 2000). In *sly1-2*, a similar staining pattern was seen in a 0.5-month-old seed lot (#35) showing 2% germination as in a 26-month-old seed lot (#21) showing 91.1% seed germination (Figures 2A, 2B, and 3A). A similar staining pattern was also seen in *Ler* (wild-type), *ga1-3*, and *sly1-10* seeds of comparable age (data not shown). By contrast, reduced staining (66.8% stained) was seen in *sly1-2* seeds stored for 63 months (#7) and giving 10% germination (Figures 2B and 3A). By contrast, 63-month-old *Ler* seeds showed 100% germination and 84.3% tetrazolium-stained seeds. These data suggest that the low percentage of germination in 0.5-month-old *sly1-2* seeds is due to seed dormancy, whereas the reduction in germination capacity of older *sly1-2* seeds is due, at least in part, to loss of longevity. Consistent with this, germination of 0.5-month-old *sly1-2* seeds is rescued by cutting the seed coat as is the germination of 0.5-month-old *ga1-3* seeds and *sly1-t2* (Figure 3B). Failure to achieve 100% germination is probably due to inadvertent damage to the embryo.

***sly1* Mutants After-Ripen Slowly**

The fact that *sly1* seed lots subjected to longer periods of dry storage show better seed germination suggests that dormant *sly1* seeds are able to after-ripen slowly (Figure 2). *Ler* (wild-type) seeds are fully after-ripened and show 100% germination after 0.5 months of dry storage (Figure 4). During an after-ripening time course, a single *sly1-2* seed lot (#37) showed 18.7% germination




Figure 3. Determination of Seed Viability by Tetrazolium Staining and Germination Rescue by Cutting the Seed Coat.

(A) Seeds of the wild type (0.5, 26, and 63 months old), after-ripened *sly1-2* (26 months old, seed lot #21), and dormant *sly1-2* (0.5 and 63 months old, seed lots #35 and #7, respectively) were stained with 0.5% tetrazolium for 10 h. Fully stained seeds were counted as viable.

(B) To examine seed viability, dormant *sly1-2* (seed lot #35, 0.5 months old) and *sly1-t2* (seed lot #12, 0.5 months old) seed coats were nicked with forceps to stimulate seed germination. *ga1-3* (0.5 months old) seeds were used as a positive control for seed germination rescue.

Age of each seed stock is shown below the graph (months). Germination rates (G) are shown below the figure (%). Error bars show SE ($n = 30$).




Figure 4. *sly1* Mutant Seeds Germinate More Slowly Than the Wild Type, and Germination Is Rescued by a Mutation in *RGL2*.

Seeds were plated on MS-agar and the number of germinated seeds scored daily after transfer to 22°C. The wild type (28 months old), *sly1-10* (27 months old, #3), after-ripened (A) *sly1-2* (26 months old, #21), dormant (D) *sly1-2* (0.5 months old, #35), 1-month-old wild type (Col), *sly1-t2 rgl2-13* (1 month old), and *sly1-t2* (1 month old, crossed to wild-type Col) were used. Error bars show SE ($n = 3$).

(A) Percentage of radicle emergence based on radicle protrusion from seed coat.

(B) Percentage of cotyledon emergence based on appearance of green cotyledons outside the seed coat.

(C) Rescue of *sly1-t2* seed germination by the *rgl2-13* mutation based on percentage of radicle emergence.

after 0.5 months of dry storage, 53.2% germination after 1 month of dry storage, and 64.2% after 8 months of dry storage (Figure 2B; see Supplemental Figure 1 online). The period required to after-ripen also appears to depend on the severity of the allele. The *sly1-10* seeds reached ~97.0% germination after 1 to 5 months after-ripening, whereas *sly1-2* seeds reached 73.4 to 91.1% germination after 26 to 28 months after-ripening (Figure 2).

Segregation analysis was used to examine the dormant seed phenotype of *sly1-t2*. The mutant was crossed with Ws (wild-type), and F₂ progeny were sown to check the segregation ratio of germinated-to-ungerminated seeds. Of 133 F₂ progeny, 29

failed to germinate, giving a 3:1 segregation ratio ($\chi^2 = 0.72$; $1.0 < P < 0.5$). These results suggested that the seed dormancy phenotype in *sly1-t2* was caused by a single recessive mutation. When germination was rescued by cutting the seed coat, all of the ungerminated seeds gave rise to dwarf *sly1* plants (data not shown). None of the 104 germinated seedlings gave rise to dwarves. The *sly1-t2* allele shows 0 to 9.9% seed germination following 18 months of after-ripening. Thus, this severe truncation of the *SLY1* gene results in strong seed dormancy.

sly1 Seeds Germinate Slowly and Show Increased Sensitivity to Polyethylene Glycol and Paclobutrazol in Germination

After-ripened *sly1* mutant seed lots showed reduced germination capacity compared with wild-type *Ler* seeds. This was first evaluated based on percentage of germination over time. To accurately examine the effect of the *sly1* mutations on germination rate, both the percentage of radicle emergence (Figure 4A) and percentage of cotyledon emergence (Figure 4B) were determined over time comparing wild-type *Ler* with *sly1-10* (#3) and dormant and after-ripened *sly1-2* seeds (#35 and #21, respectively). Similar curves were obtained using percentage of radicle emergence and cotyledon emergence, except that cotyledon emergence occurred ~1 d (24 h) after radicle emergence. Wild-type *Ler* and Ws (data not shown) reached 100% radicle emergence within 2 d. Germination was considerably slower in *sly1* mutants. The *sly1-10* seeds required 5 d to reach 100% radicle emergence, whereas after-ripened *sly1-2* seeds (#21) required 7 d to reach a final 85.6% radicle emergence.

Plating experiments using increasing concentrations of polyethylene glycol (PEG) 8000 were used to examine the effect of increasing media osmotic potential on *sly1* mutant and wild-type seed germination. We have observed that when *sly1-10* and *sly1-2* seeds are sown directly on moistened soil, virtually no seeds germinated (data not shown). This observation suggested that the *sly1* mutant seeds have difficulty imbibing water. To examine this, after-ripened wild-type and *sly1* mutant seeds were sown on Murashige and Skoog (MS)-agar media containing various low concentrations of PEG 8000. This experiment showed that *sly1* seeds were more sensitive to PEG 8000 than the wild type and that *sly1-2* showed a more severe germination phenotype than *sly1-10* (see Supplemental Figure 2 online). Next, the sensitivity of *sly1* seeds to GA biosynthesis inhibitor paclobutrazol (PAC) was examined. After-ripened wild-type, *sly1*, and *ga1-3* mutant seeds were first incubated in 100 μ M PAC solution for 3 d at 4°C, and then these seeds were sown on different concentrations of GA₃. This experiment showed that *sly1* seeds were more sensitive to PAC than the wild type and that *sly1-2* showed a more severe germination phenotype than *sly1-10* (see Supplemental Figure 3 online).

A *sly1-t2 rgl2-13* double mutant was constructed to determine the effect of the *rgl2-13* mutation on the *sly1* seed germination phenotype. Although the germination of double mutant seeds was somewhat slower than the wild type, the 1-month-old *sly1-t2 rgl2-13* seeds reached 100% germination by day 4, whereas the 1-month-old *sly1-t2* seeds crossed into the Columbia (Col) background never germinated (Figure 4C). Thus, it appears

that the *RGL2* mutation is sufficient to rescue *sly1-t2* germination. Similar suppression of the *sly1-2* germination phenotype was seen in a *sly1-2 rgl2-13* double mutant (data not shown). This result suggests that the reduced germination in *sly1* mutants is due to overaccumulation of RGL2 protein. This is consistent with data of Tyler et al. (2004) showing that RGL2 protein disappears within 5 h of GA treatment in *ga1-3* seeds but not in *sly1* mutant seeds.

RGL2 Does Not Disappear during *sly1* Mutant Seed Germination

This section examines the hypothesis that the decreased germination capacity of GA mutants is due to increased RGL2 protein accumulation. The expression of RGL2 protein was examined over a time course of seed imbibition in wild-type, *ga1-3*, and *sly1* mutant seeds to determine whether RGL2 disappears prior to wild-type seed germination and to determine whether RGL2 protein levels correlate with the severity of the *sly1* seed germination phenotype (Figure 5). This experiment was conducted using polyclonal antibody to RGL2 that was found to be specific




Figure 5. Accumulation of RGL2 Protein during a Time Course of Seed Imbibition.

RGL2 protein was detected by protein gel blot analysis.

(A) *Ler* seeds were incubated for 6 d at 4°C and then time points were at indicated intervals after transfer to 22°C (blot exposure >30 min). The correlation between RGL2 protein accumulation and radicle emergence (%) was determined. RGL2 protein disappeared within 12 h, before radicle emergence.

(B) Time-course analysis of RGL2 protein in wild-type and mutant backgrounds. Protein gel blot analysis (exposure time 5 to 10 min) of protein isolated from seeds imbibed at 4°C for 1, 3, 5, and 6 d and from seeds first imbibed for 3 d at 4°C then transferred to 22°C for indicated number of days. Protein from dry seeds was also examined. Seed samples used were *Ler* (28 months old), *ga1-3* (25 months old), *sly1-10* (27 months old, #3), after-ripened (A) *sly1-2* (26 months old, #21), dormant (D) *sly1-2* (0.5 months old, #35), *Ws* (0.5 months old), and *sly1-t2* (0.5 months old, #12). Asterisks indicate nonspecific bands.

using double mutant analysis (Hussain et al., 2005; see Supplemental Figure 4 online)

It was expected that if RGL2 is a major negative regulator of seed germination, RGL2 protein should disappear well before wild-type radicle emergence. Following 6 d of imbibition at 4°C, *Ler* (wild-type) seeds were transferred to 22°C, and time points were taken for protein blot analysis using RGL2 antibody (Figure 5A; see Supplemental Figure 4C online). RGL2 protein was present at low levels immediately after transfer to 22°C, persisted at very low levels through 5 h of incubation, and disappeared by 12 h of incubation. At 12 h of incubation, only 2% of seeds showed radicle emergence. Thus, RGL2 protein disappeared well before completion of radicle emergence at 24 h incubation at 22°C. This is consistent with data from previous studies suggesting that RGL2 protein must disappear prior to wild-type seed germination (Tyler et al., 2004).

If the seed germination phenotype of GA mutants is due to overaccumulation of RGL2 protein, then we expect the severity of the germination phenotype to positively correlate with the level of RGL2 protein accumulation. To test this hypothesis, a time course was performed examining RGL2 protein accumulation during imbibition of wild-type, *ga1-3*, and *sly1* mutant seeds (1) over 6 d of stratification at 4°C and (2) over 2 to 6 d of imbibition at 22°C following 3 d of stratification at 4°C (Figure 5B). In both *Ler* and *Ws* seeds, RGL2 protein accumulation was barely detected within 5 to 6 d of seed imbibition at 4°C (Figure 5; see Supplemental Figure 4C online). When wild-type seeds were transferred to 22°C following 3 d of stratification, RGL2 protein was not detected after 24 h of transfer. In the *ga1-3* seeds, RGL2 protein was detected earlier and at higher levels than in *Ler* during stratification at 4°C and did not decrease during 6 d of incubation at 22°C. During this time, *ga1-3* seeds did not germinate.

RGL2 protein accumulation in after-ripened *sly1-10* (seed lot #3, 27 months old) and *sly1-2* (seed lot #21, 26 months old) seeds known to show efficient seed germination were compared with dormant *sly1-2* (seed lot #35, 0.5 months old) and *sly1-t2* seeds (seed lot #12, 0.5 months old) (Figures 2 and 5B). The percentage of radicle emergence for each time point in this experiment can be found in Figure 4A. When *sly1* mutant seeds were stratified at 4°C, the RGL2 protein gradually accumulated to high levels similar to the pattern of expression seen in *ga1-3* (Figure 5B). Following transfer of seeds to 22°C, RGL2 protein accumulated at higher levels in *sly1* than in *ga1-3* mutants, regardless of allele, degree of seed dormancy, or seed lot (Figure 5B; see Supplemental Figures 4B and 4D online). Interestingly, although the *sly1-10* seeds showed 73.7% radicle emergence after 3 d of incubation at 22°C, RGL2 protein levels remained high. Rather than decreasing as germination approached, RGL2 protein levels gradually increased, reaching a maximum at day 3, and remained high after 100% radicle emergence was reached at day 5 (Figures 4 and 5B). It was expected that since *sly1-2* shows a more severe germination phenotype than *sly1-10*, the RGL2 protein levels would be higher than in *sly1-10*. However, the accumulation of RGL2 protein in the after-ripened *sly1-2* seeds was similar to or less than that seen in *sly1-10*. It was expected that RGL2 protein would accumulate at higher levels in dormant *sly1-2* seeds than after-ripened. However, almost no difference was seen between after-ripened and dormant *sly1-2* through 4 d

of incubation at 22°C. At 5 and 6 d of incubation at 22°C, slightly higher RGL2 levels were seen in dormant *sly1-2* than in after-ripened *sly1-2*. At day 6, dormant *sly1-2* seeds showed 7.4% radicle emergence, whereas after-ripened *sly1-2* seeds showed 85.6%. Thus, higher germination levels may partly correlate with reduced RGL2 protein levels in after-ripened *sly1-2* seeds. However, since after-ripened *sly1-2* reached near-maximum radicle emergence (71.2%; Figure 4) after 4 d at 22°C, it does not appear that decreased RGL2 protein levels are required prior to radicle emergence in the *sly1-2* or *sly1-10* genotype. The *sly1-t2* allele had the strongest seed germination phenotype, showing 0% germination at the conclusion of the experiment. Compared with wild-type *Ws*, the *sly1-t2* allele also showed increased RGL2 protein accumulation. After the seeds were transferred at 22°C, the RGL2 protein persisted throughout the time course.

Next we examined the expression of another DELLA protein, RGA, during seed imbibition. In a double mutant study examining four of the five DELLA genes, mutations in *RGA* rescued the germination of *ga1-3* seeds better than any DELLA mutation but *rgl2* (Cao et al., 2005). When we examined RGA protein accumulation during seed germination, RGA protein showed a very similar expression pattern to RGL2 (see Supplemental Figure 5C online). The RGA protein accumulated at high levels after 6 d of stratification and did not disappear after transfer to 22°C in *ga1-3* and *sly1* mutant seeds. Somewhat higher protein accumulation was found in after-ripened *sly1-2* than in dormant *sly1-2* after 4 d at 22°C (see Supplemental Figure 5D online). These results indicate that at least two major DELLA proteins involved in seed germination do not disappear during *sly1* seed germination.

***sly1* Mutations Cause Increased RGL2 mRNA Accumulation**

The increased level of RGL2 protein in *sly1* mutants could be either due to a requirement for *SLY1* in triggering the disappearance of RGL2 protein or mRNA. It was expected that *sly1* mutants act by stabilizing the RGL2 protein since *SLY1* is the F-box subunit of an SCF E3 ubiquitin ligase required for DELLA protein disappearance (McGinnis et al., 2003; Dill et al., 2004; Fu et al., 2004). To determine if this is true, the effect of *sly1* mutants on the accumulation of *RGL2* mRNA was determined by RT-PCR analysis of the same seed lots and imbibition time points used to examine RGL2 protein expression in Figure 5B (Figure 6). Control RT-PCR reactions using the *UBQ11* primers are shown in Supplemental Figure 6A online. As previously shown, the *RGL2* mRNA levels gradually increased during imbibition of wild-type and *ga1-3* mutant seeds over 6 d at 4°C (Lee et al., 2002). A similar pattern of expression was seen during cold imbibition of *sly1-10*, *sly1-t2*, and dormant and after-ripened *sly1-2* seeds. Next, *RGL2* mRNA expression was examined following transfer to 22°C. *RGL2* mRNA levels decreased within 1 d (24 h) of transfer to 22°C in *Ler* and *Ws* seeds, the time point corresponding to 72.5 and 81.6% radicle emergence, respectively. *RGL2* mRNA levels never decreased during imbibition of *ga1-3* seeds that failed to germinate.

In *sly1* mutants, *RGL2* mRNA levels appeared to decrease after radicle emergence. *RGL2* mRNA levels decreased after 4 d of imbibition at 22°C in *sly1-10* and after 5 d of imbibition in after-ripened *sly1-2* seeds. These time points corresponded to 90.3




Figure 6. *RGL2* mRNA Accumulation during a Time Course of Seed Imbibition for Wild-Type, *ga1-3*, and *sly1* Mutant Seeds.

RT-PCR analysis of *RGL2* mRNA expression in seeds imbibing using the same time courses in Figure 6 of stratified (4°C for 1, 3, 5, and 6 d) seeds and seeds imbibing at 22°C after 3 d of stratification at 4°C. Seed samples used were identical to Figure 6 and include *Ler* (28 months old), *ga1-3* (25 months old), *sly1-10* (27 months old, #3), after-ripened (A) *sly1-2* (26 months old, #21), dormant (D) *sly1-2* (0.5 months old, #35), *Ws* (0.5 months old), and *sly1-t2* (0.5 months old, #12).

and 81.6% radicle emergence, respectively. Like *ga1-3*, the *RGL2* mRNA levels remained high through 6 d of imbibition of dormant *sly1-2* and *sly1-t2* seeds showing 9.4 and 0% final radicle emergence, respectively (Figure 4). The decline in *RGL2* mRNA levels during this time course occurred prior to the decrease in RGL2 protein levels (Figure 5B). In *sly1-10*, *RGL2* mRNA levels decreased after 4 d of imbibition at 22°C, while RGL2 protein levels remained high at least through 5 d of imbibition. In after-ripened *sly1-2* seeds, *RGL2* mRNA levels decreased after 5 d of imbibition at 22°C, whereas RGL2 protein levels remained high through 6 d of imbibition. This suggests that the high levels of RGL2 protein accumulation following seed germination of *sly1-10* and *sly1-2* seeds are due at least in part to loss of SCF^{SLY1}-stimulated proteolysis of RGL2 protein.

GA Partially Rescues *sly1* Germination without Causing Reduced RGL2 Protein Levels

The current model is that GA stimulates germination via degradation of DELLA proteins, including RGL2, RGA, and GAI (Dill et al., 2004; Fu et al., 2004; Tyler et al., 2004). The strongest evidence of this is that RGL2 protein disappears within 5 h of adding GA to imbibing *ga1-3* seeds but not *sly1* mutant seeds. This result raised the question of whether GA treatment can stimulate *sly1* mutant seed germination and the disappearance of RGL2 protein over a time course of imbibition. Previous work showed that GA treatment did not cause an increase in *sly1-2* hypocotyl elongation (Steber and McCourt, 2001). As shown in Figure 7A, GA treatment increased the percentage of germination of all *sly1* alleles tested,




Figure 7. GA Partially Rescued *sly1* Germination without Reducing RGL2 Protein Levels.

(A) Percentage of germination on indicated GA₃ concentration in *ga1-3* biosynthesis mutant (25 months old), *sly1-10* (5 months old, #13), after-ripened (A) *sly1-2* (26 months old, #21), dormant (D) *sly1-2* (0.5 months old, #35), and *sly1-t2* (0.5 months old, #12; 18 months old, #2). Germination was calculated based on cotyledon appearance. Age of the stock is shown below the line (months). Error bars show SE ($n = 30$ to 60).

(B) Percentage of radicle emergence determined during time-course experiment examining RGL2 protein accumulation in the presence of 10 μM GA₃ using the same seed lots used in **(A)** except for 0.5-month-old *sly1-t2*. Radicle emergence was determined at 60, 72, and 84 h at 22°C following 3 d of 4°C.

(C) RGL2 protein expression in the absence and presence of 10 μM GA₃. Protein was extracted from imbibing seeds at indicated time points at 22°C following 3 d at 4°C.

(D) RGL2 mRNA accumulation in the absence and presence of GA₃ during the same time-course experiment in **(C)**. Left panel, RGL2 mRNA; right panel, ACTIN2 mRNA.

even dormant *sly1-2* and *sly1-t2*. Improvement of the germination was dependent on the concentration of GA applied, and the best recovery was found at 10 μM GA₃ (Figure 7A). Next, we examined the RGL2 protein levels after GA treatment of the same seed lots used for the germination experiments. Seeds were plated either in the absence or presence of 10 μM GA₃ and protein extracts prepared after the indicated incubation at 22°C after 3 d of stratification at 4°C. As expected, RGL2 protein disappeared after 60 h of incubation on GA in *ga1-3* but not in the *sly1* mutants (Figures 7B and 7C). Further time points showed that RGL2 protein levels did not decrease in *sly1* mutants after 84 h of incubation on GA. When RGL2 mRNA levels were examined by RT-PCR analysis, the apparent level of RGL2 mRNA remained constant in dormant *sly1-2* and *sly1-t2* and decreased slightly after 84 h of incubation in the presence of GA in after-ripened *sly1-2* and *sly1-10* (Figure 7D).

Overall, these results suggest that GA treatment improves seed germination without reducing RGL2 protein levels in *sly1-10*, *sly1-2*, and *sly1-t2* backgrounds and suggests the existence of an RGL2-independent pathway for stimulating seed germination.

GA-Regulated Genes Show Altered Expression in *sly1* Seeds

The fact that *sly1* mutant seeds can germinate in the presence of high levels of RGL2 protein suggested that RGL2 protein disappearance may not be required to derepress GA-dependent gene expression in the *sly1* mutant background. To investigate this notion, the mRNA levels of GA-regulated genes were examined during the imbibition of *sly1* seeds and *ga1-3* after 60 h of incubation at 4°C in water followed by 24 h of imbibition at 22°C on MS-agar and then 12 h of imbibition at 22°C in the presence and

absence of GA₃. At this time, none of the seeds had germinated. We examined seven transcripts showing GA induction in *ga1-3* based on previous reports (Herzog et al., 1995; Ogawa et al., 2003; Yamauchi et al., 2004; Cao et al., 2006). The mRNA accumulation in these seven genes was upregulated after GA was applied to *ga1-3* seeds (Figure 8). Dormant *sly1-2* seeds show reduced accumulation of GA-induced genes in the presence and sometimes in the absence of GA compared with *ga1-3*. This inability to induce GA-regulated genes is consistent with a GA-insensitive phenotype. Interestingly, the expression of many GA-induced genes is increased in after-ripened *sly1-2* seeds compared with dormant seeds both in the presence and absence of exogenous GA. After-ripened seeds of the weaker *sly1-10* allele did not appear to have as strong a GA-insensitive phenotype as *sly1-2* since some GA-regulated transcripts showed GA-induced expression. In after-ripened seeds, four transcripts, α -EXPANSIN1 (*EXP1*), XYLOGLUCAN ENDOTRANGLYCOSYLASE5 (*XTH5*), *XTH31*, and CYSTEINE PROTEINASE (*CP*), showed increased expression in the absence of GA in after-ripened *sly1-2* and *sly1-10* compared with *ga1-3*. Three transcripts, PECTIN METHYLESTERASE (*PME*), GIBBERELLIC ACID STIMULATED ARABIDOPSIS4 (*GASA4*), and SERINE CARBOXYPEPTIDASE S10 FAMILY PROTEIN (*SCS10*), showed higher expression in after-ripened *sly1-2* than in *ga1-3* in the




Figure 8. Effect of *sly1* Mutations on Accumulation of GA-Regulated Transcripts in Imbibing Seeds.

mRNA accumulation of seven GA-induced genes was determined by RT-PCR. mRNA was isolated from ungerminated imbibing seeds.

absence of GA and showed lower accumulation in the presence of GA. These patterns of expression are consistent with the notion that the RGL2 protein accumulating in after-ripened *sly1* seeds is not fully active as a repressor of GA-regulated gene expression.

DISCUSSION

Mutations in *SLY1* Result in Allele-Dependent Defects in Seed Germination

The data presented here provide clear evidence that *sly1* mutations cause increased seed dormancy, delayed after-ripening, and decreased germination potential (Figures 2 to 5). Even *sly1* mutant seed lots showing 100% seed germination germinate more slowly and have increased sensitivity to inhibition of germination by ABA, PEG 8000, and PAC (Strader et al., 2004; Figure 4; see Supplemental Figures 2 and 3 online). In addition, alleles that caused loss of more of the *SLY1* protein sequence also caused a more severe seed germination defect (Figures 2 to 4; see Supplemental Figures 2 and 3 online). For example, the *sly1-2* mutation resulting in loss of the C-terminal 40 amino acids showed stronger seed dormancy and required more time to after-ripen than the *sly1-10* mutation, resulting in loss of the last eight amino acids of the protein (Figures 1 and 2). These results also are consistent with the fact that *sly1-2* shows more severe dwarfism and reduced fertility compared with *sly1-10* (McGinnis et al., 2003; Dill et al., 2004; see Supplemental Table 1 online). The new *sly1-t2* allele exhibited the strongest seed dormancy of the three alleles, suggesting that *sly1-t2* may be a null allele. However, a direct comparison is not possible because different ecotypes show differences in seed dormancy (Debeaujon and Koornneef, 2000). Tetrazolium staining and after-ripening experiments showed that dormant *sly1-2* seeds are viable. However, *sly1-2* seeds stored for ~5 years show reduced viability, whereas *Ler* wild-type seeds of the same age show no decrease in germination or viability (Figure 3). The *sly1-2* and *sly1-10* seeds showed variability in the degree of seed coat-imposed dormancy (Figure 2). This variability is most likely due to environmental effects, possibly through differences in accumulation and sensitivity to other plant hormones, such as brassinosteroid, ethylene, and ABA (Koornneef and Karssen, 1994; Steber and McCourt, 2001; Okamoto et al., 2006). Both the *sly1-2* and *sly1-10* mutations result in increased sensitivity to inhibition of seed germination by ABA (Strader et al., 2004). Thus, it is possible that the variability in the *sly1* germination phenotype is due to differences in ABA levels or signaling. It is interesting to note that transcript levels of GA-induced genes were lower in imbibing dormant *sly1-2* seeds than in after-ripened *sly1-2* seeds (Figure 8). It is possible that dormancy somehow represses GA-induced genes in *sly1-2* seeds. This suggests that *SLY1* contributes to, but is not essential for, the derepression of these genes during after-ripening.

The fact that *sly1* mutants show increased sensitivity to PEG 8000 inhibition of seed germination compared with the wild type does suggest that *sly1* mutations cause altered permeability to water. While light and transmission electron microscopy

revealed no obvious morphological differences in the seed coat structure of wild-type and *sly1* seeds (K.M. McGinnis and C.M. Steber, unpublished data), we cannot exclude the possibility that changes in the seed coat are involved. Many of the GA-induced genes that show decreased expression in dormant *sly1-2* seeds, such as *EXP1*, *XTH5*, *XTH31*, and *PME*, are involved in cell wall loosening and may help to release the testa or endosperm barriers during seed germination (Ogawa et al., 2003; Yamauchi et al., 2004). Thus, it is possible that the *sly1* mutations cause increased seed dormancy in part through effects on seed coat loosening.

Mechanisms Underlying the *sly1* Mutant Seed Germination Phenotype

Analysis of *RGL2* mRNA accumulation in *sly1* mutants showed that the *SLY1* gene is required for GA-regulated disappearance of *RGL2* mRNA. Previous studies demonstrated that (1) *RGL2* mRNA levels increase during cold imbibition of wild-type and *ga1-3* seeds, (2) *RGL2* mRNA levels decrease in wild-type seeds imbibed at 23°C during radicle emergence, (3) *RGL2* mRNA levels persist at high levels in *ga1-3* seeds imbibing at 23°C, and (4) GA treatment of *ga1-3* seeds stimulates germination and triggers reduced accumulation of *RGL2* mRNA (Lee et al., 2002; Bassel et al., 2004). Bassel et al. (2004) showed that *RGL2* mRNA did not fully disappear prior to radicle emergence, indicating that *RGL2* mRNA disappearance is not required for seed germination. This work further suggested that *RGL2* activity might be controlled by multiple mechanisms. Here, we demonstrated that *RGL2* mRNA overaccumulated in both dormant and after-ripened *sly1* seeds imbibed at 22°C in a manner similar to that seen in *ga1-3* seeds (Figure 6). Unlike *ga1-3* seeds, however, *sly1* seeds showed no significant decrease in *RGL2* mRNA levels after 60 h of GA treatment (Figure 7), suggesting that *SLY1* is required for GA-stimulated disappearance of *RGL2* mRNA. *RGL2* mRNA accumulation did decrease after germination of *sly1-10* and after-ripened *sly1-2* seeds (Figures 6 and 7). This suggests that *RGL2* mRNA accumulation is subject to *SLY1*-independent downregulation after seed germination. *RGL2* mRNA decreased markedly when wild-type *Ler* reached 72.5% radicle emergence (day 1), whereas *RGL2* transcript levels remained quite high when *sly1-10* (73.7%) and *sly1-2* (71.2%) showed comparable radicle emergence at 3 and 4 d, respectively (Figures 4 and 6). Thus, *RGL2* mRNA appears to be downregulated during seedling establishment in *sly1* mutants.

Previous studies postulated that GA stimulates seed germination by causing the SCF^{SLY1} complex to trigger the proteolysis of the RGL2 protein via ubiquitination (Tyler et al., 2004). This model predicted that the degree of seed dormancy should directly correlate with the level of RGL2 protein in the seeds. The *sly1* mutants that showed variable seed dormancy and that required a long period of after-ripening provided an excellent tool for examining this prediction. This study showed that *sly1* mutants accumulate high levels of RGL2 and RGA protein during imbibition regardless of the *sly1* allele or the dormancy status of the seed (Figure 5B; see Supplemental Figure 5 online). What this result does show is that the degree of *sly1* dormancy is due to factors other than the absolute level of RGL2 protein accumu-

lation. Since mutations in *RGL2* rescue the increased dormancy and reduced germination capacity of dormant *sly1* mutant seeds, it appears that these phenotypes are due to RGL2 overaccumulation. The increased germination of *sly1* seeds due to after-ripening may be due to inactivation of RGL2 or to competition with parallel pathways. Based on previous work, we considered the possibility that other E3 ubiquitin ligases, such as the *SLY1* homolog *SNEEZY* and the *PHOR1* U-box E3 ubiquitin ligase, could replace *SLY1* in GA signaling and lead to *SLY1*-independent fluctuations in RGL2 levels during seed germination (Monte et al., 2003; Fu et al., 2004; Strader et al., 2004). No such fluctuations were seen, suggesting that SCF^{SLY1} is the major E3 ubiquitin ligase transmitting the GA signal under the imbibition conditions used. RGL2 protein does not disappear in GA-treated *sly1* mutant seeds, but GA treatment does somewhat stimulate seed germination (Figure 7). This suggests that GA may stimulate *sly1* mutant seed germination via an RGL2-independent pathway, either through a gene that stimulates seed germination or a gene that controls RGL2 activity (Figure 9A). This notion that there is an RGL2-independent mechanism for GA regulation of




Figure 9. Model for Control of Seed Germination by SLY1 and RGL2.

(A) There is an alternative pathway controlling seed germination that is stimulated by GA. This pathway may bypass the *sly1* mutation either by inhibiting RGL2 activity without causing decreased RGL2 protein levels possibly by posttranslational modification or by activating seed germination independently of SLY1/RGL2.

(B) In the *ga1-3* mutant (no GA), RGL2 does not interact with the GA receptor GID1. In this case, RGL2 functions purely as a negative regulator. Addition of GA rescues *ga1-3* seed germination by causing RGL2 to be destroyed following interaction with GA-GID1 and SCF^{SLY1}. In the *sly1* mutant (GA present), RGL2 cannot be destroyed because SLY1 is defective. In this case, RGL2 acts as a repressor of GA responses when not bound to GID1, and GA-GID1-RGL2 may be inactive as a repressor or act as a positive regulator of seed germination.

seed germination is consistent with a recent microarray study showing that there are GA-regulated genes with DELLA-independent expression (Cao et al., 2006).

Mechanisms Controlling RGL2 Function

The fact that loss-of-function mutations in DELLA genes rescue the defects of the *ga1-3* GA biosynthesis mutant forms the genetic basis for the model that DELLA proteins act as negative regulators of GA signaling and that GA acts via negative regulation of DELLA proteins (Dill and Sun, 2001; King et al., 2001; Cheng et al., 2004). Previous work has also shown that DELLA proteins disappear in response to GA and that this disappearance requires the SCF^{SLY1} E3 ubiquitin ligase (Silverstone et al., 2001; McGinnis et al., 2003; Dill et al., 2004; Fu et al., 2004; Tyler et al., 2004). Prior to this study, it was known that GA treatment of *ga1-3* seeds causes RGL2 protein to disappear prior to the disappearance of *RGL2* mRNA (Tyler et al., 2004). These data together with the evidence that RGL2 persists at high levels after GA treatment of *sly1* mutants are evidence that the RGL2 DELLA protein is also subject to GA-stimulated destruction via the SCF^{SLY1} and ubiquitin-proteasome pathway (Figure 5).

This article and others have revealed that *RGL2* function in seed germination must be regulated at multiple levels, including transcript and protein accumulation (Lee et al., 2002; Bassel et al., 2004; Tyler et al., 2004; Cao et al., 2005; Hussain et al., 2005; Figures 5 to 7). This study suggests that RGL2 functions as a repressor of seed germination in dormant *sly1* seeds and that the RGL2 repressor is deactivated by after-ripening either due to effects of dry storage itself or due to posttranslational regulation. Transcript levels of GA-induced genes are higher in imbibing after-ripened *sly1* seeds than in *ga1-3* seeds in the absence of GA (Figure 8). These data suggest that the RGL2 protein accumulating in after-ripened *sly1* mutant seeds is either less active as a repressor of GA-induced genes or is competing with a transcriptional activator of GA-induced genes (Figure 9A). One possibility is that RGL2 activity is controlled by posttranslational regulation. It appears that RGL2 is phosphorylated and that phosphorylation stabilizes the protein (Hussain et al., 2005). Possibly a kinase either activates or inactivates the RGL2 repressor (Figure 9A). It is also known that the SPINDLY O-Glc-NAC transferase acts as a negative regulator of GA signaling during seed germination (Jacobsen and Olszewski, 1993). It is possible that O-Glc-NAC modification is needed for RGL2 to act as a repressor and that the RGL2 protein accumulating in *sly1* mutants is incompletely modified. Accumulation of unmodified RGL2 in *sly1* mutants may be due to overaccumulation of the protein or to indirect effects of the *sly1* mutation. Alternatively, RGL2 activity may be regulated by protein-protein interaction. For example, RGL2 alone may act as a negative regulator of GA signaling, while RGL2 bound to the GA receptor GID1 may act as a positive regulator of GA signaling (Figure 9B). Positive regulation of GA signaling by GA-GID1-RGL2 in *sly1* mutants may compete with negative regulation by RGL2, leading to an intermediate germination phenotype in spite of the fact that RGL2 is present at high levels. This may also account for the fact that the *sly1* phenotypes in plant height and in fertility are less severe than those of *ga1-3* (Steber et al., 1998).

METHODS

Plant Materials and Growth Conditions

Seeds of wild-type *Arabidopsis thaliana* (accessions Ler and/or Ws), *sly1-10*, *sly1-2*, *sly1-DS*, and *sly1-t2* were cultivated according to McGinnis et al. (2003). The *sly1-t2* line was isolated by searching the Sussman Basta lines by PCR using the T-DNA left border primer JL202 (5'-CAT-TTTATAATAACGCTGCGGACATCTAC-3') and SLY1-specific primer sly1T-DNAr (5'-ACTTTCCGATTACGATTGTCCAGGTGA-3') to obtain a 2.0-kb product (L. Strader and C.M. Steber, unpublished data; Krysan et al., 1999). Homozygous mutant plants were identified by PCR using the primers above and the SLY1-specific primers 2-62f (5'-AAGGCATCT-GAGAAACCC-3') (2.1-kb product) and sly1T-DNAr that do not amplify the large T-DNA insertion in *sly1-t2*. The *sly1-DS* line was obtained from ABRC (stock No. CS101918; Sundaresan et al., 1995). The *sly1-t2 rgl2-13* double mutant was made by crossing the *sly1-t2* allele in the Ws ecotype to *rgl2-13* in the Col ecotype. For comparison in germination experiments, we also crossed *sly1-t2* once to Col wild type. Failure to germinate was seen in all independent isolates of *sly1-t2* recovered from the cross to Col (data not shown).

Germination Experiments

For germination experiments, 30 to 60 seeds were sterilized with 10% bleach for 15 to 20 min and plated per repetition on 0.5 × MS salts (Sigma-Aldrich)/0.8% agar (MS-agar). Percentage of germination was determined following 3 d of incubation at 4°C followed by 7 d of incubation under constant fluorescent light at 22°C. The average germination rate was calculated using three independent experiments. After harvest, seeds were stored in open tubes for 2 weeks at room temperature and then stored at room temperature in closed tubes in a desiccation cabinet. For viability tests, seeds were soaked in 0.5% 2,3,5-triphenyltetrazolium chloride (Sigma-Aldrich) solution at 37°C in darkness for 10 h. Fully red-stained seeds were counted as viable (Debeaujon et al., 2000). For preparation of the PEG 8000-supplemented media, a sterile 40% stock solution of PEG 8000 was prepared and then mixed into the MS-agar media to give 4, 6, and 8% PEG 8000 plates. Seeds from homozygous *sly1-10*, after-ripened *sly1-2*, and wild-type Ler were plated on this PEG-containing media. After 3 d of incubation at 4°C, they were transferred to 22°C under constant light, and the germination rate was determined over 7 d. For experiments in which GA was applied, seeds from homozygous *ga1-3*, *sly1-10*, after-ripened *sly1-2*, dormant *sly1-2*, *sly1-t2*, and the wild type (Ler and Ws) of similar age were used. Seeds were sown on MS-agar plates containing 0, 0.1, 1.0, and 10 μM of GA₃. After 3 d of incubation at 4°C, they were transferred to 22°C under constant light, and germination rate was determined at 60, 72, and 84 h. For PAC experiments, homozygous *ga1-3*, *sly1-10*, after-ripened *sly1-2*, and the wild type (Ler) were used. First, these seeds were incubated in 100 μM PAC solution at 4°C for 3 d, washed five times with distilled water, and sown on MS-agar plates containing 0, 0.1, 1.0, and 10 μM of GA₃. The MS-agar plates were then incubated at 22°C under constant light, and germination rate was determined after 7 d.

RT-PCR Analysis of mRNA Expression

RT-PCR analysis was used to analyze the accumulation of SLY1 mRNA expression in different mutant alleles. Total RNA was isolated from flower buds using an RNA easy kit (Qiagen), and genomic DNA contamination was removed using the DNA-Free RNA kit (ZYMO Research). cDNA was generated from 5 μg total RNA using a first-strand cDNA synthesis kit (GE Healthcare). cDNA was diluted and then used as a template for PCR amplification with the SLY1-2f (5'-CTCCGCTGATGGTTTGCTT-3') and

Sly1-2r (5'-CCTCTCAGAGAATTTCTCACGC-3') primers to obtain a 147-bp product and with the 2-63f and sly1-r primers to obtain a 213-bp product. Sly1-2f and Sly1-2r primers detect mRNA in *sly1-10*, *sly1-2*, and the wild type but not in *sly1-t2* because they flank the T-DNA insertion. 2-63f and sly-r primers detect transcript in *sly1-t2*, *sly1-2*, and the wild type but not in *sly1-10* because these primers flank the DNA rearrangement in this allele. *ACTIN2* was amplified as a loading control using actin2-F (5'-CTGGATTCTGGTGATGGTGTGTC-3') and actin2-R (5'-TCTTTGCTCATAACGGTCAGCG-3') primers (An et al., 1996). PCR was performed with Takara *Ex Taq* polymerase (Takara Shuzo) for 35 cycles of denaturation for 30 s at 94°C, annealing for 30 s at 59°C, and extension for 1 min at 72°C, followed by final extension for 5 min at 72°C. PCR reaction for *ACTIN2* amplification was performed for 23 cycles.

RGL2 mRNA accumulation was analyzed by RT-PCR using total RNA extracted from whole seeds imbibed as indicated in Figure 7 using a phenol-sodium dodecyl sulfate extraction method (Sambrook et al., 1989). Genomic DNA contamination was removed, and cDNA strand synthesis was performed as described above. To determine the transcript level of *RGL2*, PCR was performed on cDNA using RGL2int-F (5'-CACGAGGAGAATCTCAACTTAG-3') and RGL2orf3R (5'-GGCGAGTTCCACGCGAGG-3') primers. The ubiquitin gene was amplified as a constitutive control as described by Tyler et al. (2004). Primer specificity was confirmed using the *rgl2-13* mutant (see Supplemental Figure 6B online). To determine the transcript level of the seven GA-regulated genes, PCR was performed on cDNA using the specific primers (see Supplemental Table 2 online). cDNA synthesis was performed on mRNA extracted from seeds imbibed first for 60 h at 4°C in water followed by 24 h at 22°C on MS-agar plates, and then seeds were incubated for 12 h at 22°C on MS-agar plates in the presence and absence of 10 μ M GA₃. PCR reactions were performed with Mango-Taq DNA polymerase (Bioline) for 30 cycles of denaturation for 1 min at 94°C, annealing for 1 min at 60°C, and extension for 1 min at 72°C, followed by final extension for 5 min at 72°C.

Protein Gel Blot Analysis

Total protein from whole seeds was extracted in extraction buffer (phosphate buffer with 1 \times protease inhibitor cocktail [Roche]), and 60 μ g protein was separated on an 8% SDS-PAGE and transferred onto a polyvinylidene difluoride membrane. The protein concentration was determined using Bradford reagent (Bio-Rad protein assay; Bio-Rad Laboratories). Time points for *RGL2* protein extraction were taken at 24-h intervals during imbibition as indicated in Figures 5 and 7. Protein detection was performed using an enhanced chemiluminescence system (ECL; GE Healthcare) according to the manufacturer's protocol. The polyclonal *RGL2* antibody (Hussain et al., 2005) was used as primary antibody (1:20,000). The anti-rabbit IgG-horseradish peroxidase (GE Healthcare) was used as a secondary antibody (1:200,000). The specificity of the *RGL2* antibody was confirmed by determining that the 60.5-kD *RGL2* band was absent in the *sly1-2 rgl2-13* double mutant after 24 h of imbibition at 22°C (see Supplemental Figures 4A and 4B online). To detect RGA protein, polyclonal SLR1 antibody (Itoh et al., 2002) was used as primary antibody (1:15,000). The same secondary antibody was used as described above. The specificity of the SLR1 antibody was confirmed by determining that the 64.0-kD RGA band was absent in the *ga1-3 rgl2-2* and *sly1-2 rga24* double mutants after 24 h of imbibition at 22°C (see Supplemental Figures 5A and 5B online). Time points for RGA protein extraction were taken at 24-h intervals during imbibition as indicated (see Supplemental Figure 5D online).

Accession Numbers

Arabidopsis Genome Initiative locus identifiers for the genes mentioned in this article are as follows: *Sly1* (At4g24210), *RGL2* (At3g03450), *EXP1*

(At1g69530), *XTH5* or *EXGT-A4* (At5g13870), *XTH31* (At3g44990), *AtCP1* (At4g36880), *PME* (At3g10720), *GASA4* (At5g15230), and *SCS10* (At2g27920).

Supplemental Data

The following materials are available in the online version of this article.

Supplemental Table 1. Phenotypes of Different *sly1* Mutant Alleles.

Supplemental Table 2. Primer Sequences of GA-Regulated Genes for RT-PCR.

Supplemental Figure 1. Time-Course After-ripening Experiment in Dormant *sly1-2*.

Supplemental Figure 2. *sly1* Mutant Seed Germination Is More Sensitive to PEG 8000.

Supplemental Figure 3. *sly1* Mutant Seed Germination Is More Sensitive to PAC.

Supplemental Figure 4. *RGL2* Protein Detection by Protein Gel Blot Analysis.

Supplemental Figure 5. RGA Protein Detection by Protein Gel Blot Analysis.

Supplemental Figure 6. Ubiquitin Control RT-PCR Reaction and Confirmation of Amplification in *RGL2* in Different *Arabidopsis* Ecotypes.

ACKNOWLEDGMENTS

We thank J. Peng and A. Hussain for kindly providing the *RGL2* antibody and *ga1-3 rga-t2* double mutant. We are grateful to M. Matsuoka and T.-p. Sun for providing the SLR antibody and *rgl2-13* mutant allele, respectively. We thank H. Nonogaki and members of the Steber lab for helpful suggestions and comments on the manuscript. This work was supported by a Japanese Society for the Promotion of Science Postdoctoral Fellowship for Research Abroad to T.A. and by USDA-Agricultural Research Service funds and USDA National Research Initiative 2002-01351 to C.M.S.

Received October 5, 2006; revised February 15, 2007; accepted March 5, 2007; published March 23, 2007.

REFERENCES

- Alonso-Blanco, C., Bentsink, L., Hanhart, C.J., Blankestijn-de Vries, H., and Koornneef, M. (2003). Analysis of natural allelic variation at seed dormancy loci of *Arabidopsis thaliana*. *Genetics* **164**: 711–729.
- An, Y.Q., McDowell, J.M., Huang, S., McKinney, E.C., Chambliss, S., and Meagher, R.B. (1996). Strong, constitutive expression of the *Arabidopsis ACT2/ACT8* actin subclass in vegetative tissues. *Plant J.* **10**: 107–121.
- Bassel, G.W., Zielinska, E., Mullen, R.T., and Bewley, J.D. (2004). Down-regulation of DELLA genes is not essential for germination in tomato, soybean, and *Arabidopsis* seeds. *Plant Physiol.* **136**: 2782–2789.
- Bewley, J.D. (1997). Seed germination and dormancy. *Plant Cell* **9**: 1055–1066.
- Bewley, J.D., and Black, M. (1994). *Seeds: Physiology of Development and Germination*. (New York: Plenum Press).
- Cao, D., Cheng, H., Wu, W., Meng Soo, H., and Peng, J. (2006). Gibberellin mobilizes distinct DELLA-dependent transcriptomes to

- regulate seed germination and floral development in *Arabidopsis*. *Plant Physiol.* **142**: 509–525.
- Cao, D., Hussain, A., Cheng, H., and Peng, J.** (2005). Loss of function of four DELLA genes leads to light- and gibberellin-independent seed germination in *Arabidopsis*. *Planta* **223**: 105–113.
- Cheng, H., Qin, L., Lee, S., Fu, X., Richards, D.E., Cao, D., Luo, D., Harberd, N.P., and Peng, J.** (2004). Gibberellin regulates *Arabidopsis* floral development via suppression of DELLA protein function. *Development* **131**: 1055–1064.
- Debeaujon, I., and Koornneef, M.** (2000). Gibberellin requirement for *Arabidopsis* seed germination is determined both by testa characteristics and embryonic abscisic acid. *Plant Physiol.* **122**: 415–424.
- Debeaujon, I., Leon-Kloosterziel, K.M., and Koornneef, M.** (2000). Influence of the testa on seed dormancy, germination, and longevity in *Arabidopsis*. *Plant Physiol.* **122**: 403–414.
- Dill, A., and Sun, T.** (2001). Synergistic derepression of gibberellin signaling by removing RGA and GAI function in *Arabidopsis thaliana*. *Genetics* **159**: 777–785.
- Dill, A., Thomas, S.G., Hu, J., Steber, C.M., and Sun, T.P.** (2004). The *Arabidopsis* F-box protein *SLEEPY1* targets gibberellin signaling repressors for gibberellin-induced degradation. *Plant Cell* **16**: 1392–1405.
- Finkelstein, R.R., and Rock, C.D.** (September 30, 2002). Abscisic acid biosynthesis and response. In *The Arabidopsis Book*, C.R. Somerville and E.M. Meyerowitz, eds (Rockville, MD: American Society of Plant Biologists), doi/10.1199/tab.0058, <http://www.aspb.org/publications/arabidopsis/>.
- Footitt, S., Slocumbe, S.P., Lerner, V., Kurup, S., Wu, Y., Larson, T., Graham, I., Baker, A., and Holdsworth, M.** (2002). Control of germination and lipid mobilization by *COMATOSE*, the *Arabidopsis* homologue of human ALDP. *EMBO J.* **21**: 2912–2922.
- Fu, X., Richards, D.E., Fleck, B., Xie, D., Burton, N., and Harberd, N.P.** (2004). The *Arabidopsis* mutant *sleepy1gar2-1* protein promotes plant growth by increasing the affinity of the SCF^{S_{LY}1} E3 ubiquitin ligase for DELLA protein substrates. *Plant Cell* **16**: 1406–1418.
- Griffiths, J., Murase, K., Rieu, I., Zentella, R., Zhang, Z.L., Powers, S.J., Gong, F., Phillips, A.L., Hedden, P., Sun, T.P., and Thomas, S.G.** (2006). Genetic characterization and functional analysis of the GID1 gibberellin receptors in *Arabidopsis*. *Plant Cell* **18**: 3399–3414.
- Hartweck, L.M., and Olszewski, N.E.** (2006). Rice *GIBBERELLIN INSENSITIVE DWARF1* is a gibberellin receptor that illuminates and raises questions about GA signaling. *Plant Cell* **18**: 278–282.
- Herzog, M., Dorne, A.M., and Grellet, F.** (1995). *GASA*, a gibberellin-regulated gene family from *Arabidopsis thaliana* related to the tomato *GAST1* gene. *Plant Mol. Biol.* **27**: 743–752.
- Hussain, A., Cao, D., Cheng, H., Wen, Z., and Peng, J.** (2005). Identification of the conserved serine/threonine residues important for gibberellin sensitivity of *Arabidopsis* *RGL2* protein. *Plant J.* **44**: 88–99.
- Itoh, H., Ueguchi-Tanaka, M., Sato, Y., Ashikari, M., and Matsuoka, M.** (2002). The gibberellin signaling pathway is regulated by the appearance and disappearance of SLENDER RICE1 in nuclei. *Plant Cell* **14**: 57–70.
- Jacobsen, S.E., and Olszewski, N.E.** (1993). Mutations at the *SPINDLY* locus of *Arabidopsis* alter gibberellin signal transduction. *Plant Cell* **5**: 887–896.
- Keith, K., Kraml, M., Dengler, N.G., and McCourt, P.** (1994). *fusca3*: A heterochronic mutation affecting late embryo development in *Arabidopsis*. *Plant Cell* **6**: 589–600.
- King, K.E., Moritz, T., and Harberd, N.P.** (2001). Gibberellins are not required for normal stem growth in *Arabidopsis thaliana* in the absence of GAI and RGA. *Genetics* **159**: 767–776.
- Koornneef, M., and Karssen, C.M.** (1994). Seed dormancy and germination. In *Arabidopsis*, C.R. Somerville and E.M. Meyerowitz, eds (Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press), pp. 313–334.
- Krysan, P.J., Young, J.C., and Sussman, M.R.** (1999). T-DNA as an insertional mutagen in *Arabidopsis*. *Plant Cell* **11**: 2283–2290.
- Kushiro, T., Okamoto, M., Nakabayashi, K., Yamagishi, K., Kitamura, S., Asami, T., Hirai, N., Koshiba, T., Kamiya, Y., and Nambara, E.** (2004). The *Arabidopsis* cytochrome P450 CYP707A encodes ABA 8'-hydroxylases: Key enzymes in ABA catabolism. *EMBO J.* **23**: 1647–1656.
- Lee, S., Cheng, H., King, K.E., Wang, W., He, Y., Hussain, A., Lo, J., Harberd, N.P., and Peng, J.** (2002). Gibberellin regulates *Arabidopsis* seed germination via *RGL2*, a *GAI/RGA*-like gene whose expression is up-regulated following imbibition. *Genes Dev.* **16**: 646–658.
- McGinnis, K.M., Thomas, S.G., Soule, J.D., Strader, L.C., Zale, J.M., Sun, T.P., and Steber, C.M.** (2003). The *Arabidopsis* *SLEEPY1* gene encodes a putative F-box subunit of an SCF E3 ubiquitin ligase. *Plant Cell* **15**: 1120–1130.
- Monte, E., Amador, V., Russo, E., Martínez-García, J., and Prat, S.** (2003). PHOR1: A U-Box GA signaling component with a role in proteasome degradation? *J. Plant Growth Regul.* **22**: 152–162.
- Nakajima, M., et al.** (2006). Identification and characterization of *Arabidopsis* gibberellin receptors. *Plant J.* **46**: 880–889.
- Nambara, E., Keith, K., McCourt, P., and Naito, S.** (1995). A regulatory role for the *ABI3* gene in the establishment of embryo maturation in *Arabidopsis thaliana*. *Development* **121**: 629–636.
- Ogawa, M., Hanada, A., Yamauchi, Y., Kuwahara, A., Kamiya, Y., and Yamaguchi, S.** (2003). Gibberellin biosynthesis and response during *Arabidopsis* seed germination. *Plant Cell* **15**: 1591–1604.
- Okamoto, M., Kuwahara, A., Seo, M., Kushiro, T., Asami, T., Hirai, N., Kamiya, Y., Koshiba, T., and Nambara, E.** (2006). CYP707A1 and CYP707A2, which encode abscisic acid 8'-hydroxylases, are indispensable for proper control of seed dormancy and germination in *Arabidopsis*. *Plant Physiol.* **141**: 97–107.
- Olszewski, N., Sun, T.P., and Gubler, F.** (2002). Gibberellin signaling: Biosynthesis, catabolism, and response pathways. *Plant Cell* **14** (suppl.): S61–S80.
- Ooms, J.J.J., Leon-Kloosterziel, K.M., Bartels, D., Koornneef, M., and Karssen, C.M.** (1993). Acquisition of desiccation tolerance and longevity in seeds of *Arabidopsis thaliana*. *Plant Physiol.* **102**: 1185–1191.
- Papi, M., Sabatini, S., Altamura, M.M., Hennig, L., Schafer, E., Costantino, P., and Vittorioso, P.** (2002). Inactivation of the phloem-specific Dof zinc finger gene *DAG1* affects response to light and integrity of the testa of *Arabidopsis* seeds. *Plant Physiol.* **128**: 411–417.
- Papi, M., Sabatini, S., Bouchez, D., Camilleri, C., Costantino, P., and Vittorioso, P.** (2000). Identification and disruption of an *Arabidopsis* zinc finger gene controlling seed germination. *Genes Dev.* **14**: 28–33.
- Parcy, F., and Giraudat, J.** (1997). Interactions between the *ABI1* and the ectopically expressed *ABI3* genes in controlling abscisic acid responses in *Arabidopsis* vegetative tissues. *Plant J.* **11**: 693–702.
- Penfield, S., Li, Y., Gilday, A.D., Graham, S., and Graham, I.A.** (2006). *Arabidopsis* *ABA INSENSITIVE4* regulates lipid mobilization in the embryo and reveals repression of seed germination by the endosperm. *Plant Cell* **18**: 1887–1899.
- Peng, J., et al.** (1999). 'Green revolution' genes encode mutant gibberellin response modulators. *Nature* **400**: 256–261.
- Pysh, L.D., Wysocka-Diller, J.W., Camilleri, C., Bouchez, D., and Benfey, P.N.** (1999). The GRAS gene family in *Arabidopsis*: Sequence characterization and basic expression analysis of the *SCARECROW-LIKE* genes. *Plant J.* **18**: 111–119.

- Russell, L., Lerner, V., Kurup, S., Bougourd, S., and Holdsworth, M.** (2000). The Arabidopsis *COMATOSE* locus regulates germination potential. *Development* **127**: 3759–3767.
- Sambrook, J., Fritsch, E.F., and Maniatis, T.** (1989). *Molecular Cloning: A Laboratory Manual*. (Cold Spring Harbor, NY: Cold Spring Harbor Laboratory Press).
- Silverstone, A.L., Jung, H.S., Dill, A., Kawaide, H., Kamiya, Y., and Sun, T.P.** (2001). Repressing a repressor: Gibberellin-induced rapid reduction of the RGA protein in Arabidopsis. *Plant Cell* **13**: 1555–1566.
- Steber, C.M., Cooney, S.E., and McCourt, P.** (1998). Isolation of the GA-response mutant *sly1* as a suppressor of *ABI1-1* in *Arabidopsis thaliana*. *Genetics* **149**: 509–521.
- Steber, C.M., and McCourt, P.** (2001). A role for brassinosteroids in germination in Arabidopsis. *Plant Physiol.* **125**: 763–769.
- Strader, L.C., Ritchie, S., Soule, J.D., McGinnis, K.M., and Steber, C.M.** (2004). Recessive-interfering mutations in the gibberellin signaling gene *SLEEPY1* are rescued by overexpression of its homologue, *SNEEZY*. *Proc. Natl. Acad. Sci. USA* **101**: 12771–12776.
- Sundaresan, V., Springer, P., Volpe, T., Haward, S., Jones, J.D., Dean, C., Ma, H., and Martienssen, R.** (1995). Patterns of gene action in plant development revealed by enhancer trap and gene trap transposable elements. *Genes Dev.* **9**: 1797–1810.
- Thomas, S.G., Rieu, I., and Steber, C.M.** (2005). Gibberellin metabolism and signaling. *Vitam. Horm.* **72**: 289–338.
- Tyler, L., Thomas, S.G., Hu, J., Dill, A., Alonso, J.M., Ecker, J.R., and Sun, T.P.** (2004). DELLA proteins and gibberellin-regulated seed germination and floral development in Arabidopsis. *Plant Physiol.* **135**: 1008–1019.
- Ueguchi-Tanaka, M., Ashikari, M., Nakajima, M., Itoh, H., Katoh, E., Kobayashi, M., Chow, T.Y., Hsing, Y.I.C., Kitano, H., Yamaguchi, I., and Matsuoka, M.** (2005). *GIBBERELLIN INSENSITIVE DWARF1* encodes a soluble receptor for gibberellin. *Nature* **437**: 693–698.
- Yamaguchi, S., Kamiya, Y., and Sun, T.** (2001). Distinct cell-specific expression patterns of early and late gibberellin biosynthetic genes during Arabidopsis seed germination. *Plant J.* **28**: 443–453.
- Yamauchi, Y., Ogawa, M., Kuwahara, A., Hanada, A., Kamiya, Y., and Yamaguchi, S.** (2004). Activation of gibberellin biosynthesis and response pathways by low temperature during imbibition of *Arabidopsis thaliana* seeds. *Plant Cell* **16**: 367–378.

**Seed Germination of GA-Insensitive *sleepy1* Mutants Does Not Require RGL2 Protein
Disappearance in *Arabidopsis***

Tohru Ariizumi and Camille M. Steber

Plant Cell 2007;19;791-804; originally published online March 23, 2007;

DOI 10.1105/tpc.106.048009

This information is current as of January 16, 2018

Supplemental Data	/content/suppl/2007/03/20/tpc.106.048009.DC1.html
References	This article cites 49 articles, 37 of which can be accessed free at: /content/19/3/791.full.html#ref-list-1
Permissions	https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&issn=1532298X&WT.mc_id=pd_hw1532298X
eTOCs	Sign up for eTOCs at: http://www.plantcell.org/cgi/alerts/ctmain
CiteTrack Alerts	Sign up for CiteTrack Alerts at: http://www.plantcell.org/cgi/alerts/ctmain
Subscription Information	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: http://www.aspb.org/publications/subscriptions.cfm