

## REVIEW

# Calcium Signals: The Lead Currency of Plant Information Processing

Jörg Kudla,<sup>1</sup> Oliver Batistič, and Kenji Hashimoto

Institut für Botanik, Universität Münster, 48149 Münster, Germany

**Ca<sup>2+</sup> signals are core transducers and regulators in many adaptation and developmental processes of plants. Ca<sup>2+</sup> signals are represented by stimulus-specific signatures that result from the concerted action of channels, pumps, and carriers that shape temporally and spatially defined Ca<sup>2+</sup> elevations. Cellular Ca<sup>2+</sup> signals are decoded and transmitted by a toolkit of Ca<sup>2+</sup> binding proteins that relay this information into downstream responses. Major transduction routes of Ca<sup>2+</sup> signaling involve Ca<sup>2+</sup>-regulated kinases mediating phosphorylation events that orchestrate downstream responses or comprise regulation of gene expression via Ca<sup>2+</sup>-regulated transcription factors and Ca<sup>2+</sup>-responsive promoter elements. Here, we review some of the remarkable progress that has been made in recent years, especially in identifying critical components functioning in Ca<sup>2+</sup> signal transduction, both at the single-cell and multicellular level. Despite impressive progress in our understanding of the processing of Ca<sup>2+</sup> signals during the past years, the elucidation of the exact mechanistic principles that underlie the specific recognition and conversion of the cellular Ca<sup>2+</sup> currency into defined changes in protein–protein interaction, protein phosphorylation, and gene expression and thereby establish the specificity in stimulus response coupling remain to be explored.**

## INTRODUCTION

Calcium (Ca<sup>2+</sup>) likely represents the most versatile ion in eukaryotic organisms. It is involved in nearly all aspects of plant development and participates in many regulatory processes. Because of its flexibility in exhibiting different coordination numbers and complex geometries, Ca<sup>2+</sup> can easily form complexes with proteins, membranes, and organic acids. On the one hand, this feature renders Ca<sup>2+</sup> a toxic cellular compound at higher concentrations because it would readily form insoluble complexes with phosphate (as present in ATP), but on the other hand, the required tight spatial and temporal control of cellular Ca<sup>2+</sup> concentration may have paved the way for the evolutionary emergence of Ca<sup>2+</sup> signaling.

Considerable interest and research on this ion has been sparked by the apparent antagonism between the obvious cellular abundance of Ca<sup>2+</sup> in certain organelles and cell structures and its required rareness in the cytoplasm. Since the first report in the green algae *Chara* that changes of cytosolic Ca<sup>2+</sup> indicate a function of Ca<sup>2+</sup> as a second messenger in plants (Williamson and Ashley, 1982), transient elevations in cytosolic Ca<sup>2+</sup> concentration have been documented to be involved in a multitude of physiological processes, including responses to abiotic stresses, hormones, and pathogens. During the last two decades of the 20th century, advances in Ca<sup>2+</sup> monitoring techniques have allowed detailed analyses of cellular Ca<sup>2+</sup> dynamics. Several groups reported that defined changes of cytosolic Ca<sup>2+</sup> concentration are triggered by cellular second

messengers, such as NAADP, IP<sub>3</sub>, IP<sub>6</sub>, Sphingosine-1-Phosphate, and cADPR (Drøbak and Ferguson, 1985; Schumaker and Sze, 1987; Blatt et al., 1990; Gilroy et al., 1990; Allen and Sanders, 1995; Navazio et al., 2000; Lemtiri-Chlieh et al., 2003), and it became evident that the identity and intensity of a specific stimulus impulse results in stimulus-specific and dynamic alterations of cytosolic Ca<sup>2+</sup> concentration (Allen et al., 1995; McAinsh et al., 1995). This heterogeneity of increases in cytosolic-free Ca<sup>2+</sup> ion concentration in terms of duration, amplitude, frequency, and spatial distribution lead A.M. Hetherington and coworkers to formulate the concept of “Ca<sup>2+</sup> signatures” (Webb et al., 1996). Herein, signal information would be encoded by a specific Ca<sup>2+</sup> signature that is defined by precise control of spatial, temporal, and concentration parameters of alterations in cytosolic Ca<sup>2+</sup> concentration. The spectrum of stimuli that evoke such Ca<sup>2+</sup> elevations and their stimulus-specific characteristics has been cataloged and critically discussed in a number of informative reviews (Rudd and Franklin-Tong, 1999; Sanders et al., 1999; Knight and Knight, 2001; Sanders et al., 2002; Scrase-Field and Knight, 2003). Subsequent research suggested that while the shape and spatio-temporal distribution of Ca<sup>2+</sup> elevations could be of critical importance for stimulus response coupling (Allen et al., 2001), an additional level of regulation and specificity is achieved by Ca<sup>2+</sup> binding proteins that function as signal sensor proteins (Batistič and Kudla, 2004). These proteins decode and relay the information encoded by Ca<sup>2+</sup> signatures into specific protein–protein interactions, defined phosphorylation cascades, or transcriptional responses (Luan et al., 2002; Sanders et al., 2002; Finkler et al., 2007a). Consequently, the dynamic interplay between Ca<sup>2+</sup> signatures and Ca<sup>2+</sup> sensing proteins contributes to generating stimulus specificity of Ca<sup>2+</sup>

<sup>1</sup> Address correspondence to jkudla@uni-muenster.de.  
www.plantcell.org/cgi/doi/10.1105/tpc.109.072686

signaling. Since the principles and cellular tool kits of  $\text{Ca}^{2+}$  signaling were last reviewed in this journal (Luan et al., 2002; Sanders et al., 2002), remarkable progress has been achieved especially in elucidating the mechanisms that contribute to decoding of  $\text{Ca}^{2+}$  signals, and complete  $\text{Ca}^{2+}$ -triggered regulatory modules have been identified. In this review, we will focus on the description of scientific insights and the discussion of emerging concepts that have been arising over the past few years.

## FUNCTIONS OF $\text{Ca}^{2+}$ SIGNALING


$\text{Ca}^{2+}$  is involved in various responses to abiotic and biotic stimuli, including light, high and low temperature, touch, salt and drought, osmotic stress, plant hormones, fungal elicitors, and nodulation factors (Sanders et al., 1999). These stimuli induce a distinct spatio-temporal pattern of changes in cytosolic-free  $\text{Ca}^{2+}$  concentration ( $[\text{Ca}^{2+}]_{\text{cyt}}$ ). Single-cell systems, such as guard cells, growing pollen tubes, or root hairs, represent excellent models to investigate primary and autonomous  $\text{Ca}^{2+}$  responses. However, the final response of the plant to external stimuli is manifested by regulation of complex growth processes in distinct tissues and organs. Concurrently to the diversity of stimulus-specific  $\text{Ca}^{2+}$  signatures at the single-cell level, differentiation gives rise to another layer of cell type-specific  $\text{Ca}^{2+}$  responses in tissues or organs. This additional level of complex-

ity may contribute to more diversity in local or systemic responses. Therefore, research on plant  $\text{Ca}^{2+}$  signaling has taken advantage of single-cell model systems but in parallel moves forward to elucidate  $\text{Ca}^{2+}$  dynamics in the tissue context and in the whole organism. Consequently, here, we review and discuss how  $\text{Ca}^{2+}$  signatures contribute to signaling processes at the single-cell level and the multicellular level.

## $\text{Ca}^{2+}$ Signaling at the Single Cell Level

### Stomatal Closure and Opening

In *Arabidopsis thaliana* guard cells, stimuli for stomatal closure, including abscisic acid (ABA), hydrogen peroxide, cold, elevation of external  $\text{Ca}^{2+}$ , and atmospheric  $\text{CO}_2$ , induce cytosolic  $\text{Ca}^{2+}$  oscillations (Allen et al., 2001; Young et al., 2006). Artificially imposed  $\text{Ca}^{2+}$  oscillations have revealed that cytosolic  $\text{Ca}^{2+}$  regulates stomatal closure by two mechanisms: short-term  $\text{Ca}^{2+}$ -reactive closure and long-term  $\text{Ca}^{2+}$ -programmed closure (Figure 1A) (Allen et al., 2001; Sanders et al., 2002). The short-term  $\text{Ca}^{2+}$ -reactive closure is a rapid reaction to exceeding a certain level of  $[\text{Ca}^{2+}]_{\text{cyt}}$  elevation and is irrespective of the elevation pattern ( $\text{Ca}^{2+}$  functions likely in a threshold manner). Meanwhile, long-term  $\text{Ca}^{2+}$ -programmed closure (i.e., prevention of stomatal reopening) is controlled by  $\text{Ca}^{2+}$  oscillations within a defined range of amplitude, frequency, duration, and



**Figure 1.**  $\text{Ca}^{2+}$  Signaling in Guard Cell Regulation.

**(A)** Schematic representation of artificially imposed  $\text{Ca}^{2+}$  oscillations/plateau and the corresponding temporal changes in stomatal aperture. The long-term  $\text{Ca}^{2+}$ -programmed closure is caused by  $\text{Ca}^{2+}$  oscillations with a defined pattern ( $\text{Ca}^{2+}$  signature), whereas the short-term  $\text{Ca}^{2+}$ -reactive closure is induced by  $[\text{Ca}^{2+}]_{\text{cyt}}$  elevation, regardless of the pattern.

**(B)** Simplified model for  $\text{Ca}^{2+}$  controlled stomatal closure and opening. Stomatal closing stimuli, such as ABA, high  $\text{CO}_2$ , and cold, induce spontaneous  $\text{Ca}^{2+}$  oscillations, resulting in  $\text{Ca}^{2+}$ -reactive closure and subsequent  $\text{Ca}^{2+}$ -programmed closure. On the other hand, low  $\text{CO}_2$ -induced  $\text{Ca}^{2+}$  oscillations result not in  $\text{Ca}^{2+}$ -reactive closure but stomatal opening, implying that the stimulus may desensitize the signaling pathways of the  $\text{Ca}^{2+}$ -reactive closure ( $\text{Ca}^{2+}$  sensitivity priming hypothesis). It is unknown whether low  $\text{CO}_2$ -induced  $\text{Ca}^{2+}$  oscillation itself encrypts specific information of stomatal opening.

overall transient number ( $\text{Ca}^{2+}$  functions as a signature). Recent studies have revealed several elements involved in these mechanisms. The *Arabidopsis*  $\text{Ca}^{2+}$ -dependent protein kinase (CDPK) double mutant *cpk3 cpk6* exhibits impaired activation of S-type anion channels in response to cytosolic  $\text{Ca}^{2+}$ . Guard cells in the double mutant are defective in short-term closure but not in long-term closure (Mori et al., 2006). Mutation of the SLOW ANION CHANNEL-ASSOCIATED1 guard cell anion efflux channel abrogates  $\text{Ca}^{2+}$ -reactive stomatal closure and impairs stomatal responses to  $\text{CO}_2$ , ABA, ozone, light/dark transitions, humidity change, calcium ions, hydrogen peroxide, and nitric oxide (Negi et al., 2008; Vahisalu et al., 2008). By contrast, overexpression of At GLR3.1, a glutamate (Glu) receptor homolog, causes a defect in long-term closure but not in short-term closure (Cho et al., 2009). These findings indicate that the short-term and long-term responses are functionally separable. Interestingly, the translational inhibitor cycloheximide partially inhibits long-term closure but not short-term closure, suggesting no requirement of de novo protein synthesis in the short-term response (Cho et al., 2009). Taken together, these findings imply that certain threshold levels of  $\text{Ca}^{2+}$  elevations activate the preexisting cellular machinery, including  $\text{Ca}^{2+}$  sensor proteins and ion channels, to regulate rapid stomatal closure. Moreover, defined patterns of  $\text{Ca}^{2+}$  oscillations likely activate preexisting proteins and also induce the expression of required genes, resulting in inhibition of stomatal reopening (Figure 1A).

However, it is important to note that nonoscillatory increases in cytosolic  $\text{Ca}^{2+}$  are also important features of stomatal responses that result in stomatal closure and that the actual pattern of oscillations evoked by a specific stimulus exhibits some variation and is not "tight" (Hetherington and Brownlee, 2004). Consequently, the  $\text{Ca}^{2+}$  decoding machinery that translates these  $\text{Ca}^{2+}$  elevations and oscillations into defined downstream responses must be able to decode these variable oscillations (Hetherington and Brownlee, 2004).

Surprisingly, recent studies have shown that  $\text{Ca}^{2+}$  SENSING-RECEPTOR (CAS), a thylakoid membrane-localized protein, is crucial for the  $\text{Ca}^{2+}$  response, suggesting involvement of chloroplasts in  $[\text{Ca}^{2+}]_{\text{cyt}}$ -mediated stomatal closure (Nomura et al., 2008; Weinl et al., 2008). The elucidation of the functional interplay between  $\text{Ca}^{2+}$  releases from different intra- and extracellular sources certainly represents an important goal to further our understanding of guard cell signaling.

$\text{Ca}^{2+}$  signaling likely also contributes to the regulation of stomatal opening (Irving et al., 1992; Schroeder et al., 2001). For example, low atmospheric  $\text{CO}_2$ , which mediates stomatal opening, induces high-frequency  $\text{Ca}^{2+}$  oscillation. Treatment with the cytosolic  $\text{Ca}^{2+}$  chelator BAPTA-AM abolishes the oscillation and attenuates the stomatal opening in response to low  $\text{CO}_2$  (Young et al., 2006). It would be interesting to investigate whether the rapid oscillation pattern encrypts specific information. As described above, artificially imposed  $\text{Ca}^{2+}$  oscillations induce the short-term  $\text{Ca}^{2+}$ -reactive closure regardless of their pattern. However, the rapid  $\text{Ca}^{2+}$  oscillation induced by low  $\text{CO}_2$  does not cause such reaction (Young et al., 2006). A model was derived in which low  $\text{CO}_2$  stimuli may negatively modulate  $\text{Ca}^{2+}$  sensitivities of signaling components, which regulate the short-term closure, such as CDPKs. On the other hand, during the

stomatal closure response, preexposure to ABA enhances the response of S-type anion channels to  $[\text{Ca}^{2+}]_{\text{cyt}}$  elevation, implying that ABA may positively modulate sensitivities of  $\text{Ca}^{2+}$  sensor proteins (Siegel et al., 2009). A  $\text{Ca}^{2+}$  sensitivity priming model has been proposed as a hypothesis to explain these observations in the context of mechanisms that contribute to the generation of specificity in  $\text{Ca}^{2+}$  signaling (Figure 1B) (Young et al., 2006).

### The Establishment of Symbiosis in Root Hairs

In legume root hair cells, exposure to rhizobial-derived nodulation (Nod) factors induces biphasic changes  $[\text{Ca}^{2+}]_{\text{cyt}}$  that comprise an initial  $\text{Ca}^{2+}$  influx and a subsequent (10 to 20 min later) long-term  $\text{Ca}^{2+}$  oscillation (also designated as  $\text{Ca}^{2+}$  spiking) in the perinucleus (Shaw and Long, 2003). The *Medicago truncatula* mutants *does not make infection1 (dmi1)* and *dmi2* are defective in the  $\text{Ca}^{2+}$  spiking but retain the initial  $\text{Ca}^{2+}$  influx. Furthermore, low concentration ( $10^{-11}$  to  $10^{-12}$  M) of Nod factor induces  $\text{Ca}^{2+}$  spiking but not  $\text{Ca}^{2+}$  influx, suggesting that they are separable responses (Shaw and Long, 2003). Induction of *Early Nodulation 11 (ENOD11)* is impaired in the  $\text{Ca}^{2+}$  spiking-defective mutants *dmi1* and *dmi2*, as well as in *dmi3*, which is defective in a gene encoding  $\text{Ca}^{2+}$  calmodulin-dependent kinase (CCaMK). Remarkably, specific removal of the autoinhibitory domain of this kinase leads to autoactivation of the downstream nodulation signaling pathway with the resultant induction of nodules and nodulation gene expression in the absence of bacterial elicitation (Gleason et al., 2006). This demonstrates not only the essential function of this CCaMK in the regulation of nodule development but also indicates that the activation of this kinase through the oscillatory  $\text{Ca}^{2+}$  signal is necessary and sufficient to elicit this process (Gleason et al., 2006). Although the mechanism of generation of  $\text{Ca}^{2+}$  spikes is not well understood, some blockers for  $\text{Ca}^{2+}$  channels and  $\text{Ca}^{2+}$  pumps have been shown to inhibit both  $\text{Ca}^{2+}$  spiking (Engstrom et al., 2002) and *ENOD11* expression (Charron et al., 2004). This evidence suggests that  $\text{Ca}^{2+}$  spiking is essential for regulation of nodulation, raising the question of how the spiking could encode specific information. Miwa et al. (2006) proposed a correlation between the number of  $\text{Ca}^{2+}$  spikes and *ENOD11* expression levels. In this work, *ENOD11* inductions were observed only when the  $\text{Ca}^{2+}$  spiking lasted for at least 60 min. The average period between each spike was  $\sim 100$  s; therefore, the authors estimated that a minimum of  $\sim 36$  spikes is required for *ENOD11* induction. Jasmonic acid treatment lengthens the period between each  $\text{Ca}^{2+}$  spike but does not affect the required number of spikes for *ENOD11* expression, further supporting the notion that the number of  $\text{Ca}^{2+}$  spikes functions to encode information (Miwa et al., 2006).

$\text{Ca}^{2+}$  also plays important roles in the symbiosis of legumes with arbuscular mycorrhizal fungi. Incubation with mycorrhizae induces  $\text{Ca}^{2+}$  spiking in legume root hair cells.  $\text{Ca}^{2+}$  spiking is abolished in *M. truncatula dmi1* and *dmi2*, suggesting that nodulation and mycorrhizal infection involve common signaling components. Despite the overlap, mycorrhizal-induced  $\text{Ca}^{2+}$  spiking exhibits a shorter period and smaller amplitudes compared with that in the Nod factor response (Kosuta et al., 2008).

Remarkably, bacterial elicitors of plant pathogens also induce a  $\text{Ca}^{2+}$  release in the cytosol and nucleus. Here, the nitric oxide signal following elicitor application is important for the  $\text{Ca}^{2+}$  release in the cytosol, but it does not trigger a nuclear  $\text{Ca}^{2+}$  response (Lamotte et al., 2004). The nucleus also exhibits specific calcium signals in response to different elicitors. Harpin and flagellin resulted in a different  $\text{Ca}^{2+}$  release than observed in response to carbohydrate elicitors such as oligogalacturonides. Remarkably, the cytosolic  $\text{Ca}^{2+}$  response to these elicitors was comparable (Lecourieux et al., 2005). These observations suggest that the nucleus does indeed harbor an independent  $\text{Ca}^{2+}$  machinery that may involve P-ATPases and nucleotide gated channels located at the inner membrane of the nucleus to regulate the nuclear  $\text{Ca}^{2+}$  reservoir (Mazars et al., 2009).

#### Tip Growth in Pollen Tubes and Root Hair Cells

Pollen tubes are one of the most extensively studied tip-growing model systems. In pollen tubes,  $[\text{Ca}^{2+}]_{\text{cyt}}$  is highly concentrated at the apex by an extracellular influx (Sanders et al., 1999; Hepler et al., 2001). In the  $\text{Ca}^{2+}$  gradient,  $[\text{Ca}^{2+}]_{\text{cyt}}$  oscillates with a lag phase behind the corresponding growth rate oscillation, suggesting that stretch-activated  $\text{Ca}^{2+}$  channels may be involved in the maintenance of the  $\text{Ca}^{2+}$  gradient (Dutta and Robinson, 2004). In accordance with an essential function of stretch-activated channels, pharmacological inhibition of channel activity disrupts the  $\text{Ca}^{2+}$  influx at the apex and terminates pollen tube elongation (Picton and Steer, 1985). Moreover, *Petunia inflata* and *M. truncatula* mutant lines that are defective in CDPKs exhibit loss of polarity (Ivashuta et al., 2005; Yoon et al., 2006). Therefore, the formation of a  $\text{Ca}^{2+}$  gradient as well as the decoding of  $\text{Ca}^{2+}$  signals by  $\text{Ca}^{2+}$  sensing proteins appear to play important roles in generating spatial determinants for pollen tube growth. Moreover, several studies have suggested a close interaction of intracellular  $\text{Ca}^{2+}$  and the cytoskeleton in that the growth regulatory effect of this ion is exerted by  $\text{Ca}^{2+}$ -dependent regulation of the structure and activity of F-actin (Snowman et al., 2002; Cardenas et al., 2008).

Recent studies have also revealed the importance of  $\text{Ca}^{2+}$  signaling in the root hair single-cell model system. Similar to growing pollen, root hair cells exhibit a tip-focused  $\text{Ca}^{2+}$  gradient with  $\text{Ca}^{2+}$  oscillation lagging behind growth oscillation (Monshausen et al., 2008). Moreover, the  $\text{Ca}^{2+}$  oscillation is followed by oscillation of apoplastic reactive oxygen species (ROS) production (Monshausen et al., 2007, 2008). Intriguingly, the RHD2 NADPH oxidase (also known as RBOH C) is localized to the plasma membrane of the tip-growing site and is required for appropriate growth of root hair cells (Takeda et al., 2008). RHD2 contains an EF hand-like  $\text{Ca}^{2+}$  binding domain and phosphorylation sites that appear to be targets of calcium-regulated protein kinases (Takeda et al., 2008). Therefore, the  $\text{Ca}^{2+}$  oscillation may regulate RHD2 activity. Inhibition of the  $\text{Ca}^{2+}$  oscillation by  $\text{La}^{3+}$  causes bursting growth, while increasing the  $\text{Ca}^{2+}$  gradient by  $\text{Ca}^{2+}$  ionophore arrests growth of root hairs (Monshausen et al., 2008). Consequently, it has been proposed that  $\text{Ca}^{2+}$  and ROS compose a positive feedback loop to sustain tip growth of root hair cells (Takeda et al., 2008).

## $\text{Ca}^{2+}$ Signaling at the Multicellular Level

### Abiotic Stress Responses

In the endodermis and the pericycle of *Arabidopsis* roots, salt stress causes a biphasic  $\text{Ca}^{2+}$  response consisting of an initial transient elevation and a subsequent oscillation that differs in phase and/or period in individual cells (Kiegle et al., 2000). By contrast, only a monophasic  $[\text{Ca}^{2+}]_{\text{cyt}}$  elevation is induced in the epidermis and the cortex (Kiegle et al., 2000). Interestingly, the initial  $\text{Ca}^{2+}$  response in the pericycle occurs a few seconds later than that in the other cell types. Moreover, the magnitude in the pericycle is significantly lower. This is likely because the endodermis limits the flow of water and ions to the xylem. Accordingly, whole-plant  $[\text{Ca}^{2+}]_{\text{cyt}}$  measurements have suggested a direct correlation between the strength of NaCl stress and the magnitude of  $[\text{Ca}^{2+}]_{\text{cyt}}$  elevation (Tracy et al., 2008). The function of the subsequent  $\text{Ca}^{2+}$  oscillation in the endodermis and the pericycle remains to be elucidated.

In contrast with salt stress, cold stress causes only monophasic  $[\text{Ca}^{2+}]_{\text{cyt}}$  elevation in the four types of *Arabidopsis* root cells without significant temporal difference, suggesting that all cells sense temperature changes simultaneously (Kiegle et al., 2000). The magnitude of cold-induced  $\text{Ca}^{2+}$  responses is dependent on the cooling rate ( $\Delta T/\Delta t$ ) (Plieth et al., 1999). However, the pericycle exhibits a more pronounced  $\text{Ca}^{2+}$  response to cold stress (Kiegle et al., 2000), suggesting that different cell types harbor different  $\text{Ca}^{2+}$  homeostasis and signaling components. Interestingly, when cold-induced  $\text{Ca}^{2+}$  responses were analyzed by specific expression of the reporter protein Aequorin in guard cells, Dodd et al. (2006) observed circadian modulations of the cold-induced  $\text{Ca}^{2+}$  elevation that were significantly more pronounced during the mid-photoperiod than at the beginning or the end of the day. Therefore, it would be interesting to determine if circadian modulation of  $\text{Ca}^{2+}$  responses also occurs in root tissues and during responses to other environmental cues.

Dodd et al. (2006) also performed mathematical modeling analysis of calcium signatures to investigate the relationship between single-cell calcium analyses and whole population (or whole plant) calcium monitoring. This study revealed that the population  $\text{Ca}^{2+}$  signature may be diagnostic for underlying single-cell  $\text{Ca}^{2+}$  oscillations, but it is unlikely to be representative of single-cell  $\text{Ca}^{2+}$  signatures. This is a very important finding that needs to be considered when interpreting work of  $\text{Ca}^{2+}$  dynamics that has used Aequorin as a (whole plant) reporter protein. For example, the second sustained  $\text{Ca}^{2+}$  elevation of biphasic  $\text{Ca}^{2+}$  responses that has been observed in such studies may in fact represent an amalgamation of asynchronous  $\text{Ca}^{2+}$  oscillations that occur in individual cells of the plant after the initial synchronous  $\text{Ca}^{2+}$  transient (Dodd et al., 2006).

Recently, it has been demonstrated that heat shock induces prolonged  $[\text{Ca}^{2+}]_{\text{cyt}}$  elevation ( $\sim 20$  min duration) via putative heat-sensitive plasma membrane  $\text{Ca}^{2+}$ -permeable channels in the moss *Physcomitrella patens* (Saidi et al., 2009). It appears that a larger increase in temperature induces a more intense  $[\text{Ca}^{2+}]_{\text{cyt}}$  elevation and results in a stronger response (e.g., expression of heat shock proteins) compared with a less pronounced temperature elevation shift. In *Arabidopsis*,  $\text{Ca}^{2+}$  and

calmodulin proteins have been shown to be involved in the heat shock response (Zhang et al., 2009). It is likely that flowering plants also sense temperature increments with heat-sensitive channels. It would be interesting to elucidate whether heat shock induces cell type-specific  $[Ca^{2+}]_{cyt}$  elevations in *Arabidopsis* and to compare the  $Ca^{2+}$ -related mechanisms of heat stress responses with that of the cold response.

### Response to Mechanical Stimuli

There is ample evidence that mechanical stimuli, such as touch or wind, induce  $[Ca^{2+}]_{cyt}$  elevations in plant cells (Braam, 2005). Recently, Monshausen et al. (2009) reported that different types of mechanical stimuli, such as touch and bending, induce distinct patterns of  $Ca^{2+}$  responses in the *Arabidopsis* root. Whereas touch stimuli induce monophasic  $[Ca^{2+}]_{cyt}$  elevations in the cells at the touch site, bending elicits biphasic transient elevations in the cells on the convex (stretching) side. These  $Ca^{2+}$  responses are essential for apoplastic alkalization as well as RBOH C-dependent apoplastic ROS production that may contribute to resistance of plants to mechanical stresses (Monshausen et al., 2009). Bending of the root recruits pericycle cells on the convex side to become founder cells of a new lateral root (LR) primordium. Interestingly, blocking the bending-induced  $[Ca^{2+}]_{cyt}$  elevation inhibits the recruitment of new LRs (Richter et al., 2009), indicating an essential role of  $Ca^{2+}$  signals in this process. However, it remains to be elucidated how the biphasic pattern of  $Ca^{2+}$  elevation encrypts specific information for bending-induced LR production.

### GENERATION OF $Ca^{2+}$ SIGNALS

A  $Ca^{2+}$  signal is defined by the balanced activation of  $Ca^{2+}$  channels at different cellular membranes, which is followed by the subsequent inactivation of channels and activation of efflux transporters to terminate  $Ca^{2+}$  influx and to rebalance the cellular  $Ca^{2+}$  homeostasis. Both processes are strictly regulated and define the physiological outcome of  $Ca^{2+}$  signaling.

### Influx of $Ca^{2+}$

Distinct ion channel types capable of mediating  $Ca^{2+}$  fluxes coexist in different cell types and tissues. According to their activation mechanism, these channels can be classified as voltage-dependent, voltage-independent/ligand-dependent, and stretch-activated  $Ca^{2+}$  channels (Cosgrove and Hedrich, 1991; White et al., 2002; White and Broadley, 2003; Dutta and Robinson, 2004; Nakagawa et al., 2007). Depending on their specific activation properties,  $Ca^{2+}$  channels can shape the parameters of  $Ca^{2+}$  influx and the resulting  $Ca^{2+}$  signature (White et al., 2002; Demidchik and Maathuis, 2007). This enables the plant to translate a wide range of different signals into distinct  $Ca^{2+}$  signatures (Miedema et al., 2001, 2008). Moreover, variability in the specific abundance of the different channel types likely reflects the special needs of a cell type or tissue (Demidchik et al., 2002). A schematic summary of the channels and transporters that are discussed in this review is illustrated in Figure 2.


### Voltage-Dependent Channels at the Plasma Membrane

Voltage-dependent  $Ca^{2+}$ -permeable channels have been classified as depolarization-activated  $Ca^{2+}$ -permeable channels (DACCs) and hyperpolarization-activated  $Ca^{2+}$ -permeable channels (HACCs) (White et al., 2002). Although the properties of DACCs and HACCs are well studied electrophysiologically (Grabov and Blatt, 1998; Thion et al., 1998; Thuleau et al., 1998; Hamilton et al., 2000; Pei et al., 2000; Klusener et al., 2002), the molecular identity of these channels is still unknown (White et al., 2002). It is assumed that DACCs contribute to the short transient influx of  $Ca^{2+}$  in response to various stimuli, including chilling and microbe interaction (Thion et al., 1998). HACCs contribute to a sustained  $Ca^{2+}$  influx in response to ABA (Hamilton et al., 2000; Pei et al., 2000), blue light (Harada and Shimazaki, 2009), and  $Ca^{2+}$  nutrition (Miedema et al., 2001, 2008).

Plant annexins are ubiquitous, soluble proteins capable of  $Ca^{2+}$ -dependent and  $Ca^{2+}$ -independent binding to endomembranes and the plasma membrane (Demidchik and Maathuis, 2007; Mortimer et al., 2008). Surprisingly, a recent study indicated that cytosolic annexins create a  $Ca^{2+}$  influx pathway directly, particularly during stress responses involving acidosis in *Zea mays* (Laohavisit et al., 2009). This suggests that annexins modulate or even represent a part of voltage-gated  $Ca^{2+}$ -permeable channels. However, further characterization of plant annexins and especially the molecular identification of voltage-gated  $Ca^{2+}$  channels are required to enable more insights of the contribution of annexins and voltage-gated channels to  $Ca^{2+}$  signaling (Cheong et al., 2007).

### Ligand-Gated Channels at the Plasma Membrane

Cyclic nucleotide-gated channels (CNGCs) are ligand-gated channels that are important for cellular homeostasis of cations and can mediate fluxes of  $Ca^{2+}$  ions (Hua et al., 2003a; Ali et al., 2006). Twenty CNGC genes have been identified in *Arabidopsis* (White et al., 2002). CNGCs are activated by binding of the cyclic nucleotides cAMP and cGMP and harbor a binding site for calmodulin that partially overlaps with the binding domain for cyclic nucleotides (cNMPs). Consequently, binding of  $Ca^{2+}$ /calmodulin results in inactivation of CNGCs due to blocking of the cNMP binding domain (Hua et al., 2003b; Ali et al., 2006). Thereby,  $Ca^{2+}$  itself, by binding to calmodulins/calmodulin-like proteins (CMLs), can modulate the influx of  $Ca^{2+}$  mediated by CNGCs. Several CNGCs, including CNGC4, CNGC11, and CNGC12, have been implicated in response reactions to pathogens (Balague et al., 2003; Yoshioka et al., 2006; Urquhart et al., 2007). CNGC2 represents an extensively characterized channel that was originally identified by cloning of the gene responsible for the *defense no death1 (dnd1)* mutant phenotype. Mutant plants fail to induce the  $Ca^{2+}$ -mediated hypersensitive response to an avirulent strain of the pathogen *Pseudomonas syringae* and exhibit enhanced resistance to pathogens (Yu et al., 1998; Clough et al., 2000). Specifically, *cngc2* mutants are impaired in nitric oxide production that is required for the occurrence of a hypersensitive response (Ali et al., 2007) and that was previously reported to depend specifically on  $Ca^{2+}$  influx from extracellular stores (Lamotte et al., 2004).



**Figure 2.** Overview of Ca<sup>2+</sup> Transport Systems in an *Arabidopsis* Cell.

Shown are Ca<sup>2+</sup> influx/efflux pathways that have been identified at the molecular level. See text for further details. CNGC, cyclic nucleotide channel; GLR, glutamate receptor; TPC1, two pore channel 1; CAS, Ca<sup>2+</sup>-sensing receptor; ACA, autoinhibited calcium ATPase; ECA, ER type calcium ATPase; HMA1, heavy metal ATPase1; CAX, cation exchanger.

The function of CNGCs appears not to be restricted to plant pathogen interactions. CNGC3 and CNGC10 are implicated in regulating ion homeostasis especially for establishing the proper Na<sup>+</sup>/K<sup>+</sup> ratio during salt stress adaptation (Gobert et al., 2006; Guo et al., 2008). CNGC18 is asymmetrically localized to the tip region of growing pollen tubes and has been shown to be indispensable for proper tip growth of pollen, suggesting that this channel is important to establish the tip-focused gradient of Ca<sup>2+</sup> (Frietsch et al., 2007). Consequently, CNGC18 provides a mechanism for direct transduction of a cNMP signal into an ion flux that can produce a localized signal capable of regulating the pollen tip growth machinery. Considering the mechanistic similarities in the signal transduction processes governing tip growth in pollen tubes and root hairs, it would be most interesting to address the function of CNGCs in root hairs. Moreover, the investigation of Ca<sup>2+</sup> dynamics in *cngc* mutant lines that would express the Ca<sup>2+</sup> reporter protein cameleon would be most helpful for further elucidating the contribution of CNGC toward the generation of cellular Ca<sup>2+</sup> signals.

Similar to CNGCs, glutamate receptors (GLRs) are nonselective cation channels that have attracted considerable attention as important regulators of Ca<sup>2+</sup> influx. In *Arabidopsis*, 20 genes encode for GLRs that are differentially activated by Glu and Gly, as well as by other amino acids, and mediate an increase of cytosolic Ca<sup>2+</sup> (Chiu et al., 2002; Qi et al., 2006; Stephens et al., 2008). It is assumed that GLRs are important for plant Ca<sup>2+</sup> nutrition (Kim et al., 2001; Demidchik and Maathuis, 2007) but also in mediating Ca<sup>2+</sup> responses upon cold stress (Meyerhoff

et al., 2005) or excess aluminum (Sivaguru et al., 2003). At GLR1.1 regulates the expression of enzymes involved in carbon and nitrogen metabolism and regulates ABA biosynthesis (Kang and Turano, 2003; Kang et al., 2004). At GLR1.1 antisense lines contain elevated ABA levels and are hypersensitive to ABA or glucose (Kang and Turano, 2003; Kang et al., 2004). Overexpression of another GLR, At GLR3.1, impaired long-term stomatal closure but did not affect the short-term stomatal closure response or the kinetics of Ca<sup>2+</sup> oscillations that were imposed by extracellular Ca<sup>2+</sup> (Cho et al., 2009). In rice (*Oryza sativa*), disruption of the *Os GLR3.1* gene results in reduced growth of the primary root and of adventitious roots especially in the early seedling stage due to reduced mitotic activity of the root apical meristem (Li et al., 2006a). GLRs also have been implicated in root development in *Arabidopsis* since Glu can affect root architecture (Walch-Liu et al., 2006). Additionally, GLRs might have a role in Ca<sup>2+</sup>-dependent generation of electrical potentials in the root (Masi et al., 2009). Moreover, plants that have been treated with antagonists of GLRs display impaired light-induced signal transduction, harbor enlarged hypocotyls, and accumulate less chlorophyll when grown in light, suggesting that GLRs may also function in photomorphogenesis (Lam et al., 1998; Brenner et al., 2000).

These findings suggest that extracellular Glu not only provides a signal for Ca<sup>2+</sup> nutrition but in addition regulates different developmental aspects, electrical transmission, supply of nutrients, and responses to abiotic stress. Therefore, it would be of considerable interest to investigate the functional interrelation of

plant CNGCs and GLRs and to investigate whether they act in concert to mediate  $\text{Ca}^{2+}$  signaling.

### **Voltage-Dependent Channels at Other Cellular Membranes**

The coexistence of voltage-dependent channels and ligand-gated channels appears not to be restricted to the plasma membrane since compelling evidence points to a function of both channel types also in the vacuolar membrane. An important voltage-dependent channel activity was identified as a slow vacuolar type (SV) channel under depolarized conditions (Johannes et al., 1992; Allen and Sanders, 1994). This SV channel originally was described as voltage-dependent channel that is regulated by cytoplasmic  $\text{Ca}^{2+}$  concentrations (Hedrich and Neher, 1987), and subsequent work reported the  $\text{Ca}^{2+}$  release capability of this channel (Allen and Sanders, 1994). In addition, this channel seems to be regulated by  $\text{Ca}^{2+}$  in two opposing ways. While elevation of cytoplasmic  $\text{Ca}^{2+}$  can activate the channel, an increase in vacuolar luminal  $\text{Ca}^{2+}$  inactivates the channel (Pottosin and Schonknecht, 2007). This electrophysiologically characterized channel activity was recently shown to be conferred by the protein Two-pore channel1 (TPC1) (Peiter et al., 2005). Although the SV channel is the most abundant vacuolar channel and *TPC1* represents a single gene in the *Arabidopsis* genome, loss of SV channel function in *Arabidopsis* does not result in a lethal phenotype or severe growth defects. However, loss of *TPC1* function has a significant effect on germination inhibition by ABA, and while there is no effect on guard cell ABA signaling, there is a highly significant effect on stomatal responses to external  $\text{Ca}^{2+}$  (Peiter et al., 2005; Ranf et al., 2008). Although the SV channels appear to make only a minor contribution to a global  $\text{Ca}^{2+}$  release into the cytosol by  $\text{Ca}^{2+}$  influx from the vacuole, it was proposed that TPC1 channels could form clusters within the tonoplast to form local  $\text{Ca}^{2+}$  microdomains (Perez et al., 2008). In contrast with the situation in *Arabidopsis*, knockout mutation of *TPC1* in rice results in reduced plant growth on soil (Kadota et al., 2004). Moreover, overexpression of Os *TPC1* resulted in prolonged activation of the mitogen-activated protein kinase (MAPK) pathway, indicating that in rice (Kurusu et al., 2005) and also in tobacco (*Nicotiana tabacum*; Kadota et al., 2004), channel function could have a role in pathogen response. Similarly, knockout of *TPC1* in *Arabidopsis* results in weaker induction of defense genes after pathogen infection but does not significantly affect pathogen resistance (Bonaventure et al., 2007b). By contrast, a hyperactive gain-of-function allele of *TPC1*, designated as *fou2*, leads to enhanced defense gene expression and resistance to pathogens (Bonaventure et al., 2007a, 2007b). Although the exact physiological role of TPC1 in *Arabidopsis* is not fully understood, the fact that TPC1 mediates the flux of  $\text{K}^+$  and other cations through the vacuolar membrane (Bihler et al., 2005; Ivashikina and Hedrich, 2005) may point to an important role of TPC1 in regulating cytosolic  $\text{K}^+$  homeostasis in a  $\text{Ca}^{2+}$ -dependent manner (Pottosin and Schonknecht, 2007). In agreement with such a function, SV channel activity is down-regulated at low vacuolar potassium concentrations (Pottosin et al., 2005), and the *fou2* transcriptome resembles the transcript profile of plants after potassium starvation (Bonaventure et al., 2007b). Moreover, *fou2* mutant plants contain less  $\text{K}^+$  and more

$\text{Ca}^{2+}$  within the vacuole (Beyhl et al., 2009). Several recent investigations reported that NAADP binds to and confers the activation of animal TPC channels that results in  $\text{Ca}^{2+}$  release from lysosomal and endosomal compartments (Brailoiu et al., 2009; Calcraft et al., 2009). Since these animal TPC channels are structurally closely related to plant TPC1 but lack the EF hand motif in their central domain (Ishibashi et al., 2000), it will be most interesting to investigate if the respective plant channels are also responsive to this second messenger.

In addition to TPC1, several other less well characterized voltage-dependent channels appear to exist at the vacuolar membrane (Allen and Sanders, 1994), which were characterized as a fast vacuolar channel (Hedrich and Neher, 1987) and as a  $\text{Ca}^{2+}$ -insensitive vacuolar channel (Ranf et al., 2008).

### **Ligand-Gated Channels at the Vacuolar and Endoplasmic Reticulum Membranes**

Several electrophysiological studies supported the existence of ligand-gated channels at the tonoplast and suggested regulation of these channels by IP3/IP6 and cADPR (Schumaker and Sze, 1987; Allen et al., 1995; Muir and Sanders, 1996; Martinec et al., 2000; Navazio et al., 2000; Lemtiri-Chlieh et al., 2003). However, despite the existence of multiple full-genome sequences for several higher plant species, no genes encoding for an ADP ribosyl cyclase that would be required for cADPR production have been identified. In addition, ryanodine receptors that are the targets of cADPR in animal cells appear to be missing in higher plants. In this regard, experimental determination and direct proof of cADPR abundance by gas chromatography–mass spectrometry techniques are urgently needed.

Similarly, no sequences with significant similarity to animal IP3 receptors have been identified in genomes of higher plants. Interestingly, while higher plants lack classical endoplasmic reticulum (ER)-localized IP3 receptors, which are well known and characterized in animals (Berridge, 2009), several algae species, such as *Volvox* and *Chlamydomonas*, appear to harbor these receptor channels, suggesting that this channel type was present in the ancient plant progenitor of the evolutionary plant lineage and may have been lost during the further evolution of the plant lineage (Wheeler and Brownlee, 2008). This raises the obvious question of how IP3 can function in plant  $\text{Ca}^{2+}$  signaling in the absence of canonical IP3 receptors. The absence of IP3 receptors in higher plants in combination with the extremely low levels of phosphatidylinositol-4,5-bisphosphate in plant tissues, that in animal cells is converted by PLC into IP3, has raised concerns about the simple assignability of mechanistic principles of IP3-induced  $\text{Ca}^{2+}$  release from animal systems to the situation in plant cells (Munnik and Testerink, 2009). Therefore, future research in this field of plant  $\text{Ca}^{2+}$  signaling may unravel unexpected mechanisms that interconnect  $\text{Ca}^{2+}$  and IP3.

In addition to the previously mentioned ligand gated channels, a unique ligand-gated channel that appears to reside in the ER membrane is activated by NAADP (Navazio et al., 2000). However, since the molecular identity of all these channels is still unknown, it remains to be established if these ligands directly activate channels or, alternatively, function via receptors that indirectly modulate the activity of channels.

### Efflux of Ca<sup>2+</sup>

Much of the research on Ca<sup>2+</sup> signaling has focused on advancing our understanding of the generation of Ca<sup>2+</sup> elevations by Ca<sup>2+</sup>-releasing channels. However, to represent a distinct signal, the regulation of Ca<sup>2+</sup> efflux that not only terminates but also shapes the Ca<sup>2+</sup> signature is as important as the Ca<sup>2+</sup>-releasing events. During the past years, we have witnessed significant insights into the regulation of Ca<sup>2+</sup> extrusion. However, in the future, it will be most important to reveal the interconnected regulation of Ca<sup>2+</sup> release and extrusion that is required to generate defined signals.

Extrusion of Ca<sup>2+</sup> from the cytosol is achieved by P-type Ca<sup>2+</sup>-ATPases and by the Ca<sup>2+</sup>/proton antiporter systems. While antiporters mediate a high-affinity low turnover efflux, ATPases mediate a low-affinity high-capacity efflux. Therefore, it is assumed that antiporters reduce the Ca<sup>2+</sup> concentration back to a few micromolar after signal mediated influx, while Ca<sup>2+</sup>-ATPases are important to maintain the low resting concentration of Ca<sup>2+</sup> (Hirschi, 1999).

The coordinative regulation of the cellular extrusion systems still is not fully understood. Transport activity is clearly activated after influx of Ca<sup>2+</sup>, as in response to different hormones (Erdei et al., 1979; Bush et al., 1993; Zocchi and Rabotti, 1993), salt stress (Gao et al., 2004), or mechanical stimulation (Bourgeade et al., 1991). Enhanced Ca<sup>2+</sup> extrusion was reported to occur in senescent potato (*Solanum tuberosum*; Fakhrai and Hall, 1984). Moreover, enhanced transcription of ATPases was observed in response to ABA (Cerana et al., 2006), sugar (Mito et al., 1996), or by sodium (Wimmers et al., 1992).

Hormones can also differentially activate transporter systems of different cellular membranes. Ca<sup>2+</sup> released by GA seems to be transported mainly via ER transporters out of the cytoplasm, while ABA activates transport activity at the ER as well as at the tonoplast (Bush and Sze, 1986; Bush et al., 1989a, 1993). Interestingly, overall, the Ca<sup>2+</sup> extrusion system seems to be less effective in protoplasts or cells from epidermal strips than in intact leaves (Levchenko et al., 2008). This important observation should be considered in the design of future experiments addressing the regulation of Ca<sup>2+</sup> extrusion.

### Ca<sup>2+</sup>-Proton Antiporter

In the *Arabidopsis* genome, six genes encode for putative Ca<sup>2+</sup>-proton antiporters, also named cation exchangers (CAX) (Maser et al., 2001; Shigaki et al., 2006), which regulate the homeostasis of Ca<sup>2+</sup> and other divalent cations, such as cadmium (Catala et al., 2003; Cheng et al., 2003; Koren'kov et al., 2007; Zhao et al., 2008). Five additional antiporters that were previously designated as CAX7-11 resemble potassium (Na<sup>+</sup>)-dependent Na<sup>+</sup>/Ca<sup>2+</sup> antiporters and were therefore renamed cation Ca<sup>2+</sup> exchanger proteins (Shigaki et al., 2006). Additionally, four putative antiporters encoded in the *Arabidopsis* genome contain EF hand Ca<sup>2+</sup> binding motifs, implicating that these transporters are directly regulated by Ca<sup>2+</sup> (Shigaki et al., 2006). It is unknown to which extent the latter two types of antiporters are important for Ca<sup>2+</sup> extrusion.

The antiporters CAX1 to CAX4 are localized to the vacuole (Hirschi et al., 2000; Cheng et al., 2002, 2003, 2005),

but antiporter activity was also reported to reside at the plasma membrane (Kasai and Muto, 1990; Luo et al., 2005). CAX proteins harbor an N-terminal regulatory/autoinhibitory domain, which binds to an adjacent region within the N terminus (Pittman et al., 2002; Mei et al., 2007). The exact mechanistic regulation by the N terminus is not well understood, and differential N-terminal-dependent regulation of CAX protein activity may occur. Autoinhibition can be relieved by formation of heteromers of, for example, CAX1 and CAX3 (Zhao et al., 2009), and different regulatory proteins could additionally interact with CAX proteins to regulate transport activity (Cheng and Hirschi, 2003; Cheng et al., 2004a, 2004b). The importance of specific transport activity regulation by the N-terminal region was demonstrated by overexpressing an N-terminal truncated, deregulated version of the vacuolar Ca<sup>2+</sup>/proton antiporter CAX1 from *Arabidopsis* in tobacco. Although tobacco plants contained more total Ca<sup>2+</sup>, these plants showed Ca<sup>2+</sup> deficiency symptoms and displayed hypersensitivity to magnesium, sodium, and cold shock (Hirschi, 1999). It was assumed that overexpression of At CAX1 leads to overloading of Ca<sup>2+</sup> into the vacuole and to a severe reduction of cytosolic Ca<sup>2+</sup> concentration, which caused the observed deficiency symptoms.

The formation of functional heteromers does not exclude the possibility that single CAX proteins are important for responses to specific signals. While both *cax1* and *cax3* mutant plants are hypersensitive to ABA during germination (Zhao et al., 2008), *cax3* mutants exhibited increased sensitivity toward NaCl or LiCl and low pH levels, while *cax1* mutants displayed normal wild-type responses (Catala et al., 2003; Zhao et al., 2008). This difference can be partially explained by the observation that CAX3 is prominently expressed in roots, whereas CAX1 is predominantly expressed in leaves (Zhao et al., 2008).

### P-ATPases

Classical Ca<sup>2+</sup> P-ATPases belong to the second subclass (II) of phosphorylated (P)-type ATPases. They are classified as P<sub>IIA</sub>- or ER-type Ca<sup>2+</sup> ATPases (ECAs; which include four members in *Arabidopsis*) and P<sub>IIIB</sub> ATPases (10 family members in *Arabidopsis*, all of which contain an autoinhibitory N-terminal region; Sze et al., 2000). Therefore, the latter ATPases also have been named autoinhibited Ca<sup>2+</sup> ATPases (ACAs) (Sze et al., 2000). The autoinhibitory domain in P<sub>IIIB</sub>-type proteins can be relieved by binding of calmodulin to the regulatory domain, which results in activation of the pump (Harper et al., 1998). On the other hand, the activity of the P<sub>IIIB</sub>-type Ca<sup>2+</sup> ATPase ACA2 can be inhibited by phosphorylation within the N-terminal regulatory domain by other Ca<sup>2+</sup>-regulated proteins, such as CDPKs (Hwang et al., 2000).

P<sub>IIA</sub>-type ATPases are localized at the ER (ECA1) (Liang et al., 1997), the Golgi (ECA3) (Mills et al., 2008), and endosomes (also ECA 3) (Li et al., 2008), implicating that the latter organelles can function as mobile Ca<sup>2+</sup> stores that could contribute to the spatial specificity of Ca<sup>2+</sup> signaling (Menteyne et al., 2006).

P<sub>IIIB</sub> types are localized at the ER (ACA2) (Harper et al., 1998), vacuole (ACA4 and ACA11) (Geisler et al., 2000; Lee et al., 2007b), plasma membrane (ACA8, ACA9, and ACA10) (Bonza et al., 2000; Schiott et al., 2004; George et al., 2008), and at the plastid envelope (ACA1) (Huang et al., 1993). The importance of a

$P_{11a}$ -type  $Ca^{2+}$ -ATPase activity regulating the cytoplasmic  $Ca^{2+}$  dynamics was recently impressively exemplified by the analysis of a  $Ca^{2+}$ -ATPase loss-of-function mutant in the moss *P. patens* (Qudeimat et al., 2008). Whereas wild-type plants exhibit a transient cytosolic  $Ca^{2+}$  signature after applying sodium stress, loss-of-function mutant lines displayed a sustained elevation of  $Ca^{2+}$  (Qudeimat et al., 2008). Interestingly, the sustained level of  $Ca^{2+}$  leads to a reduced upregulation of salt stress-induced genes and renders mutant plants specifically less tolerant to sodium stress (Qudeimat et al., 2008). These findings not only establish the importance of regulated  $Ca^{2+}$  extrusion for appropriate formation of  $Ca^{2+}$  signatures, they also implicate a direct interaction between the shape of a  $Ca^{2+}$  signature and proper stress responsiveness. In *Arabidopsis*, analysis of loss-of-function mutants of ACA9 and ACA10 indicated that the pumps specifically function in pollen tube growth and in inflorescence development of plants, respectively (Schiott et al., 2004; George et al., 2008).

Besides ECAs and ACAs,  $P_1$ -type ATPases, which are known as heavy metal transporters, were recently implicated in  $Ca^{2+}$  transport. At HMA1, a heavy metal transporter involved in detoxification processes for heavy metals, is a  $P_1$ -ATPase that localizes to the chloroplast envelope. At HMA1 transports  $Ca^{2+}$  and heavy metals, such as copper, with high affinity and, similar to  $Ca^{2+}$  pumps from animals, is specifically inhibited by thapsigargin (Seigneurin-Berny et al., 2006; Moreno et al., 2008).

All of these observations support the notion that finely regulated  $Ca^{2+}$  extrusion is as important as the influx of  $Ca^{2+}$ , as the deregulated hyperextrusion of  $Ca^{2+}$  as well as downregulated extrusion result in deregulated  $Ca^{2+}$  dynamics and homeostasis. Certainly, a minimal concentration of the toxic cation  $Ca^{2+}$  is required to sustain correct signaling and metabolic functions, while decelerated extrusion of  $Ca^{2+}$  disturbs the formation of defined  $Ca^{2+}$  signatures and impairs the capabilities of the plants to correctly respond to a signal.

### Calcium Signal Modulation by Organelles

Plastids can accumulate high levels of  $Ca^{2+}$  (Nobel et al., 1966) in the millimolar range and thereby can contribute to the homeostasis of cellular  $Ca^{2+}$  and other ions (Portis and Heldt, 1976).  $Ca^{2+}$  within plastids, especially the correct distribution between stroma and thylakoid lumen, is important for the regulation of plastidial enzymes (Brand and Becker, 1984; Kreimer et al., 1988). The level of  $Ca^{2+}$  within plastids is regulated and can raise upon illumination (Muto et al., 1982; Kreimer et al., 1988) or during transition to dark (Sai and Johnson, 2002).

A protein identified as the  $Ca^{2+}$  binding protein CAS may influence the loading capacity of chloroplasts or may be important for sensing the loading status of  $Ca^{2+}$  within the chloroplasts, thereby modulating cytoplasmic  $Ca^{2+}$  signaling (Figure 2) (Han et al., 2003; Nomura et al., 2008; Weini et al., 2008). CAS was first reported as an extracellular  $Ca^{2+}$ -sensing receptor, exhibiting a high capacity to bind  $Ca^{2+}$  (10 to 12  $Ca^{2+}$  ions per molecule) (Han et al., 2003). However, the protein contains an N-terminal chloroplast targeting signal peptide, and subsequent reports identified CAS as a chloroplast-localized protein regulating cytoplasmic  $Ca^{2+}$  levels (Nomura et al., 2008; Vainonen et al., 2008;

Weini et al., 2008). Within the chloroplast, CAS is targeted to the thylakoid membrane (Vainonen et al., 2008; Weini et al., 2008). CAS can be phosphorylated in a light-dependent manner depending on the activity of the light-regulated kinase STN8 (Vainonen et al., 2005, 2008). Although the activity of the photosystem is unaltered, *cas* knockout plants show retarded growth. Especially when grown under low  $Ca^{2+}$  conditions, plants show delayed bolting and impaired induction of flowering (Han et al., 2003). Mutant plants also show a strong deficit in regulating cytoplasmic  $Ca^{2+}$  levels (Vainonen et al., 2008) and are not able to induce stomatal closure provoked by extracellular  $Ca^{2+}$  (Han et al., 2003; Nomura et al., 2008; Weini et al., 2008). However, *cas* knockout plants can respond to externally imposed  $Ca^{2+}$  oscillations and then display normal stomatal closure compared with the wild type, indicating a defect in the generation of  $Ca^{2+}$  transients that are required for stomatal closure, rather than the response machinery per se (Weini et al., 2008). This indicates that the chloroplast-targeted  $Ca^{2+}$  sensor protein somehow connects cytoplasmic and chloroplast  $Ca^{2+}$  levels. This function resembles that of the  $Ca^{2+}$  buffer protein CRT within the ER lumen (Persson et al., 2001; Jia et al., 2009). Similarly, loss of CAS may lead to a reduced buffer capacity of the chloroplasts, suggesting that less  $Ca^{2+}$  can be allocated from the chloroplasts to the transient cytoplasmic increase of  $Ca^{2+}$ . Therefore, it will be of interest to analyze the concentration and dynamics of  $Ca^{2+}$  in chloroplasts of *cas* mutant lines.

### DECODING AND RELAY OF $Ca^{2+}$ SIGNALS

$Ca^{2+}$  signals evoked by a specific stimulus are presented as defined  $Ca^{2+}$  signatures on the cellular level (as for example in guard cells) as well as in distinct cell types and tissue regions (for example, in roots after mechanical stimulation). It is apparent that the ability of a given cell or tissue to translate these  $Ca^{2+}$  signals into defined molecular and biochemical responses primarily depends on the presence, concentration, cellular localization, and  $Ca^{2+}$  binding affinity of signaling components that can sense such  $Ca^{2+}$  signatures and convey specific output reactions for further information processing (McAinsh and Hetherington, 1998). In plants, a diverse and extensive set of  $Ca^{2+}$  binding proteins that function as cellular  $Ca^{2+}$  sensors represent these first information translation points (Luan et al., 2002; Batistič and Kudla, 2004; McCormack et al., 2005; Kim et al., 2009).

Plant  $Ca^{2+}$  sensor proteins have been classified conceptually into sensor relays and sensor responders (Sanders et al., 2002). Sensor responder proteins, for example, CDPKs, combine within one protein a sensing function ( $Ca^{2+}$  binding and  $Ca^{2+}$ -induced conformational changes) with a response activity (e.g., protein kinase activity). By contrast, sensor relay proteins, like calmodulin, also effectively bind  $Ca^{2+}$  ions and usually undergo  $Ca^{2+}$ -induced conformational changes but lack other effector domains. To transmit the  $Ca^{2+}$  signal, sensor relay proteins interact with target proteins and regulate their activity (Luan et al., 2002; Sanders et al., 2002).

Initially, this concept of functional classification for  $Ca^{2+}$  sensing proteins was applied to the conversion of  $Ca^{2+}$  signals into phosphorylation responses (Sanders et al., 2002). However,

considering new insights into the function of  $\text{Ca}^{2+}$ -sensing proteins, we propose extending this concept to the conversion of  $\text{Ca}^{2+}$  signals into transcriptional responses. Whereas  $\text{Ca}^{2+}$  binding to calmodulin 7 (CAM7) appears to result in direct promoter interaction and regulation, other calmodulins are likely to mediate gene regulation by interacting with calmodulin binding transcription factors (CAMTAs) that function as transcriptional coregulators (see below). Consequently, CAM7 would represent a bona fide sensor responder directly contributing to activating gene expression, while other calmodulins function as sensor relays modulating gene expression via their interaction with proteins that function as transcription factors (Kushwaha et al., 2008).

In addition to calmodulins, CMLs, which are represented by >50 diverse calcium sensor proteins in *Arabidopsis*, appear to fulfill important functions in plant development and responses to environmental cues (McCormack et al., 2005). Loss of CML42 function leads to aberrant trichomes with increased branching, while mutation of CML24 causes alterations in flowering time (Delk et al., 2005; Tsai et al., 2007; Dobney et al., 2009). Moreover, CML24 functions in responses to ABA and ionic stress, and mutation of CML9 alters plant responses to ABA and abiotic stresses (Delk et al., 2005; Magnan et al., 2008). Exploring the function of the other members of this diverse protein family and especially identifying targets of these proteins therefore represents a promising area of plant signaling research.

Many biological processes, such as metabolic starch degradation by  $\alpha$ -amylase (Bush et al., 1989b); biosynthetic processes, such as brassinosteroid synthesis (Du and Poovaiah, 2005); or even the mechanical occlusion of sieve tube elements (Furch et al., 2009) are likely important targets of direct  $\text{Ca}^{2+}$ -dependent modulation. However, with respect to the signaling function of  $\text{Ca}^{2+}$  ions,  $\text{Ca}^{2+}$ -dependent phosphorylation events and  $\text{Ca}^{2+}$ -dependent gene regulation represent the major cellular currencies for converting defined  $\text{Ca}^{2+}$  signatures into specific downstream reactions. Recent advances in these two facets of  $\text{Ca}^{2+}$  signaling will be discussed in detail here.

### Translating $\text{Ca}^{2+}$ Signatures into Protein Phosphorylation

Signaling requires messengers whose concentration varies in time and space.  $\text{Ca}^{2+}$  ions and phosphate ions have come to dominate cellular signaling.  $\text{Ca}^{2+}$  binding triggers changes in protein shape and charge. Similarly, phosphorylation imparts a negative charge, modulating protein conformations and protein interactions (Hunter, 1995; Soderling, 1999; Clapham, 2007).  $\text{Ca}^{2+}$ -dependent kinases and protein kinases regulated by interaction with  $\text{Ca}^{2+}$  binding proteins functionally combine these two major cellular currencies of signal transduction and allow for the perception and transmission of  $\text{Ca}^{2+}$  signatures directly into phosphorylation events that orchestrate downstream signaling responses (Harmon et al., 2000; Batistič and Kudla, 2004). Plants possess an extensive repertoire of calcium-regulated protein kinases, which are represented by the three families of CCaMKs, CDPKs, and CBL-interacting protein kinases (CIPKs) (Sanders et al., 2002; Batistič and Kudla, 2004; Gleason et al., 2006). While CDPKs and CCaMKs (the later appear not to exist in the *Arabidopsis* genome) represent typical sensor responders, the CIPKs are targets of Calcineurin B-like (CBL) sensor relay pro-


teins. These kinases and  $\text{Ca}^{2+}$  binding proteins form an intricate cellular network for decoding  $\text{Ca}^{2+}$  signals in diverse cellular processes (Batistič and Kudla, 2009; Weinl and Kudla, 2009).

### The CDPK Signaling System

The *Arabidopsis* genome encodes 34 CDPKs and eight additional CDPK-related kinases (Hrabak et al., 2003). The biochemistry and regulation of CDPKs have been comprehensively reviewed (Harper et al., 2004; Ludwig et al., 2004; Harper and Harmon, 2005). Binding of  $\text{Ca}^{2+}$  to the C-terminal EF hand-containing regulatory domain leads to conformational changes relieving the active site of the kinase domain from masking by an autoinhibitory domain, resulting in activation of the respective CDPK. This process is enhanced by autophosphorylation of the CDPKs that contribute to full activation of the kinases (Ludwig et al., 2004).

The first conclusive evidence that linked the activation of CDPKs with the induction of signaling responses *in vivo* was performed on the Cf9-avr9 disease response of tomato (*Solanum lycopersicum*) and was obtained by suppression of Nt CDPK2 by viral-induced gene silencing in *Nicotiana benthamiana* (Romeis et al., 2001). CDPK-silenced plants displayed a reduced and delayed hypersensitive response after race-specific Avr9 elicitation in a gene-for-gene interaction and lacked an accompanying wilting phenotype. Subsequent further analysis of Nt CDPK2 function by T. Romeis and coworkers additionally indicated that elevated CDPK signaling compromises stress-induced MAPK activation and that this inhibition requires ethylene synthesis and perception (Ludwig et al., 2005). These important findings suggest that CDPK and MAPK pathways do not function independently and that a concerted regulation of both pathways controls response specificity to biotic and abiotic stress. Clearly, such aspects of interconnection and interplay between different plant signaling systems will need to be investigated in more depth in the future.

The application of reverse genetics techniques has recently extended our knowledge of the physiological function of several members of the CDPK protein family (see Figure 3 for an illustrated summary). *Arabidopsis* CPK3 and CPK6 function synergistically in the regulation of stomatal closure in response to ABA and external  $\text{Ca}^{2+}$  elevation (Mori et al., 2006). Moreover, in independent alleles of single and double mutants of *cpk3* and *cpk6*, ABA and  $\text{Ca}^{2+}$  activation of slow-type anion channels and ABA activation of plasma membrane  $\text{Ca}^{2+}$ -permeable channels were impaired in guard cells (Mori et al., 2006). In addition to CPK3 and CPK6, *Arabidopsis* CPK4 and CPK11 are critical for proper ABA responsiveness of guard cells and have been shown to phosphorylate the ABA-responsive transcription factors ABF1 and ABF4 *in vitro* (Zhu et al., 2007). Research in tobacco revealed that CDPK1 regulates the transcription factor RSG in response to gibberellins (Ishida et al., 2008). Phosphorylation of Ser-114 in RSG by CDPK1 promotes interaction with 14-3-3 proteins that appears to regulate the intracellular localization of this transcription factor. Work in potato suggests that two CDPKs (St CDPK4 and St CDPK5) phosphorylate NADPH oxidases and thereby positively regulate the production of ROS (Kobayashi et al., 2007). Accordingly, ectopic expression of a constitutively active



**Figure 3.** The CDPK Signaling System for Translating  $\text{Ca}^{2+}$  Signatures into Protein Phosphorylation.

Shown are characterized components of the CDPK system. See text for further details. RSG, repression of shoot growth; ABF, ABA response element-binding factor.

mutant of St CDPK5 provoked ROS production in *N. benthamiana* leaves, and this CDPK-mediated ROS production was disrupted by knockdown of St CDPK5. Together, these findings point to a critical role of CDPK-mediated  $\text{Ca}^{2+}$  signaling in a diverse set of biological processes. The identification of additional CDPK target proteins and CDPK-regulated process should in the near future allow us to address more precisely how specific CDPKs mechanistically contribute to a specific decoding of distinct  $\text{Ca}^{2+}$  signatures.

### The CBL/CIPK Signaling System

CBL proteins and their interacting protein kinases (CIPKs) were originally identified in *Arabidopsis* (Kudla et al., 1999; Shi et al., 1999). Subsequent comprehensive bioinformatics analyses have identified a complement of 10 CBLs and 26 CIPKs in *Arabidopsis* and 10 CBLs and 30 CIPKs in the genome of rice, respectively (Kolukisaoglu et al., 2004; Weinel and Kudla, 2009). In contrast with the CDPK sensor responders, the families of CBL proteins and their interacting CIPKs separate  $\text{Ca}^{2+}$  binding function (sensor relay function) and kinase activity (response activity) into two flexible combinable modules. In addition, preferential complex formation of individual CBLs with defined subsets of CIPKs appears to be one of the mechanisms generating the temporal and spatial specificity of  $\text{Ca}^{2+}$  signals in plant cells (Albrecht et al., 2001). Together, these features allow for the formation of a complex and dynamic  $\text{Ca}^{2+}$ -decoding signaling network. Since its initial discovery, this signaling system has been subject to intensive research. Advances in our understanding of the physiological function and structural features of these proteins and functional principles of this  $\text{Ca}^{2+}$ -decoding system have been discussed in several reviews (Luan et al., 2002; Batistič and Kudla, 2004, 2009; Luan, 2009; Luan et al., 2009; Weinel and Kudla, 2009).

CBL proteins exhibit significant similarity to the regulatory B subunit of calcineurin and Neuronal  $\text{Ca}^{2+}$  Sensor proteins from animals and yeast (Kudla et al., 1999). All CBL proteins share a rather conserved core region consisting of four EF hand  $\text{Ca}^{2+}$  binding sites that are arranged in completely invariant spacing within the protein (Kolukisaoglu et al., 2004). All CIPK-type kinases are composed of a conserved N-terminal kinase domain and a C-terminal regulatory domain, which are separated by a variable junction domain. Within the rather divergent regulatory domain, the conserved NAF domain has been identified as required and sufficient for mediating CBL interaction (Albrecht et al., 2001). It is assumed that binding of CBL proteins to the NAF domain of CIPKs releases the C-terminal (autoinhibitory) domain from the kinase domain, thereby transforming the kinase into an active state (Guo et al., 2001; Gong et al., 2002). Additional phosphorylation of the activation loop within the kinase domain by an unidentified kinase further contributes to the activation of CIPKs (Gong et al., 2002). An interesting new facet of CIPK–CBL interaction is provided by recent reports of phosphorylation of CBL proteins by their interacting CIPKs that appears to enhance the CBL–CIPK interaction (Mahajan et al., 2006; Lin et al., 2009). Lin et al. (2009) reported that the kinase CIPK24/SOS2 specifically phosphorylates CBL10 (which these authors renamed S $\text{CaBP8}$ ), but not any other investigated CBL protein, at position Ser-237, a finding that is rather surprising considering that this amino acid position and the surrounding sequence motif is conserved in eight out of the 10 CBL proteins in *Arabidopsis* (Lin et al., 2009). In addition, a protein-phosphatase interaction (PPI) domain mediating CIPK interaction with phosphatases of the PP2C group has been identified within the C terminus of these kinases (Ohta et al., 2003). Unfortunately, the functional implications of this interaction are currently unknown. It appears conceivable that CIPKs may phosphorylate PP2Cs or that

PP2Cs dephosphorylate CIPKs in vivo, as it has been shown for sucrose nonfermenting-related kinases (SnRKs) of the SnRK2 family that are dephosphorylated by a subgroup of PP2Cs (Umezawa et al., 2009). Alternatively, CIPK/PP2C complexes could serve as signaling kinase/phosphatase modules allowing for rapid alternating phosphorylation/dephosphorylation of target proteins. However, crystallization studies of CBL4 (SOS3) in complex with the regulatory domain of CIPK24 (SOS2) suggest that either CBLs or PP2Cs may interact in a mutually exclusive manner with the regulatory domain of CIPKs, thereby preventing the formation of trimeric complexes (Sanchez-Barrera et al., 2007). Consequently, PP2C interaction with the PPI domain of CIPKs would lead to replacement of the CBL protein, which binds to the NAF and partly to the PPI domain, and would allow for competitive formation of either CBL/CIPK or CIPK/PP2C complexes. Considering the recent identification of PP2C phosphatases as direct targets of the PYR/RCAR ABA receptors (Ma et al., 2009; Park et al., 2009), it is tempting to speculate that PP2C-bound CIPKs contribute to early signaling steps after ABA perception, an assumption that would be in agreement with the observation of ABA response phenotypes in many analyzed CIPK mutants (see below).

### **Evolution and Functional Diversification of the CBL/CIPK Signaling System**

The increasing number of available full-genome sequences has facilitated the study of the evolution of the CBL/CIPK signaling system. Single CBL and CIPK genes have been identified in green alga species, such as *Ostreococcus tauri* and *Chlorella* sp, whereas the moss *P. patens* contains four CBLs and seven CIPKs, and the genome of the fern *Selaginella moellendorffii* possesses a complement of five CBL and five CIPK genes (Batistič and Kudla, 2009; Weinl and Kudla, 2009; Batistič et al., 2010). These observations suggest that the complexity of the CBL/CIPK system evolved concurrently with the increasing morphological and developmental sophistication of land plants that enabled the colonization of ecologically diverse and environmentally fluctuating habitats. Surprisingly, CBLs and CIPKs also were recently identified in protozoan eukaryotic species, such as *Trichomonas vaginalis* and *Naegleria gruberi*, raising the question about the function of this Ca<sup>2+</sup> decoding components in nonplant species (Batistič and Kudla, 2009). Remarkably, the occurrence and function of plant-specific Ca<sup>2+</sup> signaling components in human pathogens appears not to be restricted to CBLs and CIPKs, as CDPKs have been identified in *Plasmodium falciparum* and *Plasmodium berghei*, where Pb CDPK4 fulfills a critical function during the life cycle of this human pathogen (Billker et al., 2004; Harper and Harmon, 2005). Considering the absence of related proteins in their human host, further advancements of our understanding of the regulation of plant Ca<sup>2+</sup> sensor proteins may facilitate the identification of therapeutic inhibitors specifically affecting the plant-like proteins of these protozoan species.

Local Ca<sup>2+</sup> signals at specific microdomains are assumed to be the basis for differential Ca<sup>2+</sup> signaling (Berridge, 2006). Consequently, Ca<sup>2+</sup>-decoding systems would be expected to reflect this spatial specification of Ca<sup>2+</sup> signaling. The CBL/CIPK

network appears to meet this demand. Localization studies of all 10 *Arabidopsis* CBL proteins revealed the importance of their variable N-terminal extensions for specific subcellular targeting (Batistič et al., 2010). Four CBL proteins are localized to the plasma membrane, while another four CBLs are localized to vacuolar membrane and two CBLs are detected in the cytoplasm and nucleus (Batistič et al., 2010). These distinct subcellular localizations suggest that CBL Ca<sup>2+</sup> sensors might function as fast relays of local Ca<sup>2+</sup> release events from internal and external stores and that the spatial separation of distinct CBL/CIPK complexes contributes to spatial specificity in Ca<sup>2+</sup> signaling. Dual lipid modification by myristoylation and S-acylation are required for CBL1 function and for localization of this Ca<sup>2+</sup> sensor at the plasma membrane. This localization is achieved by a two-step targeting process in which initial myristoylation results in localization at the ER, and subsequent S-acylation is crucial for ER-to-plasma membrane trafficking (Batistič et al., 2008). As CBL4, CBL5, and CBL9 have been shown to undergo myristoylation and share the adjacent palmitoylation motif, a similar targeting mechanism can be predicted for these proteins (Batistič et al., 2008).

In contrast with CBL proteins, most CIPKs when expressed as green fluorescent protein fusions exhibit a cytoplasmic and nucleoplasmic localization (D'Angelo et al., 2006; Batistič et al., 2010). However, CBL-CIPK interaction analyses using bimolecular fluorescence complementation revealed that the subcellular localization of CIPKs incorporated into distinct CBL/CIPK complexes is determined by the identity of their CBL moiety (D'Angelo et al., 2006; Cheong et al., 2007; Batistič et al., 2008, 2010; Waadt et al., 2008). For example, CIPK1 is targeted to the plasma membrane by CBL1 or CBL9 (Cheong et al., 2007; Waadt et al., 2008). However, upon interaction with CBL2, the resulting CBL2/CIPK1 complexes are localized exclusively to the tonoplast (Batistič et al., 2008). Similarly, CIPK14/CBL2 complexes have been detected at the tonoplast, while the same kinase is targeted to the plasma membrane upon interaction with CBL8 (Batistič et al., 2010). Remarkably, the cellular targeting of plasma membrane- or tonoplast-localized CBL/CIPK complexes appears not to involve the conventional protein trafficking pathway because inhibition experiments using a dominant-negative form of the SAR1 protein (that interferes with COPII vesicle formation) or brefeldin A (that impedes COPI vesicle formation) did not affect the cellular targeting of singular CBL proteins or CBL/CIPK complexes (Batistič et al., 2008, 2010).

### **Physiological Functions of CBLs and CIPKs**

Forward genetic screens aiming to identify critical components of plant salt tolerance have provided insights into the physiological function of CBLs and CIPKs. The CBL Ca<sup>2+</sup> sensor SOS3 (At CBL4) and the CIPK-type kinase SOS2 (At CIPK24) appear to be part of a Ca<sup>2+</sup>-regulated signaling pathway that specifically mediates salt stress adaptation by regulating the Na<sup>+</sup>/H<sup>+</sup> antiporter SOS1 (Liu and Zhu, 1998; Halfter et al., 2000; Qiu et al., 2002). Recent studies revealed that mutation of CBL10 also renders plants salt sensitive and that CBL10 is also able to interact with the kinase CIPK24 (Kim et al., 2007; Quan et al., 2007). In vivo analyses revealed a tonoplast localization of the


CBL10/CIPK24 complex, thereby supporting a functional model wherein alternative complex formation of CIPK24 kinases with either CBL4 or CBL10 creates a dual functioning kinase. While CBL4/CIPK24 complexes mediate  $\text{Na}^+$  extrusion via the regulation of the  $\text{H}^+/\text{Na}^+$  antiporter SOS1 at the plasma membrane, formation of CBL10/CIPK24 likely results in  $\text{Na}^+$  sequestration into the vacuole by regulating unknown targets (Kim et al., 2007; Weint and Kudla, 2009). In addition, recent studies have suggested that CBL/CIPK24 complexes at the tonoplast membrane activate and regulate the vacuolar  $\text{Ca}^{2+}/\text{H}^+$  antiporter CAX1 independently from CBL4 (Cheng et al., 2004b) and have identified subunits of the V-ATPase complex as interacting proteins of CIPK24 (Batelli et al., 2007).

Reverse genetics analyses have greatly advanced our understanding of CBLs and CIPKs and have uncovered crucial functions of these proteins for plant mineral nutrition and for responses to abiotic stresses and hormones, such as ABA (see Figure 4 for an illustrated summary). Characterization of *cb1* loss-of-function mutant lines revealed a function of CBL1 as a central integrator of responses to abiotic stresses, including drought, cold, and salt (Albrecht et al., 2003; Cheong et al., 2003). While the mutant studies of CBL1 revealed an ABA-independent function of this protein in several abiotic stress responses, loss of function of the closely related  $\text{Ca}^{2+}$  sensor CBL9 renders plants hypersensitive to ABA (Pandey et al., 2004). Alternative complex formation of the kinase CIPK1 with either CBL1 or CBL9 mediates ABA-dependent and ABA-independent signaling responses (D'Angelo et al., 2006). In addition, CBL9 appears to form a complex with CIPK3 for modulating ABA responses (Pandey et al., 2008).

A breakthrough study in 2006 by Wei-Hua Wu and colleagues uncovered a role of the CBL/CIPK system in regulating  $\text{K}^+$  homeostasis and provided the first molecular insights into how

plant ion channels might be regulated by phosphorylation (Xu et al., 2006). CIPK23 is targeted to the plasma membrane and activated by the two highly related  $\text{Ca}^{2+}$  sensors CBL1 and CBL9 (Xu et al., 2006; Cheong et al., 2007). These complexes then regulate the activity of the shaker-like potassium channel AKT1 (Li et al., 2006b; Xu et al., 2006). CIPK23 interacts exclusively with AKT1 but not with other  $\text{K}^+$  transporters from *Arabidopsis* (Hedrich and Kudla, 2006; Lee et al., 2007a; Geiger et al., 2009). Interaction analysis in yeast in combination with electrophysiological studies identified the 2C-type protein phosphatase AIP1 as a negative regulator of AKT1 that counteracts activation by CIPK23 (Lee et al., 2007a). Considering that AIP1 can interact with both CIPK23 and AKT1, it will be most interesting to distinguish if this activation/deactivation switch is brought about by phosphorylation/dephosphorylation or, alternatively, results from competitive binding of either the phosphatase or the kinase to the potassium channel. Besides the regulation of  $\text{K}^+$  uptake in roots, the  $\text{Ca}^{2+}$ -decoding CBL1/CBL9/CIPK23 module appears to be involved in stomata regulation under dehydrating conditions (Cheong et al., 2007).

An exciting novel twist in our understanding of CIPK23 function results from the recent report that this kinase also phosphorylates the nitrate transporter CHL1 (also called NRT1.1). Taking advantage of a novel CHL1 mutant allele (*chl1-9*), Yi-Fang Tsay and colleagues provided compelling evidence that this mutation impairs the nitrate uptake function of CHL1 without affecting the signaling response to nitrate as analyzed by the transcriptional response of the NRT2.1 gene (Ho et al., 2009). Importantly, biochemical and reverse genetics analyses identified CIPK23 as a critical regulator mediating the switch between low- and high-affinity nitrate transport modes by phosphorylating residue Thr-101 of CHL1. Specifically, at low external nitrate concentrations,



**Figure 4.** The CBL/CIPK Signaling System for Translating  $\text{Ca}^{2+}$  Signatures into Protein Phosphorylation.

Shown are characterized complexes consisting of CBL proteins and their interacting CIPKs. AKT1, *Arabidopsis*  $\text{K}^+$  transporter 1; SOS1, salt overly sensitive 1; CHL, a nitrate transporter; AHA2, *Arabidopsis*  $\text{H}^+$  ATPase 2.

CIPK23-mediated phosphorylation results in low-level nitrate signaling (Ho et al., 2009). Moreover, Ho et al. reported that CIPK23 is independently involved in potassium and nitrate responses, indicating a lack of crosstalk between both ions. Considering the critical involvement of the same CBL1 (and CBL9)  $\text{Ca}^{2+}$  sensor proteins and CIPK23-dependent phosphorylation in both processes, this puzzling observation underscores the urgent need for further investigation of the primary ion-sensing mechanism(s) that confer this remarkable specificity in plant signaling.

The plasma membrane  $\text{H}^{+}$ -ATPase AHA2 has been identified as an additional target of CBL/CIPK signaling complexes (Fuglsang et al., 2007). Phosphorylation of a Ser residue within the C-terminal domain of AHA2 by CIPK11 (designated as PKS5 by these authors) prevents binding of 14-3-3 proteins to this domain and leads to downregulation of AHA2 proton transport activity. Moreover, yeast two-hybrid interaction studies suggested a potential function of CBL2 in the CIPK11-dependent regulation of AHA2 (Fuglsang et al., 2007). However, since CBL2 has so far only been detected at the tonoplast in plants cells (Batistić et al., 2008, 2010), unambiguously determining the identity of the CBL protein mediating  $\text{Ca}^{2+}$ -dependent regulation of AHA2 in plants may require additional in planta or reverse genetics analyses.

Finally, recent work performed in rice further extended the identified physiological functions of CIPKs (Lee et al., 2009). The results from this study indicate that protein kinase Os CIPK15 plays a key role in  $\text{O}_2$  deficiency tolerance in rice and is required for growth of rice under flooded conditions. Moreover, CIPK15 regulates the plant global energy and stress sensor SnRK1A, thereby integrating responses to  $\text{O}_2$  deficiency with sugar signaling and enabling rice growth under floodwater. The latter finding may point to a general role of the CBL/CIPK system in fine-tuning plant metabolism in response to adverse environmental conditions.

The results of all these studies suggest that the CBL/CIPK network represents a central and critical signaling system for decoding  $\text{Ca}^{2+}$  signals in response to a wide range of stimuli. So far, research on CBLs and CIPKs has been focused mainly on abiotic stress responses and ion uptake. However, it is safe to predict that future studies will most likely uncover a role of this  $\text{Ca}^{2+}$  decoding system in additional processes, such as plant pathogen interactions. It is now well established that each CBL and each CIPK represents a multifunctional signaling component that can undergo alternative protein interactions determining the flow of information processing through this signaling system. Therefore, furthering our understanding of this signaling system will require the combined application of state-of-the-art cell biological techniques that enable the monitoring of dynamic protein-protein interactions and protein targeting. Such approaches should also allow the elucidation of mechanistic factors that determine the decision making in this flexible interaction network and will be of primary importance to advance our understanding of  $\text{Ca}^{2+}$  decoding mechanisms.

### Converting $\text{Ca}^{2+}$ Signals into Transcriptional Responses

Transcriptional regulation is coupled with numerous signaling processes that can be conveyed by several second messengers and the function of  $\text{Ca}^{2+}$  binding proteins (Ikura et al., 2002).

Despite the obvious importance of defined  $\text{Ca}^{2+}$  signatures in eliciting specific transcriptional response reactions in plants to stimuli, such as cold, drought, and salt stress, the molecular mechanisms mediating  $\text{Ca}^{2+}$ -responsive gene expression have long remained little understood (Scrase-Field and Knight, 2003). This is due in part to the difficulty in discriminating between stress-dependent  $\text{Ca}^{2+}$  responses per se and stress-dependent but  $\text{Ca}^{2+}$ -independent transcriptional reactions (Finkler et al., 2007b). Recently, the induction of defined artificial  $\text{Ca}^{2+}$  transients in response to calmodulin antagonists like WP7 and SKF-7171 has allowed the identification of 230  $\text{Ca}^{2+}$ -responsive genes that were differentially expressed 1 h after stimulus in *Arabidopsis* (Kaplan et al., 2006). Since the microarrays used in this study covered only 25% of the known *Arabidopsis* genes, this finding suggests that ~3% of the protein-coding genes in the *Arabidopsis* genome are subject to regulation by  $\text{Ca}^{2+}$ . Remarkably, many of the genes identified in this study as being  $\text{Ca}^{2+}$  regulated were previously characterized as early stress-induced genes.

Within the promoter regions of  $\text{Ca}^{2+}$ -regulated genes, this study identified abscisic acid-responsive element (ABRE)-related *cis*-elements as being sufficient to confer transcriptional regulation in response to cytosolic  $\text{Ca}^{2+}$  signatures. This finding is significant because such ABREs have been identified in the promoter of CBF/DREB1 transcription factors that function as master regulators of abiotic stress responses (Finkler et al., 2007b). Therefore, these results point to a direct interconnection of  $\text{Ca}^{2+}$ -regulated gene transcription and abiotic stress responses. Such an immediate conversion of  $\text{Ca}^{2+}$  signatures into transcriptional regulation may be in part achieved by the function of calmodulin binding transcription factors (CAMTA proteins).


### $\text{Ca}^{2+}$ -Regulated Transcription Factors

CAMTAs are  $\text{Ca}^{2+}$ -dependent CaM binding transcription factors forming a family of six members in *Arabidopsis* (Finkler et al., 2007a). They share a conserved domain structure including an N-terminal CG-1 domain mediating binding to DNA *cis*-elements (CAMTA binding sites), including ABREs (Finkler et al., 2007a). The C-terminal CaM binding domain of CAMTAs mediates interactions with calmodulin. First insights into the physiological function of plant CAMTA proteins came from a reverse genetic analysis of *Arabidopsis* CAMTA3 function that revealed a critical role of this protein in suppressing plant responses to pathogens such as *P. syringae* and *Botrytis cinerea* (Galon et al., 2008). Most recent work on CAMTA3 (designated At SR1) uncovered that CAMTA3 directly interacts with the promoter of the *EDS1* gene, an established regulator of salicylic acid levels, and represses its expression (Figure 5) (Du et al., 2009). Moreover,  $\text{Ca}^{2+}$ /calmodulin binding to CAMTA3 was required for the suppression of plant defense, indicating a direct role of  $\text{Ca}^{2+}$ /calmodulin in regulating the function of CAMTA3 (Du et al., 2009). Important evidence for a direct link between  $\text{Ca}^{2+}$  signaling via CAMTA1 and CAMTA3 and the regulation of cold tolerance in plants was provided by the discovery that these CAMTA proteins bind to regulatory elements in the promoter region of the *DREB1c/CBF2* gene (Doherty et al., 2009). A single *camta3* mutant exhibited a

significant reduction in cold induction of a number of known cold-induced genes, including *CBF2*. Moreover, *camta1 camta3* double mutant plants were impaired in their cold acclimation to freezing tolerance (Doherty et al., 2009). These findings establish a crucial role of  $\text{Ca}^{2+}$ /calmodulin-regulated CAMTA transcription factors in controlling CBF-regulated cold-responsive gene expression in plants (Figure 5). Direct decoding of cold-induced  $\text{Ca}^{2+}$  signatures into the regulation of gene expression may occur through direct and  $\text{Ca}^{2+}$ -dependent interaction of CAMTAs with members of the *Arabidopsis* calmodulin  $\text{Ca}^{2+}$  sensors family (McCormack et al., 2005). The function of calmodulin in the regulation of gene expression by direct interaction with transcription factors appears not to be restricted to their interaction with CAMTAs because interactions of calmodulin with MYB and WRKY transcription factors has also been reported (Park et al., 2005; Yoo et al., 2005). In this regard, it will be important to investigate exactly how the stimulus-induced changes in  $\text{Ca}^{2+}$  concentration that occur in the cytoplasm are connected into transcriptional regulation that proceeds in the nucleus.

### Transcriptional Regulation by Calmodulin

Calmodulins are conserved eukaryotic  $\text{Ca}^{2+}$  sensor relay proteins that upon  $\text{Ca}^{2+}$  binding undergo extensive conformational changes that result in protein interactions with their target proteins (Luan et al., 2002; McCormack et al., 2005). In *Arabidopsis*, seven genes encode four CAM isoforms, of which CAM1/CAM4 differ by four amino acid substitutions from CAM7, whereas CAM2/3/5 and CAM6 differ in one amino acid position from CAM7 (McCormack et al., 2005). Classical microinjection experiments into hypocotyl cells of the phytochrome deficient tomato *aurea* mutant suggested a function of  $\text{Ca}^{2+}$  and calmodulin in light signal transduction that controls regulation of light-responsive gene expression (Neuhaus et al., 1993; Bowler et al., 1994). A recent study of CAM7 from *Arabidopsis* revealed surprising insights into the regulation of light-induced gene expression by calmodulin (Kushwaha et al., 2008). The authors identified specifically CAM7 but not CAM2/3/5 as a transcrip-



**Figure 5.** Converting  $\text{Ca}^{2+}$  Signals into Transcriptional Responses.

Recent studies have unveiled that  $\text{Ca}^{2+}$ /CAMs (calmodulins) regulate CAMTAs that interact with promoters of abiotic stress-responsive genes. Interestingly, *Arabidopsis* CAM7 is likely a direct converter of cytoplasmic  $\text{Ca}^{2+}$  signatures into regulation of gene expression. *EDS1*, enhanced disease susceptibility 1.

tional regulator that directly interacts with promoters of several light-inducible genes (Figure 5). In accordance with such a function of CAM7, *cam7* mutants exhibited a reduced expression of light-inducible genes. Conversely, overexpression of CAM7 resulted in an increase in the expression of light-inducible genes. The findings of this study suggest a direct translation of cytoplasmic  $\text{Ca}^{2+}$  signatures by the  $\text{Ca}^{2+}$  sensor responder protein CAM7 into regulation of gene expression by DNA binding. To further elucidate details of this regulatory circuit, it will be most interesting to investigate if this function requires the translocation of CAM7 from the cytoplasm to the nucleus and/or the interaction of CAM7 with additional (transcription factor) proteins.

### Conclusions and Future Perspectives

During the past years we have witnessed tremendous progress in our understanding of  $\text{Ca}^{2+}$  signaling processes in plants. When comparing research on  $\text{Ca}^{2+}$  signaling in animals and plants during the past decade, remarkably different developments become apparent. Whereas in animals the components and mechanisms that exercise primary  $\text{Ca}^{2+}$ -releasing events and shape  $\text{Ca}^{2+}$  signatures have seen the most significant advances, in plant research, most progress has been achieved in identifying and characterizing components that decode  $\text{Ca}^{2+}$  signals.

A central question around which  $\text{Ca}^{2+}$  signaling research in plants is centered is: How can one ion specifically control so many different processes and events? Current research is beginning to provide answers.  $\text{Ca}^{2+}$  signaling in plants involves many facets that can define and adjust responses in both time and space. The unequal distribution of this ion in the cell provides the basis for rapid  $\text{Ca}^{2+}$  fluxes and the resulting concentration changes. Many energized cation transporters and channels that were previously assumed to merely be regulators of  $\text{Ca}^{2+}$  homeostasis actually appear to play a role in  $\text{Ca}^{2+}$  signaling. However, a major bottleneck in our understanding of the generation of  $\text{Ca}^{2+}$  signals is the paucity of molecular information about true  $\text{Ca}^{2+}$  channels. It may require novel genetic screens or the application of chemical genetic screens (that have been successful, for example, in identifying plant ABA receptors) to successfully tackle this serious limitation of plant  $\text{Ca}^{2+}$  signaling research.

It is now widely accepted that complex families of  $\text{Ca}^{2+}$  binding proteins that function as calcium sensors provide the toolkit for deciphering various  $\text{Ca}^{2+}$  signatures. Calmodulins, CMLs, CDPKs, and CBL/CIPK complexes form intricate signaling networks for translating these signatures into downstream phosphorylation events and transcriptional responses. An important role of  $\text{Ca}^{2+}$ -regulated transcription factors like CAMTAs in these processes is emerging. It is very likely the network-like character of these signaling systems that warrants flexible but robust information processing. Accumulating evidence also points to two levels at which  $\text{Ca}^{2+}$  signaling is operative. Within specific cells, such as guard cell or root hairs, intracellular  $\text{Ca}^{2+}$  signaling processes occur at high spatial and temporal specificity. This layer of  $\text{Ca}^{2+}$  signaling appears to be superimposed on the tissue or organismic level where specific tissue layers or cell types exhibit defined and prominent changes in  $\text{Ca}^{2+}$  concentration.

It has become evident that parameters such as the specific  $\text{Ca}^{2+}$  binding affinity, the specific cellular concentration and

subcellular localization, and the specific interaction affinities of  $\text{Ca}^{2+}$  decoders are critical components that contribute to generating specificity in signal-to-response coupling. However, a major challenge that remains is to understand mechanistically how exactly calcium signatures or variable oscillations are sensed and transduced by calcium sensor proteins. Tackling this important task will require serious investments in structural analyses of  $\text{Ca}^{2+}$  sensors, NMR-based studies of their structural dynamics, extensive investigation of  $\text{Ca}^{2+}$  binding and  $\text{Ca}^{2+}$ -dissociation parameters of  $\text{Ca}^{2+}$  sensors, and novel cell biological approaches to visualize the dynamics of protein interactions and protein complex formation in planta. The integration of these data with advanced fluorescence resonance energy transfer-based monitoring of cellular  $\text{Ca}^{2+}$  dynamics and the mathematical modeling of such processes represent the route to be pursued in plant  $\text{Ca}^{2+}$  signaling research in the near future. Currently, despite all the exciting progress described in this review article, we are still not close to finally cracking the calcium code in plants.

#### ACKNOWLEDGMENTS

This work was supported by the Deutsche Forschungsgemeinschaft, the Alexander von Humboldt Foundation, the Deutscher Akademischer Austauschdienst, and the Human Frontier of Science Program.

Received November 11, 2009; revised March 12, 2010; accepted March 14, 2010; published March 30, 2010.

#### REFERENCES

- Albrecht, V., Ritz, O., Linder, S., Harter, K., and Kudla, J. (2001). The NAF domain defines a novel protein-protein interaction module conserved in  $\text{Ca}^{2+}$ -regulated kinases. *EMBO J.* **20**: 1051–1063.
- Albrecht, V., Weinl, S., Blazevic, D., D'Angelo, C., Batistič, O., Kolukisaoglu, U., Bock, R., Schulz, B., Harter, K., and Kudla, J. (2003). The calcium sensor CBL1 integrates plant responses to abiotic stresses. *Plant J.* **36**: 457–470.
- Ali, R., Ma, W., Lemtiri-Chlieh, F., Tsaltas, D., Leng, Q., von Bodman, S., and Berkowitz, G.A. (2007). Death don't have no mercy and neither does calcium: *Arabidopsis* CYCLIC NUCLEOTIDE GATED CHANNEL2 and innate immunity. *Plant Cell* **19**: 1081–1095.
- Ali, R., Zielinski, R.E., and Berkowitz, G.A. (2006). Expression of plant cyclic nucleotide-gated cation channels in yeast. *J. Exp. Bot.* **57**: 125–138.
- Allen, G.J., Chu, S.P., Harrington, C.L., Schumacher, K., Hoffmann, T., Tang, Y.Y., Grill, E., and Schroeder, J.I. (2001). A defined range of guard cell calcium oscillation parameters encodes stomatal movements. *Nature* **411**: 1053–1057.
- Allen, G.J., Muir, S.R., and Sanders, D. (1995). Release of  $\text{Ca}^{2+}$  from individual plant vacuoles by both InsP3 and cyclic ADP-ribose. *Science* **268**: 735–737.
- Allen, G.J., and Sanders, D. (1994). Two voltage-gated, calcium release channels coreside in the vacuolar membrane of broad bean guard cells. *Plant Cell* **6**: 685–694.
- Allen, G.J., and Sanders, D. (1995). Calcineurin, a Type 2B protein phosphatase, modulates the  $\text{Ca}^{2+}$ -permeable slow vacuolar ion channel of stomatal guard cells. *Plant Cell* **7**: 1473–1483.
- Balague, C., Lin, B., Alcon, C., Flottes, G., Malmstrom, S., Kohler, C., Neuhaus, G., Pelletier, G., Gaymard, F., and Roby, D. (2003). HLM1, an essential signaling component in the hypersensitive response, is a member of the cyclic nucleotide-gated channel ion channel family. *Plant Cell* **15**: 365–379.
- Batelli, G., Verslues, P.E., Agius, F., Qiu, Q., Fujii, H., Pan, S., Schumaker, K.S., Grillo, S., and Zhu, J.K. (2007). SOS2 promotes salt tolerance in part by interacting with the vacuolar  $\text{H}^{+}$ -ATPase and upregulating its transport activity. *Mol. Cell. Biol.* **27**: 7781–7790.
- Batistič, O., and Kudla, J. (2004). Integration and channeling of calcium signaling through the CBL calcium sensor/CIPK protein kinase network. *Planta* **219**: 915–924.
- Batistič, O., and Kudla, J. (2009). Plant calcineurin B-like proteins and their interacting protein kinases. *Biochim. Biophys. Acta* **1793**: 985–992.
- Batistič, O., Sorek, N., Schultke, S., Yalovsky, S., and Kudla, J. (2008). Dual fatty acyl modification determines the localization and plasma membrane targeting of CBL/CIPK  $\text{Ca}^{2+}$  signaling complexes in *Arabidopsis*. *Plant Cell* **20**: 1346–1362.
- Batistič, O., Waadt, R., Steinhorst, L., Held, K., and Kudla, J. (2010). CBL-mediated targeting of CIPKs facilitates the decoding of calcium signals emanating from distinct cellular stores. *Plant J.* **61**: 211–222.
- Berridge, M.J. (2006). Calcium microdomains: Organization and function. *Cell Calcium* **40**: 405–412.
- Berridge, M.J. (2009). Inositol trisphosphate and calcium signalling mechanisms. *Biochim. Biophys. Acta* **1793**: 933–940.
- Beyhl, D., Hortensteiner, S., Martinoia, E., Farmer, E.E., Fromm, J., Marten, I., and Hedrich, R. (2009). The *fou2* mutation in the major vacuolar cation channel TPC1 confers tolerance to inhibitory luminal calcium. *Plant J.* **58**: 715–723.
- Bihler, H., Eing, C., Hebeisen, S., Roller, A., Czempinski, K., and Bertl, A. (2005). TPK1 is a vacuolar ion channel different from the slow-vacuolar cation channel. *Plant Physiol.* **139**: 417–424.
- Billker, O., Dechamps, S., Tewari, R., Wenig, G., Franke-Fayard, B., and Brinkmann, V. (2004). Calcium and a calcium-dependent protein kinase regulate gamete formation and mosquito transmission in a malaria parasite. *Cell* **117**: 503–514.
- Blatt, M.R., Thiel, G., and Trentham, D.R. (1990). Reversible inactivation of  $\text{K}^{+}$  channels of *Vicia* stomatal guard cells following the photolysis of caged inositol 1,4,5-trisphosphate. *Nature* **346**: 766–769.
- Bonaventure, G., Gfeller, A., Proebsting, W.M., Hortensteiner, S., Chetelat, A., Martinoia, E., and Farmer, E.E. (2007a). A gain-of-function allele of *TPC1* activates oxylipin biogenesis after leaf wounding in *Arabidopsis*. *Plant J.* **49**: 889–898.
- Bonaventure, G., Gfeller, A., Rodriguez, V.M., Armand, F., and Farmer, E.E. (2007b). The *fou2* gain-of-function allele and the wild-type allele of *Two Pore Channel 1* contribute to different extents or by different mechanisms to defense gene expression in *Arabidopsis*. *Plant Cell Physiol.* **48**: 1775–1789.
- Bonza, M.C., Morandini, P., Luoni, L., Geisler, M., Palmgren, M.G., and De Michelis, M.I. (2000). *At-ACA8* encodes a plasma membrane-localized calcium-ATPase of *Arabidopsis* with a calmodulin-binding domain at the N terminus. *Plant Physiol.* **123**: 1495–1506.
- Bourgeade, P., de Jaeger, G., and Boyer, N. (1991). Microsomal ATP-dependent  $\text{Ca}^{2+}$  transport as affected by environmental stress in *Bryonia dioica* internodes. *Plant Sci.* **79**: 23–30.
- Bowler, C., Neuhaus, G., Yamagata, H., and Chua, N.H. (1994). Cyclic GMP and calcium mediate phytochrome phototransduction. *Cell* **77**: 73–81.
- Braam, J. (2005). In touch: plant responses to mechanical stimuli. *New Phytol.* **165**: 373–389.
- Brand, J.J., and Becker, D.W. (1984). Evidence for direct roles of calcium in photosynthesis. *J. Bioenerg. Biomembr.* **16**: 239–249.

- Brailoiu, E., Churamani, D., Cai, X., Schrlau, M.G., Brailoiu, G.C., Gao, X., Hooper, R., Boulware, M.J., Dun, N.J., Marchant, J.S., and Patel, S. (2009). Essential requirement for two-pore channel 1 in NAADP-mediated calcium signaling. *J. Cell Biol.* **186**: 201–209.
- Brenner, E.D., Martinez-Barboza, N., Clark, A.P., Liang, Q.S., Stevenson, D.W., and Coruzzi, G.M. (2000). *Arabidopsis* mutants resistant to S<sup>+</sup>-beta-methyl-alpha, beta-diaminopropionic acid, a cycad-derived glutamate receptor agonist. *Plant Physiol.* **124**: 1615–1624.
- Bush, D.R., and Sze, H. (1986). Calcium transport in tonoplast and endoplasmic reticulum vesicles isolated from cultured carrot cells. *Plant Physiol.* **80**: 549–555.
- Bush, D.S., Biswas, A.K., and Jones, R.L. (1989a). Gibberellic-acid-stimulated Ca<sup>2+</sup> accumulation in endoplasmic reticulum of barley aleurone: Ca<sup>2+</sup> transport and steady-state levels. *Planta* **178**: 411–420.
- Bush, D.S., Biswas, A.K., and Jones, R.L. (1993). Hormonal regulation of Ca<sup>2+</sup> transport in the endomembrane system of the barley aleurone. *Planta* **189**: 507–515.
- Bush, D.S., Sticher, L., van Huystee, R., Wagner, D., and Jones, R.L. (1989b). The calcium requirement for stability and enzymatic activity of two isoforms of barley aleurone alpha-amylase. *J. Biol. Chem.* **264**: 19392–19398.
- Calcraft, P.J., et al. (2009). NAADP mobilizes calcium from acidic organelles through two-pore channels. *Nature* **459**: 596–600.
- Cardenas, L., Lovy-Wheeler, A., Kunkel, J.G., and Hepler, P.K. (2008). Pollen tube growth oscillations and intracellular calcium levels are reversibly modulated by actin polymerization. *Plant Physiol.* **146**: 1611–1621.
- Catala, R., Santos, E., Alonso, J.M., Ecker, J.R., Martinez-Zapater, J.M., and Salinas, J. (2003). Mutations in the Ca<sup>2+</sup>/H<sup>+</sup> transporter CAX1 increase *CBF/DREB1* expression and the cold-acclimation response in *Arabidopsis*. *Plant Cell* **15**: 2940–2951.
- Cerana, M., Bonza, M.C., Harris, R., Sanders, D., and De Michelis, M.I. (2006). Abscisic acid stimulates the expression of two isoforms of plasma membrane Ca<sup>2+</sup>-ATPase in *Arabidopsis thaliana* seedlings. *Plant Biol.* **8**: 572–578.
- Charron, D., Pingret, J.L., Chabaud, M., Journet, E.P., and Barker, D. G. (2004). Pharmacological evidence that multiple phospholipid signaling pathways link Rhizobium nodulation factor perception in *Medicago truncatula* root hairs to intracellular responses, including Ca<sup>2+</sup> spiking and specific *ENOD* gene expression. *Plant Physiol.* **136**: 3582–3593.
- Cheng, N.H., and Hirschi, K.D. (2003). Cloning and characterization of CXIP1, a novel PICOT domain-containing *Arabidopsis* protein that associates with CAX1. *J. Biol. Chem.* **278**: 6503–6509.
- Cheng, N.H., Liu, J.Z., Nelson, R.S., and Hirschi, K.D. (2004a). Characterization of CXIP4, a novel *Arabidopsis* protein that activates the H<sup>+</sup>/Ca<sup>2+</sup> antiporter, CAX1. *FEBS Lett.* **559**: 99–106.
- Cheng, N.H., Pittman, J.K., Barkla, B.J., Shigaki, T., and Hirschi, K.D. (2003). The *Arabidopsis cax1* mutant exhibits impaired ion homeostasis, development, and hormonal responses and reveals interplay among vacuolar transporters. *Plant Cell* **15**: 347–364.
- Cheng, N.H., Pittman, J.K., Shigaki, T., and Hirschi, K.D. (2002). Characterization of CAX4, an *Arabidopsis* H<sup>+</sup>/cation antiporter. *Plant Physiol.* **128**: 1245–1254.
- Cheng, N.H., Pittman, J.K., Shigaki, T., Lachmansingh, J., LeClere, S., Lahner, B., Salt, D.E., and Hirschi, K.D. (2005). Functional association of *Arabidopsis* CAX1 and CAX3 is required for normal growth and ion homeostasis. *Plant Physiol.* **138**: 2048–2060.
- Cheng, N.H., Pittman, J.K., Zhu, J.K., and Hirschi, K.D. (2004b). The protein kinase SOS2 activates the *Arabidopsis* H<sup>+</sup>/Ca<sup>2+</sup> antiporter CAX1 to integrate calcium transport and salt tolerance. *J. Biol. Chem.* **279**: 2922–2926.
- Cheong, Y.H., Kim, K.N., Pandey, G.K., Gupta, R., Grant, J.J., and Luan, S. (2003). CBL1, a calcium sensor that differentially regulates salt, drought, and cold responses in *Arabidopsis*. *Plant Cell* **15**: 1833–1845.
- Cheong, Y.H., Pandey, G.K., Grant, J.J., Batistić, O., Li, L., Kim, B.G., Lee, S.C., Kudla, J., and Luan, S. (2007). Two calcineurin B-like calcium sensors, interacting with protein kinase CIPK23, regulate leaf transpiration and root potassium uptake in *Arabidopsis*. *Plant J.* **52**: 223–239.
- Chiu, J.C., Brenner, E.D., DeSalle, R., Nitabach, M.N., Holmes, T.C., and Coruzzi, G.M. (2002). Phylogenetic and expression analysis of the glutamate-receptor-like gene family in *Arabidopsis thaliana*. *Mol. Biol. Evol.* **19**: 1066–1082.
- Cho, D., Kim, S.A., Murata, Y., Lee, S., Jae, S.K., Nam, H.G., and Kwak, J.M. (2009). De-regulated expression of the plant glutamate receptor homolog *AtGLR3.1* impairs long-term Ca<sup>2+</sup>-programmed stomatal closure. *Plant J.* **58**: 437–449.
- Clapham, D.E. (2007). Calcium signaling. *Cell* **131**: 1047–1058.
- Clough, S.J., Fengler, K.A., Yu, I.C., Lippok, B., Smith, R.K., Jr., and Bent, A.F. (2000). The *Arabidopsis dnd1* “defense, no death” gene encodes a mutated cyclic nucleotide-gated ion channel. *Proc. Natl. Acad. Sci. USA* **97**: 9323–9328.
- Cosgrove, D.J., and Hedrich, R. (1991). Stretch-activated chloride, potassium, and calcium channels coexisting in plasma membranes of guard cells of *Vicia faba* L. *Planta* **186**: 143–153.
- D’Angelo, C., et al. (2006). Alternative complex formation of the Ca<sup>2+</sup>-regulated protein kinase CIPK1 controls abscisic acid-dependent and independent stress responses in *Arabidopsis*. *Plant J.* **48**: 857–872.
- Delk, N.A., Johnson, K.A., Chowdhury, N.I., and Braam, J. (2005). CML24, regulated in expression by diverse stimuli, encodes a potential Ca<sup>2+</sup> sensor that functions in responses to abscisic acid, day-length, and ion stress. *Plant Physiol.* **139**: 240–253.
- Demidchik, V., Bowen, H.C., Maathuis, F.J., Shabala, S.N., Tester, M.A., White, P.J., and Davies, J.M. (2002). *Arabidopsis thaliana* root non-selective cation channels mediate calcium uptake and are involved in growth. *Plant J.* **32**: 799–808.
- Demidchik, V., and Maathuis, F.J. (2007). Physiological roles of nonselective cation channels in plants: From salt stress to signalling and development. *New Phytol.* **175**: 387–404.
- Dobney, S., Chiasson, D., Lam, P., Smith, S.P., and Snedden, W.A. (2009). The calmodulin-related calcium sensor CML42 plays a role in trichome branching. *J. Biol. Chem.* **284**: 31647–31657.
- Dodd, A.N., Jakobsen, M.K., Baker, A.J., Telzerow, A., Hou, S.W., Laplaze, L., Barrot, L., Poethig, R.S., Haseloff, J., and Webb, A.A. (2006). Time of day modulates low-temperature Ca signals in *Arabidopsis*. *Plant J.* **48**: 962–973.
- Doherty, C.J., Van Buskirk, H.A., Myers, S.J., and Thomashow, M.F. (2009). Roles for *Arabidopsis* CAMTA transcription factors in cold-regulated gene expression and freezing tolerance. *Plant Cell* **21**: 972–984.
- Drobak, B.K., and Ferguson, I.B. (1985). Release of Ca<sup>2+</sup> from plant hypocotyl microsomes by inositol-1,4,5-trisphosphate. *Biochem. Biophys. Res. Commun.* **130**: 1241–1246.
- Du, L., Ali, G.S., Simons, K.A., Hou, J., Yang, T., Reddy, A.S., and Poovaiah, B.W. (2009). Ca<sup>2+</sup>/calmodulin regulates salicylic-acid-mediated plant immunity. *Nature* **457**: 1154–1158.
- Du, L., and Poovaiah, B.W. (2005). Ca<sup>2+</sup>/calmodulin is critical for brassinosteroid biosynthesis and plant growth. *Nature* **437**: 741–745.
- Dutta, R., and Robinson, K.R. (2004). Identification and characterization of stretch-activated ion channels in pollen protoplasts. *Plant Physiol.* **135**: 1398–1406.
- Engstrom, E.M., Ehrhardt, D.W., Mitra, R.M., and Long, S.R. (2002). Pharmacological analysis of nod factor-induced calcium spiking

- in *Medicago truncatula*. Evidence for the requirement of type IIA calcium pumps and phosphoinositide signaling. *Plant Physiol.* **128**: 1390–1401.
- Erdei, L., Toth, I., and Zsoldos, F.** (1979). Hormonal regulation of  $\text{Ca}^{2+}$ -stimulated  $\text{K}^+$  influx and  $\text{Ca}^{2+}$ ,  $\text{K}^+$ -ATPase in rice roots: *in vivo* and *in vitro* effects of auxins and reconstitution of the ATPase. *Physiol. Plant.* **45**: 448–452.
- Fakhr, H., and Hall, J.L.** (1984). Changes in calcium-ATPase activity associated with the washing (ageing) of potato tuber discs. *J. Plant Physiol.* **117**: 69–79.
- Finkler, A., Ashery-Padan, R., and Fromm, H.** (2007a). CAMTAs: Calmodulin-binding transcription activators from plants to human. *FEBS Lett.* **581**: 3893–3898.
- Finkler, A., Kaplan, B., and Fromm, H.** (2007b).  $\text{Ca}^{2+}$ -responsive *cis*-elements in plants. *Plant Signal. Behav.* **2**: 17–19.
- Frietsch, S., Wang, Y.F., Sladek, C., Poulsen, L.R., Romanowsky, S. M., Schroeder, J.I., and Harper, J.F.** (2007). A cyclic nucleotide-gated channel is essential for polarized tip growth of pollen. *Proc. Natl. Acad. Sci. USA* **104**: 14531–14536.
- Fuglsang, A.T., Guo, Y., Cuin, T.A., Qiu, Q., Song, C., Kristiansen, K. A., Bych, K., Schulz, A., Shabala, S., Schumaker, K.S., Palmgren, M.G., and Zhu, J.K.** (2007). *Arabidopsis* protein kinase PKS5 inhibits the plasma membrane  $\text{H}^+$ -ATPase by preventing interaction with 14-3-3 protein. *Plant Cell* **19**: 1617–1634.
- Furch, A.C., van Bel, A.J., Fricker, M.D., Felle, H.H., Fuchs, M., and Hafke, J.B.** (2009). Sieve element  $\text{Ca}^{2+}$  channels as relay stations between remote stimuli and sieve tube occlusion in *Vicia faba*. *Plant Cell* **21**: 2118–2132.
- Galon, Y., Nave, R., Boyce, J.M., Nachmias, D., Knight, M.R., and Fromm, H.** (2008). Calmodulin-binding transcription activator (CAMTA) 3 mediates biotic defense responses in *Arabidopsis*. *FEBS Lett.* **582**: 943–948.
- Gao, D., Knight, M.R., Trewavas, A.J., Sattelmacher, B., and Plieth, C.** (2004). Self-reporting *Arabidopsis* expressing pH and  $[\text{Ca}^{2+}]$  indicators unveil ion dynamics in the cytoplasm and in the apoplast under abiotic stress. *Plant Physiol.* **134**: 898–908.
- Geiger, D., Becker, D., Vosloh, D., Gambale, F., Palme, K., Rehers, M., Anschuetz, U., Dreyer, I., Kudla, J., and Hedrich, R.** (2009). Heteromeric AtKC1/AKT1 channels in *Arabidopsis* roots facilitate growth under  $\text{K}^+$ -limiting conditions. *J. Biol. Chem.* **284**: 21288–21295.
- Geisler, M., Frangne, N., Gomes, E., Martinoia, E., and Palmgren, M. G.** (2000). The *ACA4* gene of *Arabidopsis* encodes a vacuolar membrane calcium pump that improves salt tolerance in yeast. *Plant Physiol.* **124**: 1814–1827.
- George, L., Romanowsky, S.M., Harper, J.F., and Sharrock, R.A.** (2008). The *ACA10*  $\text{Ca}^{2+}$ -ATPase regulates adult vegetative development and inflorescence architecture in *Arabidopsis*. *Plant Physiol.* **146**: 716–728.
- Gilroy, S., Read, N.D., and Trewavas, A.J.** (1990). Elevation of cytoplasmic calcium by caged calcium or caged inositol triphosphate initiates stomatal closure. *Nature* **346**: 769–771.
- Gleason, C., Chaudhuri, S., Yang, T., Munoz, A., Poovaiah, B.W., and Oldroyd, G.E.** (2006). Nodulation independent of rhizobia induced by a calcium-activated kinase lacking autoinhibition. *Nature* **441**: 1149–1152.
- Gobert, A., Park, G., Amtmann, A., Sanders, D., and Maathuis, F.J.** (2006). *Arabidopsis thaliana* cyclic nucleotide gated channel 3 forms a non-selective ion transporter involved in germination and cation transport. *J. Exp. Bot.* **57**: 791–800.
- Gong, D., Guo, Y., Jagendorf, A.T., and Zhu, J.K.** (2002). Biochemical characterization of the *Arabidopsis* protein kinase SOS2 that functions in salt tolerance. *Plant Physiol.* **130**: 256–264.
- Grabov, A., and Blatt, M.R.** (1998). Membrane voltage initiates  $\text{Ca}^{2+}$  waves and potentiates  $\text{Ca}^{2+}$  increases with abscisic acid in stomatal guard cells. *Proc. Natl. Acad. Sci. USA* **95**: 4778–4783.
- Guo, K.M., Babourina, O., Christopher, D.A., Borsics, T., and Rengel, Z.** (2008). The cyclic nucleotide-gated channel, AtCNGC10, influences salt tolerance in *Arabidopsis*. *Physiol. Plant.* **134**: 499–507.
- Guo, Y., Halfter, U., Ishitani, M., and Zhu, J.K.** (2001). Molecular characterization of functional domains in the protein kinase SOS2 that is required for plant salt tolerance. *Plant Cell* **13**: 1383–1400.
- Halfter, U., Ishitani, M., and Zhu, J.K.** (2000). The *Arabidopsis* SOS2 protein kinase physically interacts with and is activated by the calcium-binding protein SOS3. *Proc. Natl. Acad. Sci. USA* **97**: 3735–3740.
- Hamilton, D.W., Hills, A., Kohler, B., and Blatt, M.R.** (2000).  $\text{Ca}^{2+}$  channels at the plasma membrane of stomatal guard cells are activated by hyperpolarization and abscisic acid. *Proc. Natl. Acad. Sci. USA* **97**: 4967–4972.
- Han, S., Tang, R., Anderson, L.K., Woerner, T.E., and Pei, Z.M.** (2003). A cell surface receptor mediates extracellular  $\text{Ca}^{2+}$  sensing in guard cells. *Nature* **425**: 196–200.
- Harada, A., and Shimazaki, K.** (2009). Measurement of changes in cytosolic  $\text{Ca}^{2+}$  in *Arabidopsis* guard cells and mesophyll cells in response to blue light. *Plant Cell Physiol.* **50**: 360–373.
- Harmon, A.C., Gribskov, M., and Harper, J.F.** (2000). CDPKs: A kinase for every  $\text{Ca}^{2+}$  signal? *Trends Plant Sci.* **5**: 154–159.
- Harper, J.F., Breton, G., and Harmon, A.** (2004). Decoding  $\text{Ca}^{2+}$  signals through plant protein kinases. *Annu. Rev. Plant Biol.* **55**: 263–288.
- Harper, J.F., and Harmon, A.** (2005). Plants, symbiosis and parasites: A calcium signalling connection. *Nat. Rev. Mol. Cell Biol.* **6**: 555–566.
- Harper, J.F., Hong, B., Hwang, I., Guo, H.Q., Stoddard, R., Huang, J.F., Palmgren, M.G., and Sze, H.** (1998). A novel calmodulin-regulated  $\text{Ca}^{2+}$ -ATPase (*ACA2*) from *Arabidopsis* with an N-terminal autoinhibitory domain. *J. Biol. Chem.* **273**: 1099–1106.
- Hedrich, R., and Kudla, J.** (2006). Calcium signaling networks channel plant  $\text{K}^+$  uptake. *Cell* **125**: 1221–1223.
- Hedrich, R., and Neher, E.** (1987). Cytoplasmic calcium regulates voltage-dependent ion channels in plant vacuoles. *Nature* **329**: 833–836.
- Hepler, P.K., Vidali, L., and Cheung, A.Y.** (2001). Polarized cell growth in higher plants. *Annu. Rev. Cell Dev. Biol.* **17**: 159–187.
- Hetherington, A.M., and Brownlee, C.** (2004). The generation of  $\text{Ca}^{2+}$  signals in plants. *Annu. Rev. Plant Biol.* **55**: 401–427.
- Hirschi, K.D.** (1999). Expression of *Arabidopsis* CAX1 in tobacco: Altered calcium homeostasis and increased stress sensitivity. *Plant Cell* **11**: 2113–2122.
- Hirschi, K.D., Korenkov, V.D., Wilganowski, N.L., and Wagner, G.J.** (2000). Expression of *Arabidopsis* CAX2 in tobacco. Altered metal accumulation and increased manganese tolerance. *Plant Physiol.* **124**: 125–133.
- Ho, C.H., Lin, S.H., Hu, H.C., and Tsay, Y.F.** (2009). CHL1 functions as a nitrate sensor in plants. *Cell* **138**: 1184–1194.
- Hrabak, E.M., et al.** (2003). The *Arabidopsis* CDPK-SnRK superfamily of protein kinases. *Plant Physiol.* **132**: 666–680.
- Hua, B.G., Mercier, R.W., Leng, Q., and Berkowitz, G.A.** (2003a). Plants do it differently. A new basis for potassium/sodium selectivity in the pore of an ion channel. *Plant Physiol.* **132**: 1353–1361.
- Hua, B.G., Mercier, R.W., Zielinski, R.E., and Berkowitz, G.A.** (2003b). Functional interaction of calmodulin with a plant cyclic nucleotide gated cation channel. *Plant Physiol. Biochem.* **41**: 945–954.
- Huang, L., Berkelman, T., Franklin, A.E., and Hoffman, N.E.** (1993). Characterization of a gene encoding a  $\text{Ca}^{2+}$ -ATPase-like protein in the plastid envelope. *Proc. Natl. Acad. Sci. USA* **90**: 10066–10070.

- Hunter, T. (1995). Protein kinases and phosphatases: the yin and yang of protein phosphorylation and signaling. *Cell* **80**: 225–236.
- Hwang, I., Sze, H., and Harper, J.F. (2000). A calcium-dependent protein kinase can inhibit a calmodulin-stimulated  $\text{Ca}^{2+}$  pump (ACA2) located in the endoplasmic reticulum of *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **97**: 6224–6229.
- Ikura, M., Osawa, M., and Ames, J.B. (2002). The role of calcium-binding proteins in the control of transcription: Structure to function. *Bioessays* **24**: 625–636.
- Irving, H.R., Gehring, C.A., and Parish, R.W. (1992). Changes in cytosolic pH and calcium of guard cells precede stomatal movements. *Proc. Natl. Acad. Sci. USA* **89**: 1790–1794.
- Ishibashi, K., Suzuki, M., and Imai, M. (2000). Molecular cloning of a novel form (two-repeat) protein related to voltage-gated sodium and calcium channels. *Biochem. Biophys. Res. Commun.* **270**: 370–376.
- Ishida, S., Yuasa, T., Nakata, M., and Takahashi, Y. (2008). A tobacco calcium-dependent protein kinase, CDPK1, regulates the transcription factor REPRESSION OF SHOOT GROWTH in response to gibberellins. *Plant Cell* **20**: 3273–3288.
- Ivashikina, N., and Hedrich, R. (2005).  $\text{K}^+$  currents through SV-type vacuolar channels are sensitive to elevated luminal sodium levels. *Plant J.* **41**: 606–614.
- Ivashuta, S., Liu, J., Lohar, D.P., Haridas, S., Bucciarelli, B., VandenBosch, K.A., Vance, C.P., Harrison, M.J., and Gantt, J.S. (2005). RNA interference identifies a calcium-dependent protein kinase involved in *Medicago truncatula* root development. *Plant Cell* **17**: 2911–2921.
- Jia, X.-Y., He, L.-H., Jing, R.-L., and Li, R.-Z. (2009). Calreticulin: Conserved protein and diverse functions in plants. *Physiol. Plant.* **136**: 127–138.
- Johannes, E., Brosnan, J.M., and Sanders, D. (1992). Parallel pathways for intracellular  $\text{Ca}^{2+}$  release from the vacuole of higher plants. *Plant J.* **2**: 97–102.
- Kadota, Y., Furuichi, T., Ogasawara, Y., Goh, T., Higashi, K., Muto, S., and Kuchitsu, K. (2004). Identification of putative voltage-dependent  $\text{Ca}^{2+}$ -permeable channels involved in cryptogin-induced  $\text{Ca}^{2+}$  transients and defense responses in tobacco BY-2 cells. *Biochem. Biophys. Res. Commun.* **317**: 823–830.
- Kang, J., Mehta, S., and Turano, F.J. (2004). The putative glutamate receptor 1.1 (AtGLR1.1) in *Arabidopsis thaliana* regulates abscisic acid biosynthesis and signaling to control development and water loss. *Plant Cell Physiol.* **45**: 1380–1389.
- Kang, J., and Turano, F.J. (2003). The putative glutamate receptor 1.1 (AtGLR1.1) functions as a regulator of carbon and nitrogen metabolism in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **100**: 6872–6877.
- Kaplan, B., Davydov, O., Knight, H., Galon, Y., Knight, M.R., Fluhr, R., and Fromm, H. (2006). Rapid transcriptome changes induced by cytosolic  $\text{Ca}^{2+}$  transients reveal ABRE-related sequences as  $\text{Ca}^{2+}$ -responsive cis elements in *Arabidopsis*. *Plant Cell* **18**: 2733–2748.
- Kasai, M., and Muto, S. (1990).  $\text{Ca}^{2+}$  pump and  $\text{Ca}^{2+}/\text{H}^+$  antiporter in plasma membrane vesicles isolated by aqueous two-phase partitioning from corn leaves. *J. Membr. Biol.* **114**: 133–142.
- Kiegle, E., Moore, C.A., Haseloff, J., Tester, M.A., and Knight, M.R. (2000). Cell-type-specific calcium responses to drought, salt and cold in the *Arabidopsis* root. *Plant J.* **23**: 267–278.
- Kim, B.G., Waadt, R., Cheong, Y.H., Pandey, G.K., Dominguez-Solis, J.R., Schultke, S., Lee, S.C., Kudla, J., and Luan, S. (2007). The calcium sensor CBL10 mediates salt tolerance by regulating ion homeostasis in *Arabidopsis*. *Plant J.* **52**: 473–484.
- Kim, S.A., Kwak, J.M., Jae, S.K., Wang, M.H., and Nam, H.G. (2001). Overexpression of the *AtGluR2* gene encoding an *Arabidopsis* homolog of mammalian glutamate receptors impairs calcium utilization and sensitivity to ionic stress in transgenic plants. *Plant Cell Physiol.* **42**: 74–84.
- Kim, M.C., Chung, W.S., Yun, D.J., and Cho, M.J. (2009). Calcium and calmodulin-mediated regulation of gene expression in plants. *Mol. Plant* **2**: 13–21.
- Klusener, B., Young, J.J., Murata, Y., Allen, G.J., Mori, I.C., Hugouvieux, V., and Schroeder, J.I. (2002). Convergence of calcium signaling pathways of pathogenic elicitors and abscisic acid in *Arabidopsis* guard cells. *Plant Physiol.* **130**: 2152–2163.
- Knight, H., and Knight, M.R. (2001). Abiotic stress signalling pathways: Specificity and cross-talk. *Trends Plant Sci.* **6**: 262–267.
- Kobayashi, M., Ohura, I., Kawakita, K., Yokota, N., Fujiwara, M., Shimamoto, K., Doke, N., and Yoshioka, H. (2007). Calcium-dependent protein kinases regulate the production of reactive oxygen species by potato NADPH oxidase. *Plant Cell* **19**: 1065–1080.
- Kolkisaoglu, U., Weinl, S., Blazevic, D., Batistić, O., and Kudla, J. (2004). Calcium sensors and their interacting protein kinases: genomics of the *Arabidopsis* and rice CBL-CIPK signaling networks. *Plant Physiol.* **134**: 43–58.
- Koren'kov, V., Park, S., Cheng, N.H., Sreevidya, C., Lachmansingh, J., Morris, J., Hirschi, K., and Wagner, G.J. (2007). Enhanced  $\text{Cd}^{2+}$ -selective root-tonoplast-transport in tobaccos expressing *Arabidopsis* cation exchangers. *Planta* **225**: 403–411.
- Kosuta, S., Hazledine, S., Sun, J., Miwa, H., Morris, R.J., Downie, J.A., and Oldroyd, G.E. (2008). Differential and chaotic calcium signatures in the symbiosis signaling pathway of legumes. *Proc. Natl. Acad. Sci. USA* **105**: 9823–9828.
- Kreimer, G., Melkonian, M., Holtum, J.A., and Lutzko, E. (1988). Stromal free calcium concentration and light-mediated activation of chloroplast fructose-1,6-bisphosphatase. *Plant Physiol.* **86**: 423–428.
- Kudla, J., Xu, Q., Harter, K., Grisse, W., and Luan, S. (1999). Genes for calcineurin B-like proteins in *Arabidopsis* are differentially regulated by stress signals. *Proc. Natl. Acad. Sci. USA* **96**: 4718–4723.
- Kurusu, T., Yagala, T., Miyao, A., Hirochika, H., and Kuchitsu, K. (2005). Identification of a putative voltage-gated  $\text{Ca}^{2+}$  channel as a key regulator of elicitor-induced hypersensitive cell death and mitogen-activated protein kinase activation in rice. *Plant J.* **42**: 798–809.
- Kushwaha, R., Singh, A., and Chattopadhyay, S. (2008). Calmodulin7 plays an important role as transcriptional regulator in *Arabidopsis* seedling development. *Plant Cell* **20**: 1747–1759.
- Lam, H.M., Chiu, J., Hsieh, M.H., Meisel, L., Oliveira, I.C., Shin, M., and Coruzzi, G. (1998). Glutamate-receptor genes in plants. *Nature* **396**: 125–126.
- Lamotte, O., Gould, K., Lecourieux, D., Sequeira-Legrand, A., Lebrun-Garcia, A., Durner, J., Pugin, A., and Wendehenne, D. (2004). Analysis of nitric oxide signaling functions in tobacco cells challenged by the elicitor cryptogin. *Plant Physiol.* **135**: 516–529.
- Laohavisit, A., Mortimer, J.C., Demidchik, V., Coxon, K.M., Stancombe, M.A., Macpherson, N., Brownlee, C., Hofmann, A., Webb, A.A., Miedema, H., Battey, N.H., and Davies, J.M. (2009). *Zea mays* annexins modulate cytosolic free  $\text{Ca}^{2+}$  and generate a  $\text{Ca}^{2+}$ -permeable conductance. *Plant Cell* **21**: 479–493.
- Lecourieux, D., Lamotte, O., Bourque, S., Wendehenne, D., Mazars, C., Ranjeva, R., and Pugin, A. (2005). Proteinaceous and oligosaccharidic elicitors induce different calcium signatures in the nucleus of tobacco cells. *Cell Calcium* **38**: 527–538.
- Lee, K.W., Chen, P.W., Lu, C.A., Chen, S., Ho, T.H., and Yu, S.M. (2009). Coordinated responses to oxygen and sugar deficiency allow rice seedlings to tolerate flooding. *Sci. Signal.* **2**: ra61.
- Lee, S.C., Lan, W.Z., Kim, B.G., Li, L., Cheong, Y.H., Pandey, G.K., Lu, G., Buchanan, B.B., and Luan, S. (2007a). A protein phosphorylation/dephosphorylation network regulates a plant potassium channel. *Proc. Natl. Acad. Sci. USA* **104**: 15959–15964.

- Lee, S.M., Kim, H.S., Han, H.J., Moon, B.C., Kim, C.Y., Harper, J.F., and Chung, W.S. (2007b). Identification of a calmodulin-regulated autoinhibited  $\text{Ca}^{2+}$ -ATPase (ACA11) that is localized to vacuole membranes in *Arabidopsis*. *FEBS Lett.* **581**: 3943–3949.
- Lemtiri-Chlieh, F., MacRobbie, E.A., Webb, A.A., Manison, N.F., Brownlee, C., Skepper, J.N., Chen, J., Prestwich, G.D., and Brearley, C.A. (2003). Inositol hexakisphosphate mobilizes an endomembrane store of calcium in guard cells. *Proc. Natl. Acad. Sci. USA* **100**: 10091–10095.
- Levchenko, V., Guinot, D.R., Klein, M., Roelfsema, M.R., Hedrich, R., and Dietrich, P. (2008). Stringent control of cytoplasmic  $\text{Ca}^{2+}$  in guard cells of intact plants compared to their counterparts in epidermal strips or guard cell protoplasts. *Protoplasma* **233**: 61–72.
- Li, J., Zhu, S., Song, X., Shen, Y., Chen, H., Yu, J., Yi, K., Liu, Y., Karplus, V.J., Wu, P., and Deng, X.W. (2006a). A rice glutamate receptor-like gene is critical for the division and survival of individual cells in the root apical meristem. *Plant Cell* **18**: 340–349.
- Li, L., Kim, B.G., Cheong, Y.H., Pandey, G.K., and Luan, S. (2006b). A  $\text{Ca}^{2+}$  signaling pathway regulates a  $\text{K}^{+}$  channel for low-K response in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **103**: 12625–12630.
- Li, X., Chanroj, S., Wu, Z., Romanowsky, S.M., Harper, J.F., and Sze, H. (2008). A distinct endosomal  $\text{Ca}^{2+}/\text{Mn}^{2+}$  pump affects root growth through the secretory process. *Plant Physiol.* **147**: 1675–1689.
- Liang, F., Cunningham, K.W., Harper, J.F., and Sze, H. (1997). ECA1 complements yeast mutants defective in  $\text{Ca}^{2+}$  pumps and encodes an endoplasmic reticulum-type  $\text{Ca}^{2+}$ -ATPase in *Arabidopsis thaliana*. *Proc. Natl. Acad. Sci. USA* **94**: 8579–8584.
- Lin, H., Yang, Y., Quan, R., Mendoza, I., Wu, Y., Du, W., Zhao, S., Schumaker, K.S., Pardo, J.M., and Guo, Y. (2009). Phosphorylation of SOS3-LIKE CALCIUM BINDING PROTEIN8 by SOS2 protein kinase stabilizes their protein complex and regulates salt tolerance in *Arabidopsis*. *Plant Cell* **21**: 1607–1619.
- Liu, J., and Zhu, J.K. (1998). A calcium sensor homolog required for plant salt tolerance. *Science* **280**: 1943–1945.
- Luan, S. (2009). The CBL-CIPK network in plant calcium signaling. *Trends Plant Sci.* **14**: 37–42.
- Luan, S., Kudla, J., Rodriguez-Concepcion, M., Yalovsky, S., and Griesem, W. (2002). Calmodulins and calcineurin B-like proteins: Calcium sensors for specific signal response coupling in plants. *Plant Cell* **14** (suppl.): S389–S400.
- Luan, S., Lan, W., and Chul Lee, S. (2009). Potassium nutrition, sodium toxicity, and calcium signaling: connections through the CBL-CIPK network. *Curr. Opin. Plant Biol.* **12**: 339–346.
- Ludwig, A.A., Romeis, T., and Jones, J.D. (2004). CDPK-mediated signalling pathways: Specificity and cross-talk. *J. Exp. Bot.* **55**: 181–188.
- Ludwig, A.A., Saitoh, H., Felix, G., Freymark, G., Miersch, O., Wasternack, C., Boller, T., Jones, J.D., and Romeis, T. (2005). Ethylene-mediated cross-talk between calcium-dependent protein kinase and MAPK signaling controls stress responses in plants. *Proc. Natl. Acad. Sci. USA* **102**: 10736–10741.
- Luo, G.Z., Wang, H.W., Huang, J., Tian, A.G., Wang, Y.J., Zhang, J.S., and Chen, S.Y. (2005). A putative plasma membrane cation/proton antiporter from soybean confers salt tolerance in *Arabidopsis*. *Plant Mol. Biol.* **59**: 809–820.
- Ma, Y., Szostkiewicz, I., Korte, A., Moes, D., Yang, Y., Christmann, A., and Grill, E. (2009). Regulators of PP2C phosphatase activity function as abscisic acid sensors. *Science* **324**: 1064–1068.
- Magnan, F., Ranty, B., Charpentreau, M., Sotta, B., Galaud, J.P., and Aldon, D. (2008). Mutations in AtCML9, a calmodulin-like protein from *Arabidopsis thaliana*, alter plant responses to abiotic stress and abscisic acid. *Plant J.* **56**: 575–589.
- Mahajan, S., Sopory, S.K., and Tuteja, N. (2006). Cloning and characterization of CBL-CIPK signalling components from a legume (*Pisum sativum*). *FEBS J.* **273**: 907–925.
- Martinez, J., Feltl, T., Scanlon, C.H., Lumsden, P.J., and Machackova, I. (2000). Subcellular localization of a high affinity binding site for D-myo-inositol 1,4,5-trisphosphate from *Chenopodium rubrum*. *Plant Physiol.* **124**: 475–483.
- Maser, P., et al. (2001). Phylogenetic relationships within cation transporter families of *Arabidopsis*. *Plant Physiol.* **126**: 1646–1667.
- Masi, E., Ciszak, M., Stefano, G., Renna, L., Azzarello, E., Pandolfi, C., Mugnai, S., Baluska, F., Arecchi, F.T., and Mancuso, S. (2009). Spatiotemporal dynamics of the electrical network activity in the root apex. *Proc. Natl. Acad. Sci. USA* **106**: 4048–4053.
- Mazars, C., Bourque, S., Mithofer, A., Pugin, A., and Ranjeva, R. (2009). Calcium homeostasis in plant cell nuclei. *New Phytol.* **181**: 261–274.
- McAinsh, M.R., and Hetherington, A.M. (1998). Encoding specificity in  $\text{Ca}^{2+}$  signalling systems. *Trends Plant Sci.* **3**: 32–36.
- McAinsh, M.R., Webb, A., Taylor, J.E., and Hetherington, A.M. (1995). Stimulus-induced oscillations in guard cell cytosolic free calcium. *Plant Cell* **7**: 1207–1219.
- McCormack, E., Tsai, Y.C., and Braam, J. (2005). Handling calcium signaling: *Arabidopsis* CaMs and CMLs. *Trends Plant Sci.* **10**: 383–389.
- Mei, H., Zhao, J., Pittman, J.K., Lachmansingh, J., Park, S., and Hirschi, K.D. (2007). In planta regulation of the *Arabidopsis*  $\text{Ca}^{2+}/\text{H}^{+}$  antiporter CAX1. *J. Exp. Bot.* **58**: 3419–3427.
- Menteyne, A., Burdakov, A., Charpentier, G., Petersen, O.H., and Cancela, J.M. (2006). Generation of specific  $\text{Ca}^{2+}$  signals from  $\text{Ca}^{2+}$  stores and endocytosis by differential coupling to messengers. *Curr. Biol.* **16**: 1931–1937.
- Meyerhoff, O., Muller, K., Roelfsema, M.R., Latz, A., Lacombe, B., Hedrich, R., Dietrich, P., and Becker, D. (2005). *AtGLR3.4*, a glutamate receptor channel-like gene is sensitive to touch and cold. *Planta* **222**: 418–427.
- Miedema, H., Bothwell, J.H., Brownlee, C., and Davies, J.M. (2001). Calcium uptake by plant cells—Channels and pumps acting in concert. *Trends Plant Sci.* **6**: 514–519.
- Miedema, H., Demidchik, V., Very, A.A., Bothwell, J.H., Brownlee, C., and Davies, J.M. (2008). Two voltage-dependent calcium channels co-exist in the apical plasma membrane of *Arabidopsis thaliana* root hairs. *New Phytol.* **179**: 378–385.
- Mills, R.F., Doherty, M.L., Lopez-Marques, R.L., Weimar, T., Dupree, P., Palmgren, M.G., Pittman, J.K., and Williams, L.E. (2008). ECA3, a Golgi-localized P2A-type ATPase, plays a crucial role in manganese nutrition in *Arabidopsis*. *Plant Physiol.* **146**: 116–128.
- Mito, N., Wimmers, L.E., and Bennett, A.B. (1996). Sugar regulates mRNA abundance of  $\text{H}^{+}$ -ATPase gene family members in tomato. *Plant Physiol.* **112**: 1229–1236.
- Miwa, H., Sun, J., Oldroyd, G.E., and Downie, J.A. (2006). Analysis of calcium spiking using aameleon calcium sensor reveals that nodulation gene expression is regulated by calcium spike number and the developmental status of the cell. *Plant J.* **48**: 883–894.
- Monshausen, G.B., Bibikova, T.N., Messerli, M.A., Shi, C., and Gilroy, S. (2007). Oscillations in extracellular pH and reactive oxygen species modulate tip growth of *Arabidopsis* root hairs. *Proc. Natl. Acad. Sci. USA* **104**: 20996–21001.
- Monshausen, G.B., Bibikova, T.N., Weisenseel, M.H., and Gilroy, S. (2009).  $\text{Ca}^{2+}$  regulates reactive oxygen species production and pH during mechanosensing in *Arabidopsis* roots. *Plant Cell* **21**: 2341–2356.
- Monshausen, G.B., Messerli, M.A., and Gilroy, S. (2008). Imaging of the Yellow Cameleon 3.6 indicator reveals that elevations in cytosolic  $\text{Ca}^{2+}$  follow oscillating increases in growth in root hairs of *Arabidopsis*. *Plant Physiol.* **147**: 1690–1698.

- Moreno, I., Norambuena, L., Maturana, D., Toro, M., Vergara, C., Orellana, A., Zurita-Silva, A., and Ordenes, V.R. (2008). AtHMA1 is a thapsigargin-sensitive  $\text{Ca}^{2+}$ /heavy metal pump. *J. Biol. Chem.* **283**: 9633–9641.
- Mori, I.C., Murata, Y., Yang, Y., Munemasa, S., Wang, Y.-F., Andreoli, S., Tiriach, H., Alonso, J.M., Harper, J.F., Ecker, J.R., Kwak, J.M., and Schroeder, J.I. (2006). CDPKs CPK6 and CPK3 function in ABA regulation of guard cell S-type anion- and  $\text{Ca}^{2+}$ -permeable channels and stomatal closure. *PLoS Biol.* **4**: 1749–1762.
- Mortimer, J.C., Laohavisit, A., Macpherson, N., Webb, A., Brownlee, C., Battey, N.H., and Davies, J.M. (2008). Annexins: Multifunctional components of growth and adaptation. *J. Exp. Bot.* **59**: 533–544.
- Munnik, T., and Testerink, C. (2009). Plant phospholipid signaling: In a nutshell. *J. Lipid Res.* **50** (suppl.): S260–S265.
- Muir, S.R., and Sanders, D. (1996). Pharmacology of  $\text{Ca}^{2+}$  release from red beet microsomes suggests the presence of ryanodine receptor homologs in higher plants. *FEBS Lett.* **395**: 39–42.
- Muto, S., Izawa, S., and Miyachi, S. (1982). Light-induced  $\text{Ca}^{2+}$  uptake by intact chloroplast. *FEBS Lett.* **139**: 250–254.
- Nakagawa, Y., et al. (2007). *Arabidopsis* plasma membrane protein crucial for  $\text{Ca}^{2+}$  influx and touch sensing in roots. *Proc. Natl. Acad. Sci. USA* **104**: 3639–3644.
- Navazio, L., Bewell, M.A., Siddiqua, A., Dickinson, G.D., Galione, A., and Sanders, D. (2000). Calcium release from the endoplasmic reticulum of higher plants elicited by the NADP metabolite nicotinic acid adenine dinucleotide phosphate. *Proc. Natl. Acad. Sci. USA* **97**: 8693–8698.
- Negi, J., Matsuda, O., Nagasawa, T., Oba, Y., Takahashi, H., Kawai-Yamada, M., Uchimiya, H., Hashimoto, M., and Iba, K. (2008).  $\text{CO}_2$  regulator SLAC1 and its homologues are essential for anion homeostasis in plant cells. *Nature* **452**: 483–486.
- Neuhaus, G., Bowler, C., Kern, R., and Chua, N.H. (1993). Calcium/calmodulin-dependent and -independent phytochrome signal transduction pathways. *Cell* **73**: 937–952.
- Nobel, P.S., Murakami, S., and Takamiya, A. (1966). Localization of light-induced strontium accumulation in spinach chloroplasts. *Plant Cell Physiol.* **7**: 263–275.
- Nomura, H., Komori, T., Kobori, M., Nakahira, Y., and Shiina, T. (2008). Evidence for chloroplast control of external  $\text{Ca}^{2+}$ -induced cytosolic  $\text{Ca}^{2+}$  transients and stomatal closure. *Plant J.* **53**: 988–998.
- Ohta, M., Guo, Y., Halfter, U., and Zhu, J.K. (2003). A novel domain in the protein kinase SOS2 mediates interaction with the protein phosphatase 2C ABI2. *Proc. Natl. Acad. Sci. USA* **100**: 11771–11776.
- Pandey, G.K., Cheong, Y.H., Kim, K.N., Grant, J.J., Li, L., Hung, W., D'Angelo, C., Weinl, S., Kudla, J., and Luan, S. (2004). The calcium sensor calcineurin B-like 9 modulates abscisic acid sensitivity and biosynthesis in *Arabidopsis*. *Plant Cell* **16**: 1912–1924.
- Pandey G.K., Grant, J.J., Cheong, Y.H., Kim, B.G., Li le, G., and Luan, S. (2008). Calcineurin-B-like protein CBL9 interacts with target kinase CIPK3 in the regulation of ABA response in seed germination. *Mol. Plant* **1**: 238–248.
- Park, C.Y., Lee, J.H., Yoo, J.H., Moon, B.C., Choi, M.S., Kang, Y.H., Lee, S.M., Kim, H.S., Kang, K.Y., Chung, W.S., Lim, C.O., and Cho, M.J. (2005). WRKY group IId transcription factors interact with calmodulin. *FEBS Lett.* **579**: 1545–1550.
- Park, S.Y., et al. (2009). Abscisic acid inhibits type 2C protein phosphatases via the PYR/PYL family of START proteins. *Science* **324**: 1068–1071.
- Pei, Z.M., Murata, Y., Benning, G., Thomine, S., Klusener, B., Allen, G.J., Grill, E., and Schroeder, J.I. (2000). Calcium channels activated by hydrogen peroxide mediate abscisic acid signalling in guard cells. *Nature* **406**: 731–734.
- Peiter, E., Maathuis, F.J., Mills, L.N., Knight, H., Pelloux, J., Hetherington, A.M., and Sanders, D. (2005). The vacuolar  $\text{Ca}^{2+}$ -activated channel TPC1 regulates germination and stomatal movement. *Nature* **434**: 404–408.
- Perez, V., Wherrett, T., Shabala, S., Muniz, J., Dobrovinskaya, O., and Pottosin, I. (2008). Homeostatic control of slow vacuolar channels by luminal cations and evaluation of the channel-mediated tonoplast  $\text{Ca}^{2+}$  fluxes in situ. *J. Exp. Bot.* **59**: 3845–3855.
- Persson, S., Wyatt, S.E., Love, J., Thompson, W.F., Robertson, D., and Boss, W.F. (2001). The  $\text{Ca}^{2+}$  status of the endoplasmic reticulum is altered by induction of calreticulin expression in transgenic plants. *Plant Physiol.* **126**: 1092–1104.
- Picton, J.M., and Steer, M.W. (1985). The effects of ruthenium red, lanthanum, fluorescein isothiocyanate and trifluoperazine on vesicle transport, vesicle fusion and tip extension in pollen tubes. *Planta* **163**: 20–26.
- Pittman, J.K., Shigaki, T., Cheng, N.H., and Hirschi, K.D. (2002). Mechanism of N-terminal autoinhibition in the *Arabidopsis*  $\text{Ca}^{2+}/\text{H}^+$  antiporter CAX1. *J. Biol. Chem.* **277**: 26452–26459.
- Plieth, C., Hansen, U.P., Knight, H., and Knight, M.R. (1999). Temperature sensing by plants: The primary characteristics of signal perception and calcium response. *Plant J.* **18**: 491–497.
- Portis, A.R., Jr., and Heldt, H.W. (1976). Light-dependent changes of the  $\text{Mg}^{2+}$  concentration in the stroma in relation to the  $\text{Mg}^{2+}$  dependency of  $\text{CO}_2$  fixation in intact chloroplasts. *Biochim. Biophys. Acta* **449**: 434–436.
- Pottosin, I.I., Martinez-Estevéz, M., Dobrovinskaya, O.R., and Muniz, J. (2005). Regulation of the slow vacuolar channel by luminal potassium: Role of surface charge. *J. Membr. Biol.* **205**: 103–111.
- Pottosin, I.I., and Schonknecht, G. (2007). Vacuolar calcium channels. *J. Exp. Bot.* **58**: 1559–1569.
- Qi, Z., Stephens, N.R., and Spalding, E.P. (2006). Calcium entry mediated by GLR3.3, an *Arabidopsis* glutamate receptor with a broad agonist profile. *Plant Physiol.* **142**: 963–971.
- Qiu, Q.S., Guo, Y., Dietrich, M.A., Schumaker, K.S., and Zhu, J.K. (2002). Regulation of SOS1, a plasma membrane  $\text{Na}^+/\text{H}^+$  exchanger in *Arabidopsis thaliana*, by SOS2 and SOS3. *Proc. Natl. Acad. Sci. USA* **99**: 8436–8441.
- Quan, R., Lin, H., Mendoza, I., Zhang, Y., Cao, W., Yang, Y., Shang, M., Chen, S., Pardo, J.M., and Guo, Y. (2007). SCABP8/CBL10, a putative calcium sensor, interacts with the protein kinase SOS2 to protect *Arabidopsis* shoots from salt stress. *Plant Cell* **19**: 1415–1431.
- Qudeimat, E., Faltusz, A.M., Wheeler, G., Lang, D., Brownlee, C., Reski, R., and Frank, W. (2008). A PIIB-type  $\text{Ca}^{2+}$ -ATPase is essential for stress adaptation in *Physcomitrella patens*. *Proc. Natl. Acad. Sci. USA* **105**: 19555–19560.
- Ranf, S., Wunnenberg, P., Lee, J., Becker, D., Dunkel, M., Hedrich, R., Scheel, D., and Dietrich, P. (2008). Loss of the vacuolar cation channel, AtTPC1, does not impair  $\text{Ca}^{2+}$  signals induced by abiotic and biotic stresses. *Plant J.* **53**: 287–299.
- Richter, G.L., Monshausen, G.B., Krol, A., and Gilroy, S. (2009). Mechanical stimuli modulate lateral root organogenesis. *Plant Physiol.* **151**: 1855–1866.
- Romeis, T., Ludwig, A.A., Martin, R., and Jones, J.D. (2001). Calcium-dependent protein kinases play an essential role in a plant defence response. *EMBO J.* **20**: 5556–5567.
- Rudd, J.J., and Franklin-Tong, V.E. (1999). Calcium signaling in plants. *Cell. Mol. Life Sci.* **55**: 214–232.
- Sai, J., and Johnson, C.H. (2002). Dark-stimulated calcium ion fluxes in the chloroplast stroma and cytosol. *Plant Cell* **14**: 1279–1291.
- Saidi, Y., Finka, A., Muriset, M., Bromberg, Z., Weiss, Y.G., Maathuis, F.J., and Goloubinoff, P. (2009). The heat shock response in moss plants is regulated by specific calcium-permeable channels in the plasma membrane. *Plant Cell* **21**: 2829–2843.

- Sanchez-Barrena, M.J., Fujii, H., Angulo, I., Martinez-Ripoll, M., Zhu, J.K., and Albert, A.** (2007). The structure of the C-terminal domain of the protein kinase AtSOS2 bound to the calcium sensor AtSOS3. *Mol. Cell* **26**: 427–435.
- Sanders, D., Brownlee, C., and Harper, J.F.** (1999). Communicating with calcium. *Plant Cell* **11**: 691–706.
- Sanders, D., Pelloux, J., Brownlee, C., and Harper, J.F.** (2002). Calcium at the crossroads of signaling. *Plant Cell* **14** (suppl.): S401–S417.
- Schiott, M., Romanowsky, S.M., Baekgaard, L., Jakobsen, M.K., Palmgren, M.G., and Harper, J.F.** (2004). A plant plasma membrane  $\text{Ca}^{2+}$  pump is required for normal pollen tube growth and fertilization. *Proc. Natl. Acad. Sci. USA* **101**: 9502–9507.
- Schroeder, J.I., Kwak, J.M., and Allen, G.J.** (2001). Guard cell abscisic acid signalling and engineering drought hardness in plants. *Nature* **410**: 327–330.
- Schumaker, K.S., and Sze, H.** (1987). Inositol 1,4,5-trisphosphate releases  $\text{Ca}^{2+}$  from vacuolar membrane vesicles of oat roots. *J. Biol. Chem.* **262**: 3944–3946.
- Scrase-Field, S.A., and Knight, M.R.** (2003). Calcium: Just a chemical switch? *Curr. Opin. Plant Biol.* **6**: 500–506.
- Seigneurin-Berry, D., Gravot, A., Auroy, P., Mazard, C., Kraut, A., Finazzi, G., Grunwald, D., Rappaport, F., Vavasseur, A., Joyard, J., Richaud, P., and Rolland, N.** (2006). HMA1, a new Cu-ATPase of the chloroplast envelope, is essential for growth under adverse light conditions. *J. Biol. Chem.* **281**: 2882–2892.
- Shaw, S.L., and Long, S.R.** (2003). Nod factor elicits two separable calcium responses in *Medicago truncatula* root hair cells. *Plant Physiol.* **131**: 976–984.
- Shi, J., Kim, K.N., Ritz, O., Albrecht, V., Gupta, R., Harter, K., Luan, S., and Kudla, J.** (1999). Novel protein kinases associated with calcineurin B-like calcium sensors in *Arabidopsis*. *Plant Cell* **11**: 2393–2405.
- Shigaki, T., Rees, I., Nakhleh, L., and Hirschi, K.D.** (2006). Identification of three distinct phylogenetic groups of CAX cation/proton antiporters. *J. Mol. Evol.* **63**: 815–825.
- Siegel, R.S., Xue, S., Murata, Y., Yang, Y., Nishimura, N., Wang, A., and Schroeder, J.I.** (2009). Calcium elevation-dependent and attenuated resting calcium-dependent abscisic acid induction of stomatal closure and abscisic acid-induced enhancement of calcium sensitivities of S-type anion and inward-rectifying K channels in *Arabidopsis* guard cells. *Plant J.* **59**: 207–220.
- Sivaguru, M., Pike, S., Gassmann, W., and Baskin, T.I.** (2003). Aluminum rapidly depolymerizes cortical microtubules and depolarizes the plasma membrane: evidence that these responses are mediated by a glutamate receptor. *Plant Cell Physiol.* **44**: 667–675.
- Snowman, B.N., Kovar, D.R., Shevchenko, G., Franklin-Tong, V.E., and Staiger, C.J.** (2002). Signal-mediated depolymerization of actin in pollen during the self-incompatibility response. *Plant Cell* **14**: 2613–2626.
- Soderling, T.R.** (1999). The  $\text{Ca}^{2+}$ -calmodulin-dependent protein kinase cascade. *Trends Biochem. Sci.* **24**: 232–236.
- Stephens, N.R., Qi, Z., and Spalding, E.P.** (2008). Glutamate receptor subtypes evidenced by differences in desensitization and dependence on the *GLR3.3* and *GLR3.4* genes. *Plant Physiol.* **146**: 529–538.
- Sze, H., Liang, F., Hwang, I., Curran, A.C., and Harper, J.F.** (2000). Diversity and regulation of plant  $\text{Ca}^{2+}$  pumps: Insights from expression in yeast. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **51**: 433–462.
- Takeda, S., Gapper, C., Kaya, H., Bell, E., Kuchitsu, K., and Dolan, L.** (2008). Local positive feedback regulation determines cell shape in root hair cells. *Science* **319**: 1241–1244.
- Thion, L., Mazars, C., Nacry, P., Bouchez, D., Moreau, M., Ranjeva, R., and Thuleau, P.** (1998). Plasma membrane depolarization-activated calcium channels, stimulated by microtubule-depolymerizing drugs in wild-type *Arabidopsis thaliana* protoplasts, display constitutively large activities and a longer half-life in *ton 2* mutant cells affected in the organization of cortical microtubules. *Plant J.* **13**: 603–610.
- Thuleau, P., Schroeder, J.I., and Ranjeva, R.** (1998). Recent advances in the regulation of plant calcium channels: Evidence for regulation by G-proteins, the cytoskeleton and second messengers. *Curr. Opin. Plant Biol.* **1**: 424–427.
- Tracy, F.E., Gilliam, M., Dodd, A.N., Webb, A.A., and Tester, M.** (2008). NaCl-induced changes in cytosolic free  $\text{Ca}^{2+}$  in *Arabidopsis thaliana* are heterogeneous and modified by external ionic composition. *Plant Cell Environ.* **31**: 1063–1073.
- Tsai, Y.C., Delk, N.A., Chowdhury, N.I., and Braam, J.** (2007). *Arabidopsis* potential calcium sensors regulate nitric oxide levels and the transition to flowering. *Plant Signal. Behav.* **2**: 446–454.
- Umezawa, T., Sugiyama, N., Mizoguchi, M., Hayashi, S., Myouga, F., Yamaguchi-Shinozaki, K., Ishihama, Y., Hirayama, T., and Shinozaki, K.** (2009). Type 2C protein phosphatases directly regulate abscisic acid-activated protein kinases in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **106**: 17588–17593.
- Urhuhart, W., Gunawardena, A.H., Moeder, W., Ali, R., Berkowitz, G.A., and Yoshioka, K.** (2007). The chimeric cyclic nucleotide-gated ion channel ATCNGC11/12 constitutively induces programmed cell death in a  $\text{Ca}^{2+}$  dependent manner. *Plant Mol. Biol.* **65**: 747–761.
- Vahisalu, T., Kollist, H., Wang, Y.F., Nishimura, N., Chan, W.Y., Valerio, G., Lamminmaki, A., Brosche, M., Moldau, H., Desikan, R., Schroeder, J.I., and Kangasjarvi, J.** (2008). SLAC1 is required for plant guard cell S-type anion channel function in stomatal signalling. *Nature* **452**: 487–491.
- Vainonen, J.P., Hansson, M., and Vener, A.V.** (2005). STN8 protein kinase in *Arabidopsis thaliana* is specific in phosphorylation of photosystem II core proteins. *J. Biol. Chem.* **280**: 33679–33686.
- Vainonen, J.P., Sakuragi, Y., Stael, S., Tikkanen, M., Allahverdiyeva, Y., Paakkari, V., Aro, E., Suorsa, M., Scheller, H.V., Vener, A.V., and Aro, E.M.** (2008). Light regulation of CaS, a novel phosphoprotein in the thylakoid membrane of *Arabidopsis thaliana*. *FEBS J.* **275**: 1767–1777.
- Waadt, R., Schmidt, L.K., Lohse, M., Hashimoto, K., Bock, R., and Kudla, J.** (2008). Multicolor bimolecular fluorescence complementation reveals simultaneous formation of alternative CBL/CIPK complexes in planta. *Plant J.* **56**: 505–516.
- Walch-Liu, P., Liu, L.H., Remans, T., Tester, M., and Forde, B.G.** (2006). Evidence that L-glutamate can act as an exogenous signal to modulate root growth and branching in *Arabidopsis thaliana*. *Plant Cell Physiol.* **47**: 1045–1057.
- Webb, A.A.R., McAinsh, M.R., Taylor, J.E., and Hetherington, A.M.** (1996). Calcium ions as intracellular second messengers in higher plants. *Adv. Bot. Res.* **22**: 45–96.
- Weinl, S., Held, K., Schlucking, K., Steinhorst, L., Kuhlert, S., Hippler, M., and Kudla, J.** (2008). A plastid protein crucial for  $\text{Ca}^{2+}$ -regulated stomatal responses. *New Phytol.* **179**: 675–686.
- Weinl, S., and Kudla, J.** (2009). The CBL-CIPK  $\text{Ca}^{2+}$ -decoding signaling network: Function and perspectives. *New Phytol.* **184**: 517–528.
- Wheeler, G.L., and Brownlee, C.** (2008).  $\text{Ca}^{2+}$  signalling in plants and green algae-changing channels. *Trends Plant Sci.* **13**: 506–514.
- White, P.J., Bowen, H.C., Demidchik, V., Nichols, C., and Davies, J.M.** (2002). Genes for calcium-permeable channels in the plasma membrane of plant root cells. *Biochim. Biophys. Acta* **1564**: 299–309.
- White, P.J., and Broadley, M.R.** (2003). Calcium in plants. *Ann. Bot. (Lond.)* **92**: 487–511.
- Williamson, R.E., and Ashley, C.C.** (1982). Free  $\text{Ca}^{2+}$  and cytoplasmic streaming in the alga *Chara*. *Nature* **296**: 647–650.

- Wimmers, L.E., Ewing, N.N., and Bennett, A.B.** (1992). Higher plant  $\text{Ca}^{2+}$ -ATPase: Primary structure and regulation of mRNA abundance by salt. *Proc. Natl. Acad. Sci. USA* **89**: 9205–9209.
- Xu, J., Li, H.D., Chen, L.Q., Wang, Y., Liu, L.L., He, L., and Wu, W.H.** (2006). A protein kinase, interacting with two calcineurin B-like proteins, regulates  $\text{K}^+$  transporter AKT1 in *Arabidopsis*. *Cell* **125**: 1347–1360.
- Yoo, J.H., et al.** (2005). Direct interaction of a divergent CaM isoform and the transcription factor, MYB2, enhances salt tolerance in *Arabidopsis*. *J. Biol. Chem.* **280**: 3697–3706.
- Yoon, G.M., Dowd, P.E., Gilroy, S., and McCubbin, A.G.** (2006). Calcium-dependent protein kinase isoforms in *Petunia* have distinct functions in pollen tube growth, including regulating polarity. *Plant Cell* **18**: 867–878.
- Yoshioka, K., Moeder, W., Kang, H.G., Kachroo, P., Masmoudi, K., Berkowitz, G., and Klessig, D.F.** (2006). The chimeric *Arabidopsis* CYCLIC NUCLEOTIDE-GATED ION CHANNEL11/12 activates multiple pathogen resistance responses. *Plant Cell* **18**: 747–763.
- Young, J.J., Mehta, S., Israelsson, M., Godoski, J., Grill, E., and Schroeder, J.I.** (2006).  $\text{CO}_2$  signaling in guard cells: Calcium sensitivity response modulation, a  $\text{Ca}^{2+}$ -independent phase, and  $\text{CO}_2$  insensitivity of the *gca2* mutant. *Proc. Natl. Acad. Sci. USA* **103**: 7506–7511.
- Yu, I.C., Parker, J., and Bent, A.F.** (1998). Gene-for-gene disease resistance without the hypersensitive response in *Arabidopsis dnd1* mutant. *Proc. Natl. Acad. Sci. USA* **95**: 7819–7824.
- Zhang, W., Zhou, R.G., Gao, Y.J., Zheng, S.Z., Xu, P., Zhang, S.Q., and Sun, D.Y.** (2009). Molecular and genetic evidence for the key role of AtCaM3 in heat-shock signal transduction in *Arabidopsis*. *Plant Physiol.* **149**: 1773–1784.
- Zhao, J., Barkla, B.J., Marshall, J., Pittman, J.K., and Hirschi, K.D.** (2008). The *Arabidopsis cax3* mutants display altered salt tolerance, pH sensitivity and reduced plasma membrane  $\text{H}^+$ -ATPase activity. *Planta* **227**: 659–669.
- Zhao, J., Shigaki, T., Mei, H., Guo, Y.Q., Cheng, N.H., and Hirschi, K.D.** (2009). Interaction between *Arabidopsis*  $\text{Ca}^{2+}/\text{H}^+$  Exchangers CAX1 and CAX3. *J. Biol. Chem.* **284**: 4605–4615.
- Zhu, S.Y., et al.** (2007). Two calcium-dependent protein kinases, CPK4 and CPK11, regulate abscisic acid signal transduction in *Arabidopsis*. *Plant Cell* **19**: 3019–3036.
- Zocchi, G., and Rabotti, G.** (1993). Calcium transport in membrane vesicles isolated from maize coleoptiles (effect of indoleacetic acid and fusicoccin). *Plant Physiol.* **101**: 135–139.

## Calcium Signals: The Lead Currency of Plant Information Processing

Jörg Kudla, Oliver Batistic and Kenji Hashimoto

*Plant Cell* 2010;22;541-563; originally published online March 30, 2010;

DOI 10.1105/tpc.109.072686

This information is current as of January 20, 2018

<b>References</b>	This article cites 257 articles, 122 of which can be accessed free at: <a href="/content/22/3/541.full.html#ref-list-1">/content/22/3/541.full.html#ref-list-1</a>
<b>Permissions</b>	<a href="https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&amp;iissn=1532298X&amp;WT.mc_id=pd_hw1532298X">https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&amp;iissn=1532298X&amp;WT.mc_id=pd_hw1532298X</a>
<b>eTOCs</b>	Sign up for eTOCs at: <a href="http://www.plantcell.org/cgi/alerts/ctmain">http://www.plantcell.org/cgi/alerts/ctmain</a>
<b>CiteTrack Alerts</b>	Sign up for CiteTrack Alerts at: <a href="http://www.plantcell.org/cgi/alerts/ctmain">http://www.plantcell.org/cgi/alerts/ctmain</a>
<b>Subscription Information</b>	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: <a href="http://www.aspb.org/publications/subscriptions.cfm">http://www.aspb.org/publications/subscriptions.cfm</a>