

# The TOR Pathway Modulates the Structure of Cell Walls in *Arabidopsis*<sup>W</sup>

Ruth-Maria Leiber,<sup>a,1,2</sup> Florian John,<sup>a,1</sup> Yves Verherbruggen,<sup>b,3</sup> Anouck Diet,<sup>a,4</sup> J. Paul Knox,<sup>b</sup> and Christoph Ringli<sup>a,5</sup>

<sup>a</sup> University of Zürich, Institute of Plant Biology, 8008 Zurich, Switzerland

<sup>b</sup> Centre for Plant Sciences, Faculty of Biological Sciences, University of Leeds, Leeds LS2 9JT, United Kingdom

Plant cell growth is limited by the extension of cell walls, which requires both the synthesis and rearrangement of cell wall components in a controlled fashion. The target of rapamycin (TOR) pathway is a major regulator of cell growth in eukaryotes, and inhibition of this pathway by rapamycin reduces cell growth. Here, we show that in plants, the TOR pathway affects cell wall structures. LRR-extensin1 (LRX1) of *Arabidopsis thaliana* is an extracellular protein involved in cell wall formation in root hairs, and *lrx1* mutants develop aberrant root hairs. *rol5* (for repressor of *lrx1*) was identified as a suppressor of *lrx1*. The functionally similar ROL5 homolog in yeast, Ncs6p (needs Cla4 to survive 6), was previously found to affect TOR signaling. Inhibition of TOR signaling by rapamycin led to suppression of the *lrx1* mutant phenotype and caused specific changes to galactan/rhamnogalacturonan-I and arabinogalactan protein components of cell walls that were similar to those observed in the *rol5* mutant. The ROL5 protein accumulates in mitochondria, a target of the TOR pathway and major source of reactive oxygen species (ROS), and *rol5* mutants show an altered response to ROS. This suggests that ROL5 might function as a mitochondrial component of the TOR pathway that influences the plant's response to ROS.

## INTRODUCTION

Plant cell growth is tightly linked to the expansion of the cell wall. Cell walls are complex structures that resist internal turgor pressure and, for cell enlargement to take place, have to incorporate new material and rearrange internal linkages between the different components (Martin et al., 2001). In dicotyledonous plants, the primary cell wall is composed of cellulose microfibrils that are interconnected by hemicelluloses, mainly xyloglucan. This is considered to be the load-bearing structure and is embedded in a matrix of pectic polysaccharides (Carpita and Gibeaut, 1993). The pectic matrix has three major components: homogalacturonan, rhamnogalacturonan-I (RGI), which contains side chains of galactan and arabinan, and rhamnogalacturonan-II. Pectins influence cell wall rigidity and strength as well as cell–cell adhesion. In addition, RGI regulates wall porosity, which in turn influences the mobility of cell wall–modifying proteins and, thus, cell wall expansion (Baron-Epel et al., 1988; Ridley et al., 2001;

Willats et al., 2001; McCartney et al., 2003). Structural cell wall proteins such as hydroxyproline-rich glycoproteins (HRGPs) influence the mechanical properties of cell walls but can also be involved in cell elongation and signaling as exemplified by arabinogalactan proteins (AGPs). These are GPI-anchored proteins of the HRGP family that are extensively glycosylated with arabinose and galactose (Ding and Zhu, 1997; Majewska-Sawka and Nothnagel, 2000; van Hengel and Roberts, 2002).

The structure of cell walls, which influences the cell walls' properties, is in a constant flow of remodeling as it adapts to the prevailing functional requirements. Therefore, plants have evolved a sensing system to monitor cell wall composition and to regulate cell wall modification and restructuring. These activities are likely to involve transmembrane or membrane-anchored proteins. Receptor-like kinases, such as THESEUS and wall-associated kinases, have been shown to sense and modify cell elongation (Kohorn et al., 2006; Hematy et al., 2007), as have lectins and GPI-anchored proteins, such as AGPs (Humphrey et al., 2007; Hematy and Höfte, 2008). LRR-extensins (LRXs) are extracellular proteins consisting of an N-terminal leucine-rich repeat domain and a C-terminal extensin domain typical of HRGPs (Baumberger et al., 2003a). This particular structure suggests that LRX proteins might have a signaling or regulatory function during cell wall development (Ringli, 2005). Indeed, *Arabidopsis thaliana* LRX1 is predominantly expressed in root hairs, and *lrx1* mutants develop defective cell walls resulting in aberrant root hair formation (Baumberger et al., 2001, 2003b).

The TOR (for target of rapamycin) pathway is a major growth regulator in eukaryotes that senses nutrient availability and growth stimulators, regulates the translational machinery, and modulates cell growth. The Ser/Thr kinase TOR is central to the TOR pathway and is inhibited by the specific inhibitor rapamycin,

<sup>1</sup> These authors contributed equally to this work.

<sup>2</sup> Current address: Tecan Switzerland, Seestrasse 103, 8708 Männedorf, Switzerland.

<sup>3</sup> Current address: Joint BioEnergy Institute, Lawrence Berkeley National Laboratory, 5885 Hollis St., Emeryville, CA 94608.

<sup>4</sup> Current address: Université Paris 7, Institut des Sciences Végétales, Centre National de la Recherche Scientifique, Avenue de la Terrasse, 91198 Gif-sur-Yvette, France.

<sup>5</sup> Address correspondence to chringli@botinst.uzh.ch.

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (www.plantcell.org) is: Christoph Ringli (chringli@botinst.uzh.ch).

<sup>W</sup> Online version contains Web-only data.

www.plantcell.org/cgi/doi/10.1105/tpc.109.073007

resulting in reduced cell growth. Rapamycin inhibits the TOR kinase by forming a ternary complex with the immunophilin protein FKBP12 (FK506 binding protein 12) and TOR (Huang et al., 2003; Wullschleger et al., 2006). An important function of the TOR pathway is the regulation of mitochondrial activity and, hence, the production of reactive oxygen species (ROS), which affect life span (Schieke and Finkel, 2006; Cunningham et al., 2007) and, in plants, have an impact on oxidative stress, cell wall extension, and cell growth (for review, see Gapper and Dolan, 2006; Rhoads et al., 2006).

Recent analyses in *Candida albicans* have provided evidence for the participation of the TOR pathway in cell wall integrity sensing in yeast (Tsao et al., 2009). Numerous components of the TOR pathway were identified in yeast based on rapamycin hypersensitivity of the corresponding mutants (Chan et al., 2000). Mutations in *NCS6* (*needs Cla4 to survive 6*) of yeast induce rapamycin hypersensitivity and influence cell growth under nutrient-limited conditions (Chan et al., 2000; Goehring et al., 2003a). Recently, *Ncs6p* has been shown to be important for the modification of cytoplasmic tRNAs. tRNAs are frequently modified, mostly at the wobble position (position 34) or next to and 3' of the anticodon (position 37). tRNAs specific for Glu, Glc, and Lys have a 2-thiouridine derivative as wobble nucleoside, which helps to effectively read the corresponding codons on the mRNAs (Björk et al., 2007). *Ncs6p* and homologous proteins in other organisms are involved in the thiolation of U34, and mutations in the corresponding genes lead to the absence of thiolation (Björk et al., 2007; Schlieker et al., 2008; Leidel et al., 2009). Even though mutating *nsc6* only affects cytoplasmic tRNAs (Noma et al., 2009), *Ncs6p* is also found in mitochondria (Huh et al., 2003). Dual localization of proteins in different compartments frequently has been observed (Krause and Krupinska, 2009). Thus, it remains to be shown whether the effect of the *nsc6* mutant on TOR signaling is an indirect effect induced by the lack of tRNA modification or a second activity of the protein, reflected by its presence in mitochondria.

The TOR pathway has also been identified in plants, and some of the proteins involved in this process have been characterized (Anderson et al., 2005; Deprost et al., 2005; Ingram and Waites, 2006; Mahfouz et al., 2006). While a *tor* knockout mutant in *Arabidopsis* is embryo-lethal, modified TOR expression strongly influences plant growth, emphasizing the importance of TOR during plant development (Menand et al., 2002; Deprost et al., 2007). *Arabidopsis* is not sensitive to the specific TOR inhibitor rapamycin as rapamycin cannot form the ternary complex with FKBP12 and TOR. However, expression of yeast *FKBP12* induces rapamycin sensitivity in *Arabidopsis* (Mahfouz et al., 2006; Sormani et al., 2007).


Here, we provide evidence for a role of the plant TOR pathway in modulating cell wall structures. A suppressor screen on the root hair cell wall formation mutant *lrx1* resulted in the identification of the *rol5* (*for repressor of lrx1*) mutant. The *rol5* mutation induced changes in cell wall structure that might be the basis of suppression of *lrx1*. *ROL5* is functionally similar to *Ncs6p*, which influences TOR signaling in yeast and is required for the modification of tRNAs in *Arabidopsis*. Interfering with TOR signaling by the addition of rapamycin in yeast *FKBP12*-expressing *lrx1* mutant plants relieved the *lrx1* root hair phenotype and induced

specific changes in cell wall structure similar to *rol5*. Together, these data indicate that interfering with TOR signaling induces changes in cell walls and provide evidence for a role of the TOR pathway in the regulation of cell wall structure and properties.

## RESULTS


### Identification of *rol5*, a Suppressor of the *lrx1* Root Hair Phenotype

As a result of the defective cell wall structure, *lrx1* mutants form root hairs that are short and deformed and frequently burst (Figures 1A and 1B). To identify new loci that are involved in regulating cell wall formation and structure, a suppressor screen was performed on the *lrx1* mutant. As described previously (Diet et al., 2006), an *lrx1* missense allele was used for ethyl methanesulfonate (EMS) mutagenesis, and M2 seedlings displaying a suppressed *lrx1* phenotype were isolated. The *rol5-1* mutant was identified in this screen as it suppressed the *lrx1* phenotype. *lrx1 rol5-1* double mutants developed root hairs that were comparable to those of wild-type seedlings (Figure 1C). The *rol5-1* mutation was found to be recessive, since the F1 generation of a backcross with *lrx1* developed an *lrx1* phenotype and seedlings



**Figure 1.** Suppression of the *lrx1* Root Hair Phenotype by Mutations in *rol5*.

Seedlings were grown for 5 d ([A] to [E]) and 10 d (F) in a vertical orientation. The wild type (A) developed regular root hairs, whereas root hairs of the *lrx1* mutant (B) were severely deformed. The EMS missense allele *rol5-1* (C) was complemented with a *ROL5* genomic clone, inducing an *lrx1*-like phenotype (D). The *rol5-2* T-DNA knockout mutant (E) also suppressed the *lrx1* mutant phenotype. The *rol5* mutation (F) leads to shorter roots as shown for *rol5-1*. Bars = 0.5 mm in (A) to (E) and 10 mm in (F).



**Figure 2.** Identification of the *ROL5* Locus.

**(A)** The *rol5* locus was identified by map-based cloning on the long arm of chromosome 2, south of *AthBio2*. BAC clones in the region of *ROL5* are indicated. For mapping, cleaved-amplified polymorphic sequence

of the F2 generation segregated 3:1 for *lrx1*:wild-type-like root hairs. To characterize the effect of the *rol5-1* mutation in more detail, a *rol5-1* single mutant was identified after backcrossing with wild-type Columbia (see Methods).

#### Map-Based Cloning of *rol5*

The *rol5* gene was identified by map-based cloning and initially localized to a region on chromosome 2, south of *AthBio2*. Further mapping revealed two flanking markers on the BAC *F411*, *F411-SphI*, and *F411-ClaI* at positions 16,700 and 43,500, respectively (Figure 2A). This interval was sequenced and a single point mutation was identified in the gene At2g44270 encoding a protein of 355 amino acids (Figure 2B). Transformation of the *lrx1 rol5-1* mutant with a wild-type genomic copy of this gene led to the development of an *lrx1* root hair phenotype, confirming that the identified gene represents the *ROL5* locus (Figure 1D). The mutation in *rol5-1* results in an amino acid change from Gly-65 to Asp. A *rol5* T-DNA mutant with the insertion site 3' adjacent to the Glu-170 codon was identified and named *rol5-2* (Figure 2B). RT-PCR on RNA isolated from wild-type and mutant seedlings revealed the presence of *ROL5* RNA in wild-type and *rol5-1* mutant seedling root and shoot tissue but not in *rol5-2* seedlings (Figure 2C). Together with the position of the T-DNA in an exon, this suggests that *rol5-2* is a null allele. The *lrx1* phenotype was also suppressed by *rol5-2* (Figure 1E), revealing that suppression is not dependent on the particular missense mutation present in the *rol5-1* allele.

The RT-PCR data indicated that *ROL5* is expressed in various tissues. For a more detailed analysis, a *ROL5:ROL5-GFP* (green fluorescent protein) fusion construct was transformed into wild-type *Arabidopsis* and roots of transgenic seedlings were analyzed. Fluorescence was found to be predominant in the

and simple sequence length polymorphism markers were established, of which *F411-Sph* and *F411-Cla* were the closest flanking markers identified.

**(B)** The *ROL5* gene consists of 10 exons encoding a protein of 355 amino acids. The G-to-A mutation in *rol5-1* is located in the second exon and changes Gly-65 to Asp. *rol5-2* represents a T-DNA insertion line that interrupts the reading frame at the amino acid codon Glu-170. Gray boxes, exons.

**(C)** RT-PCR experiments on RNA isolated from shoots (S) and roots (R) of 1-week-old seedlings demonstrated that the *ROL5* gene is expressed in the wild type and the *rol5-1* mutant but not to detectable levels in *rol5-2*. RT-PCR on the *ACTIN2* gene was performed to confirm the use of similar amounts of RNA in the different samples. One of two biological replicates is shown.

**(D)** In roots, *ROL5* is predominantly expressed in the elongation zone (ez) and in a striped pattern in the differentiation zone (dz) (top panel). A close-up of the root (GFP fluorescence in the middle panel; bright field in the bottom panel) revealed overlapping GFP fluorescence and root hair formation. Red dots, root hair-forming trichoblasts; arrow, root hair structure. Bar = 0.3 mm.

**(E)** When transiently expressed in *Arabidopsis* epidermal cells, *ROL5-GFP* (left panel) and a mitochondrial marker protein (for details, see Methods) fused to red fluorescent protein (middle panel) display overlapping fluorescence patterns (right panel). Bar = 50  $\mu$ m.

**Table 1.** Length of Roots, Trichoblasts, and Root Hairs of the *rol5-1* Mutant

Genotype	Root Length (mm)	Epidermal Cell Length (Trichoblasts) (μm)	Root Hair Length (μm)
Wild Type	15 ± 0.2	147 ± 29	700 ± 80
<i>rol5-1</i>	10 ± 0.2	126 ± 24	480 ± 120

Seedlings were grown for 5 d in a vertical orientation. Values represent means ± SD. Differences are significant (*t* test, *P* < 0.05).

elongation zone and to expand in a striped pattern into the differentiation zone. These stripes overlapped with the arrangement of root hair cells (Figure 2D), which initiate root hair elongation in the differentiation zone (Dolan et al., 1994). Thus, *ROL5* is predominantly expressed in elongating cells, suggesting an important function during cell expansion. Indeed, compared with wild-type seedlings, *rol5-1* mutants had shorter roots, root epidermal cells, and root hairs (Figure 1F, Table 1).


**ROL5 Is Structurally and Functionally Similar to the Yeast Ncs6p**

*ROL5* shows 54% identity and 70% similarity to Ncs6p/Tuc1p of yeast (*Saccharomyces cerevisiae*), subsequently referred to as Ncs6p (Figure 3). The Ncs6p-like proteins of different organisms share conserved motifs, including a PP-loop domain with ATP pyrophosphatase activity (Bork and Koonin, 1994; Björk et al., 2007), which are also conserved in *ROL5*. The Gly-65 to Asp mutation in *rol5-1* is adjacent to the PP-loop motif SGGxDS

(Figure 3). Ncs6p-like proteins have been found to be involved in the thiolation of the uridine residue 34 of a subset of cytoplasmic tRNAs (Björk et al., 2007; Schlieker et al., 2008; Leidel et al., 2009). To investigate whether *ROL5* is involved in tRNA modification in *Arabidopsis*, the tRNA fraction of wild-type and *rol5-1* mutant seedlings was isolated. tRNAs containing this modification can be detected in an acrylamide gel containing *N*-acryloylamino phenyl mercuric chloride (APM), a compound that interacts with 2-thiouridine and retards migration in the gel. While a band shift can be observed with wild-type tRNAs in gels containing APM, it is absent from *rol5-1* tRNA extracts. In the absence of APM, as expected, no shift is detectable in either of the extracts (Figure 4A). Hence, *ROL5* is involved in this tRNA modification process in *Arabidopsis*.


Since Ncs6p was also identified as a component of the TOR pathway, we assessed whether Ncs6p and *ROL5* have similar functions with respect to TOR signaling. The  $\Delta ncs6$  mutant yeast strain, which is hypersensitive to rapamycin (Chan et al., 2000; Goehring et al., 2003a), was complemented with *ROL5* under the control of a constitutive yeast promoter. On standard growth medium, the wild type, the  $\Delta ncs6$  mutant, and the complemented  $\Delta ncs6$  mutant showed comparable growth properties, while in the presence of rapamycin, growth of the  $\Delta ncs6$  mutant was considerably retarded. This effect was compensated for by expressing *ROL5* in the  $\Delta ncs6$  mutant, but not by expressing the *rol5-1* missense allele (Figure 4B).

Ncs6p appears to accumulate in mitochondria, which prompted us to investigate the subcellular localization of *ROL5*. A *ROL5*-GFP fusion construct was transiently expressed in *Arabidopsis* epidermal cells, and colocalization with well-established organellar



**Figure 3.** *ROL5* Is Homologous to Ncs6p of Yeast.

The alignment of *ROL5* with Ncs6p of *S. cerevisiae* reveals 54% identity and 70% similarity between the two proteins. The Ncs6p-like proteins of different organisms share common motifs that are indicated below the sequences [(CxxC)<sub>2</sub> - SGGxDS - CxxC - GH - PL - C - (CxxC)<sub>2</sub>], all of which are conserved between the two proteins. The motif PL is not fully conserved in Ncs6p and *ROL5*. Sequences important for protein activity (Björk et al., 2007) are boxed. The Gly-65 to Asp mutation in *rol5-1* (star) is adjacent to the PP-loop motif SGGxDS, which is important for ATP binding.



**Figure 4.** Ncs6p and ROL5 Have Similar Functions.

**(A)** tRNA was extracted from 7-d-old wild-type and *rol5-1* seedlings and separated on an acrylamide gel with (left panel) or without (right panel) APM, a compound that interacts with the 2-thiouridine and retards migration in the gel.

**(B)** Wild-type (WT) and  $\Delta ncs6$  mutant yeast was grown in the absence or presence of rapamycin. The  $\Delta ncs6$  mutant grew normally on control medium but was hypersensitive to rapamycin compared with the wild type. Expression of *ROL5* but not *rol5-1* under the control of the yeast *PHOSPHOGLYCERATE KINASE* promoter in the  $\Delta ncs6$  mutant suppressed this rapamycin hypersensitivity phenotype. Spots on a line represent serial dilutions (10-fold).

marker proteins was investigated. A clear overlap was found for ROL5-GFP and a mitochondrial protein (for details, see Methods) fused to red fluorescent protein (Figure 2E). This suggests that ROL5, similar to Ncs6p in yeast, translocates to mitochondria. Together, these data demonstrate that Ncs6p and ROL5 have very similar functions in their respective organisms.

### Interfering with TOR Signaling Leads to Suppression of *lrx1*


The functional similarity between ROL5 and Ncs6p suggested that the *rol5* mutant might be impaired in a TOR-related process. This led us to further investigate whether the *rol5* mutations suppress the *lrx1* root hair phenotype by influencing TOR signaling. To this end, the TOR-specific inhibitor rapamycin was used to interfere with TOR signaling in *Arabidopsis*. This required transformation of *Arabidopsis* with the yeast *FKBP12* under the control of the ubiquitously active 35S promoter. Wild-type Columbia plants expressing *FKBP12* were produced (Figure 5A), and the *lrx1* mutation was crossed into two independent transgenic lines. While *lrx1* mutants expressing *FKBP12* developed typical *lrx1* root hairs under normal growth conditions, the presence of rapamycin led to a clear suppression of the *lrx1* phenotype in both transgenic lines. In nontransgenic *lrx1* mutants, rapamycin had no effect on the root hair phenotype (Figure

5B). This shows that interfering with TOR signaling suppresses the *lrx1* phenotype.

The treatment with rapamycin had additional effects on root development that were similar to those observed in the *rol5-1* mutant. In the presence of rapamycin, *FKBP12*-expressing wild-type plants developed shorter roots and shorter epidermal cells (Table 2) as previously observed by Sormani et al. (2007), confirming the involvement of TOR signaling in plant cell elongation.

### *rol5-1* and Rapamycin Treatment Lead to Changes in Cell Wall Components

A possible mechanism of suppression of the *lrx1* root hair phenotype might be through compensation of the cell wall defects in *lrx1* mutants by the introduction of additional changes to cell walls. To identify potential alterations in cell wall structures, root surfaces were analyzed in the wild type and *rol5-1* mutant using a series of monoclonal antibodies targeted to different cell wall polysaccharide components. These were antipectic homogalacturonan JIM5 and JIM7 (Knox et al., 1990), anti-(1 → 4)- $\beta$ -D-galactan LM5 (Jones et al., 1997), anti-(1 → 5)- $\alpha$ -L-arabinan LM6 (Willats et al., 1998), antixyloglucan LM15 (Marcus et al., 2008), and anti-AGP LM2 (Yates et al., 1996). Four of these epitopes were detected at equivalent levels on wild-type and *rol5-1* mutant root surfaces (see Supplemental Figure 1 online), whereas the LM5 galactan and the LM2 AGP epitopes displayed differential modulation in response to the mutation. Detection of the LM5 galactan epitope decreased and that of the LM2 AGP epitope increased at the root surface of the



**Figure 5.** Rapamycin Treatment Suppresses the *lrx1* Root Hair Phenotype.

**(A)** RNA gel blot of wild-type (WT) *Arabidopsis* transformed with a 35S promoter:*FKBP12* construct.

**(B)** *lrx1* mutants expressing *FKBP12* were sensitive to rapamycin (rapam.) and showed a suppressed *lrx1* root hair phenotype (right). In nontransgenic *lrx1* mutants, the root hair phenotype was not affected by rapamycin (left). Bar = 0.5 mm.

**Table 2.** Length of Roots and Trichoblasts Due to Rapamycin Treatment

Treatment	Root Length (mm)	Epidermal Cell Length (Trichoblasts) ( $\mu\text{m}$ )
35S:ScFKBP12		
– Rapamycin	15 $\pm$ 0.2	140 $\pm$ 12
+ Rapamycin	11 $\pm$ 0.2	90 $\pm$ 9

Seedlings were grown for 5 d in a vertical orientation. Values represent means  $\pm$  SD. Differences are significant (*t* test,  $P < 0.05$ ).

*rol5-1* mutant (Figure 6A). The same monoclonal antibodies were used to analyze root surfaces upon interfering with TOR signaling by rapamycin. Treatment of seedlings expressing *FKBP12* with rapamycin resulted in alterations in immunolabeling that were similar to those observed in the *rol5-1* mutant. The LM5 galactan epitope showed a marked decrease in occurrence, whereas the LM2 AGP epitope was increased in proximal parts of the root (Figure 6B) compared with nontreated seedlings expressing yeast *FKBP12*. The other antibodies showed equal labeling for both conditions (see Supplemental Figure 2A online). Immunodetection with LM5 and LM2 was identical between nontransgenic wild-type seedlings grown with or without rapamycin and *FKBP12*-expressing seedlings grown without rapamycin (see Supplemental Figure 2B online). Thus, the observed modulations of these two cell wall epitopes were specifically induced by rapamycin and only in those seedlings that were expected to be rapamycin sensitive, suggesting that they were the result of impaired TOR signaling.

#### *rol5-1* Mutants Are Affected in Their Response to ROS and ROS Scavengers

One possible crossing point of TOR signaling and ROL5 is the mitochondrial localization of ROL5, since the TOR pathway is a regulator of mitochondrial activity and hence the production of ROS. To investigate this possibility, ROS levels in roots of wild-type and *rol5-1* mutant seedlings were analyzed using different ROS-sensitive staining substrates. Under the growth conditions used, none of these stainings revealed a clear, reproducible change in ROS levels in *rol5-1* seedlings (see Supplemental Figure 3 online). Next, the effect of ROS on seedling growth was tested in liquid culture. This experiment revealed an increased susceptibility of *rol5-1* seedlings to hydrogen peroxide. While wild-type seedlings showed a similar development with or without 8 mM  $\text{H}_2\text{O}_2$ , the development of *rol5-1* seedlings was retarded in the presence of  $\text{H}_2\text{O}_2$ . By contrast, *rol5-1* seedlings were revealed to be more tolerant to ROS scavengers. While wild-type seedlings barely grew and failed to accumulate chlorophyll in the presence of 100  $\mu\text{M}$   $\text{CuCl}_2$ , *rol5-1* seedlings turned green and grew considerably better (Figure 7). This indicates that ROL5 is important for the sensing of, and the response to, ROS.


#### DISCUSSION

The work presented here suggests that the TOR pathway is a process that can lead to the specific modification of cell wall

components. The *rol5* locus was identified in a suppressor screen on the *lrx1* mutant, which is affected in cell wall formation in root hairs (Baumberger et al., 2001, 2003b). This screen was performed with the aim of identifying novel loci involved in cell wall formation, as suppressors can reveal a functional relationship between genetic loci (Huang and Sternberg, 1995). After the previous identification of *rol1*, which encodes RHAMNOSE SYNTHASE1 (Diet et al., 2006), *rol5* is the second identified suppressor of *lrx1* and also affects cell wall structure.

#### The TOR Pathway Is a Regulator of Cell Wall Development

ROL5 is homologous to the yeast Ncs6p, and these proteins have similar functions in their respective organisms. These functions include the modification of tRNAs and the effect on TOR signaling. It is likely that suppression of *lrx1* is induced via a modification of TOR signaling, as the *lrx1* mutant root hair phenotype can be suppressed by the TOR kinase inhibitor rapamycin. Rapamycin is a macrocyclic lactone originally identified in the bacterium *Streptomyces hygroscopicus* and one of the most specific kinase inhibitors known (Heitman et al., 1991; Huang et al., 2003), making additional effects of rapamycin very unlikely. The TOR pathway is a central regulator of eukaryotic growth processes (Wullschlegel et al., 2006), and previous work




**Figure 6.** Immunolabeling of Cell Walls of *rol5-1* and Rapamycin-Treated Wild-Type Seedlings.

Immunolabeling of 4-d-old roots with monoclonal antibodies (1  $\rightarrow$  4)- $\beta$ -D-galactan side chains of RG I (LM5) and glucuronic acid-containing side chains of arabinogalactan proteins (LM2).

(A) Compared with the wild type (WT), the *rol5-1* mutant root surface revealed reduced detection of the LM5 epitope and stronger detection of the LM2 epitope.

(B) Roots of *FKBP12*-expressing wild-type seedlings grown in the absence (left) and presence (right) of rapamycin. The presence of rapamycin led to a reduced labeling with LM5 and a stronger labeling in proximal parts with LM2.

Arrowheads, root apex. Bars = 0.3 mm.



**Figure 7.** Altered Response of the *rol5-1* Mutant to ROS and ROS Scavenger.

Seedlings were grown in liquid culture for 10 d. Under control conditions (A), growth of the wild type and *rol5-1* is comparable. *rol5-1* seedlings are hypersensitive to H<sub>2</sub>O<sub>2</sub> (8 mM), revealed by the reduced growth (B), and hyposensitive to the ROS scavenger CuCl<sub>2</sub> (100 μM), indicated by better growth and the development of green cotyledons (C). col, wild-type Columbia.

in *Arabidopsis* has revealed the importance of this signaling pathway, including the TOR kinase itself, for plant development (Menand et al., 2002; Bögre et al., 2003; Mahfouz et al., 2006). Our analysis shows that the TOR pathway is able to modify cell wall components, suggesting that it is part of the regulatory process that coordinates cell wall structure with cell growth and development. In fact, recent work in *C. albicans* suggests that the TOR pathway is involved in sensing cell wall integrity. The *rhb1* mutant, affected in a small G-protein of the RAS superfamily, was found to be hypersensitive to drugs interfering with cell wall formation and showed a modified cell wall integrity signaling previously identified in this organism. The *rhb1* mutation induces rapamycin hypersensitivity in *C. albicans*, indicating a function of RHB1 in TOR signaling (Tsao et al., 2009). Plants clearly have mechanisms to monitor, sense, and modify cell wall composition and structures. Proteins involved in this process have been found to be transmembrane or membrane associated, such as wall-associated kinases, the receptor kinase THESEUS, GPI-anchored proteins, or lectins (Kohorn et al., 2006; Hematy et al., 2007; Humphrey et al., 2007; Hematy and Höfte, 2008). These proteins are probably directly involved in the regulatory or sensing process, since they localize to cell surfaces. Considering

the importance in regulating cell growth, the TOR pathway might function as a relay system that integrates the signals of sensing mechanisms into cellular responses and developmental processes.

The core component of the TOR pathway is the Ser/Thr kinase protein TOR. Only yeast encodes two separate but functionally similar TOR proteins. In yeast and mammals, TOR forms two distinct multiprotein complexes, TORC1 and TORC2, of which only TORC1 is rapamycin sensitive. While TORC1 is involved in regulating translation, nutrient import, or stress responses, TORC2 influences the actin cytoskeleton (Loewith et al., 2002; Wullschleger et al., 2006). Also in *Arabidopsis*, the diverse functions of this pathway are likely to require the establishment of distinct multiprotein complexes with the TOR protein. The TOR-interacting proteins RICTOR and RAPTOR were identified in TORC1 and TORC2 (Wullschleger et al., 2006), and RAPTOR has been shown to undergo interactions with the *Arabidopsis* TOR protein. Mutations in RAPTOR affect plant development, corroborating the importance of this protein for the TOR pathway (Deprost et al., 2005; Mahfouz et al., 2006). It is likely that TOR-interacting proteins establish further protein-protein interactions as shown for other organisms (Wullschleger et al., 2006), leading to diverse signaling outputs. A better understanding of the signaling network is required to identify the mechanism by which the TOR pathway senses and influences cell wall structures.

#### Interfering with TOR Modifies Cell Walls and Plant Growth

Plant cell walls are complex, structurally variable organelles that underpin many aspects of cell and organ growth. Cell wall extension is the limiting factor in cell enlargement (Carpita and Gibeaut, 1993; Martin et al., 2001). Monoclonal antibodies are a useful tool to analyze cell walls as they can detect changes in the composition or the accessibility of cell wall structures. The LM5 galactan epitope that is a component of RGI has been specifically detected in the elongation zone at the *Arabidopsis* root surface and implicated in the onset of the acceleration of cell elongation (McCartney et al., 2003). The occurrence of the LM5 epitope has also been observed to correlate with modified mechanical properties of cell walls and to be reduced in different mutants that show reduced root epidermal cell growth (McCartney et al., 2000, 2003; Diet et al., 2006). Thus, the short root phenotype observed in the *rol5-1* mutant and in rapamycin-treated seedlings correlates well with reduced LM5 labeling and the known involvement of this cell wall component with growth. Pectin regulates the porosity of cell walls and hence influences the mobility of cell wall modifying enzymes necessary for cell wall expansion (Baron-Epel et al., 1988). RGI is thought to modify this porosity, which would serve as a possible explanation of why changes in the RGI structure influence cell growth (Ridley et al., 2001; Willats et al., 2001). AGPs are a second cell wall component found to be altered due to modified TOR signaling. The LM2 antibody, which binds to a glucuronic acid-containing epitope of AGPs (Yates et al., 1996), showed increased immunolabeling of *rol5-1* mutants and rapamycin-treated seedlings. The modified distribution and/or abundance of AGPs has been shown to correlate with aberrant cell growth in roots, root hairs, and pollen tubes as demonstrated by the analysis of *agp* mutants as well as the use of Yariv reagent

that precipitates AGPs and blocks their action (Willats and Knox, 1996; Ding and Zhu, 1997; van Hengel and Roberts, 2002; McCartney et al., 2003; Levitin et al., 2008). Reducing root epidermal cell expansion with Yariv reagent also modifies the occurrence of the LM5 epitope, indicating some linkage between RGI and AGPs (McCartney et al., 2003) that has also now been shown in the *rol5-1* mutant and rapamycin-treated seedlings. The reduced cell elongation phenotypes observed in the *rol5* mutants and upon rapamycin treatment therefore correlate with changes to specific cell wall components. Thus, the TOR pathway might be a regulatory mechanism to modulate these two factors in cell walls. It is possible that the observed changes in root surface detection of the LM5 galactan and LM2 AGP epitopes are mechanistically involved in the suppression of the *lrx1* root hair phenotype. As *lrx1* mutants develop aberrant cell walls, it can be hypothesized that secondary modifications overcome the defects induced by the lack of LRX1.

### ROL5 Might Have Dual Functions

It remains unclear exactly how ROL5 affects TOR signaling. The *rol5-1* mutant, similar to the yeast  $\Delta ncs6$  mutant, fails to properly modify tRNAs. The uridine residue 34 of several tRNAs is modified to 5-methoxycarbonylmethyl-2-thiouridine to improve translational efficiency. Ncs6p, together with other proteins, has been shown to transfer the sulfur group during 2-thiouridine formation (Björk et al., 2007; Schlieker et al., 2008; Leidel et al., 2009; Noma et al., 2009). The TOR pathway is involved in regulating the translational machinery in different organisms, including plants (Mahfouz et al., 2006; Wullschleger et al., 2006; Dinkova et al., 2007), and the lack of tRNA modifications might trigger signals that feed back into TOR signaling. This indirect effect on the TOR pathway is a possible explanation for the rapamycin hypersensitivity of the  $\Delta ncs6$  mutant (Chan et al., 2000; Goehring et al., 2003a). Alternatively, Ncs6p and ROL5 might have an additional, so far unidentified function that links protein activity to the TOR signaling network. Indicative for this hypothesis is the localization of these proteins to mitochondria (Huh et al., 2003; this work). Previous work has shown that Ncs6p is dispensable for the thiolation of mitochondrial tRNA (Noma et al., 2009), suggesting an additional function of the protein in this organelle. The dual localization in different compartments has been shown for a number of proteins (Krause and Krupinska, 2009). Goehring et al. (2003a) reported on the influence of Ncs6p on protein conjugation by Urm1p, a ubiquitin-related modifier protein. Yeast Urm1p has recently been shown to be involved in the same sulfur carrier process as Ncs6p (Leidel et al., 2009). In addition, however, Urm1p is also conjugated to Ahp1p, which is not involved in tRNA modification but is likely to have a function in TOR signaling (Goehring et al., 2003b). Our analysis points to an additional effect of ROL5 in a ROS-related process as the *rol5-1* mutant showed an increased sensitivity to ROS and an increased tolerance to ROS scavengers compared with the wild type. The mitochondrial localization of ROL5 is consistent with this additional function since mitochondria are a major source of ROS (Rhoads et al., 2006). ROS are not just byproducts of the respiratory chain but revealed to serve as signaling molecules that can affect cell elongation and cell wall development (Liszskay

et al., 2004; Takeda et al., 2008). ROL5 might regulate the response of the cell to ROS signaling, which is in agreement with the reduced cell growth observed in *rol5-1* mutants. A major function of the TOR pathway is the regulation of mitochondrial activity (Schieke et al., 2006; Cunningham et al., 2007), and ROL5 might be part of this regulatory mechanism.

The TOR pathway is a central regulator of eukaryotic cell growth. The analyses presented here suggest that the TOR pathway has the ability to modify cell wall structure and specifically components implicated in cell elongation. The TOR pathway appears to be one mechanism of connecting plant cell growth processes with specific changes to cell wall structure. Further analyses are necessary to identify the proteins that establish the link between the TOR signaling network and the extracellular proteins that sense and survey cell wall developmental processes. Moreover, the possible multiple activities of ROL5-like proteins need to be elucidated in greater detail to identify their precise roles during cell growth.

### METHODS

#### Plant Growth, EMS Mutagenesis, and Mapping

The *lrx1* missense allele and the EMS mutagenesis procedure are described by Diet et al. (2004). The *lrx1* mutant and all other *Arabidopsis thaliana* lines used are in the Columbia genetic background, except for the line used for mapping, which is Landsberg *erecta* (*Ler*). The *rol5-2* allele (line 709D04) was obtained from the GABI collection (Rosso et al., 2003). Phenotypic analysis was performed on *lrx1 rol5-1* and *rol5-1* mutant plants backcrossed twice with the *lrx1* mutant and wild-type Columbia, respectively. *lrx1 rol5-1* and *lrx1 rol5-2* double mutants and *rol5-1* single mutants were identified with molecular markers for the mutations (see below). For growth of plants in sterile conditions, seeds were surface sterilized with 1% sodium hypochlorite and 0.03% Triton X-100, stratified for 3 d at 4°C, and grown for 5 d on half-strength Murashige and Skoog (MS) medium containing 0.6% Phytagel (Sigma-Aldrich), 2% sucrose, and 100 mg/L myo-inositol, or, for liquid culture, in half-strength MS medium, 1% sucrose, and 100 mg/L myo-inositol, with a 16-h-light/8-h-dark cycle at 22°C. For crosses and propagation of the plants, seedlings were transferred to soil and grown in growth chambers with a 16-h-light/8-h-dark cycle at 22°C. Plant transformation was performed as described by Diet et al. (2006).

For mapping, the *lrx1 rol5-1* mutant was crossed with *Ler* and propagated to the F2 generation. Five hundred F2 seedlings displaying a wild-type root hair phenotype were selected and screened for homozygous *lrx1* mutant plants with a PCR-based marker (Diet et al., 2004). These plants were assumed to be homozygous mutants for *rol5-1* and were thus used for initial mapping. Once the approximate map position of *rol5* was identified, F2 plants displaying an *lrx1* mutant phenotype (i.e., being homozygous mutant *lrx1*) were selected, and those heterozygous Columbia/*Ler* in the region containing the *rol5* locus were propagated to the F3 generation. As expected, seedlings of the F3 population segregated 3:1 for *lrx1* versus wild-type root hairs. One thousand wild-type-like F3 seedlings were selected for detailed mapping of *rol5*. Mapping was performed using standard simple sequence length polymorphism and cleaved-amplified polymorphic sequence markers developed based on the Columbia/*Ler* polymorphism databank (Jander et al., 2002).

#### Molecular Markers for Genotyping

The marker for *lrx1* was previously described (Diet et al., 2004). The *rol5-1* mutation was detected by PCR with the primers *rol5BanI\_F*

(5'-ACAATCTTAAGAGGCCAAACC-3') and *rol5BanI\_R* (5'-CATATTAAG-CAGAAGCTTGG-3'), followed by digestion with the enzyme *BanI*, which only cuts wild-type *ROL5* but not the *rol5-1* DNA. The T-DNA insertion in *rol5-2* is 3' adjacent to the sequence 5'-GTTATTGAAAGTAGAGA-3'. Homozygous *rol5-2* mutants were identified by DNA gel blotting using genomic DNA digested with *BglII* and a fragment of the *ROL5* gene 3' adjacent of the T-DNA insertion site as a specific probe for hybridization.

### DNA Constructs

For complementation of the *rol5-1* mutant, a *ROL5* genomic clone including 1.8 kb of the promoter region and 400 bp of terminator sequence was amplified by PCR using the primer pair *Rol5R1Not* (5'-ATTGCGCC-GCTGGGCTGGTGATGAAAGTTG-3'), *Rol5F1NotI* (5'-ATTGCGCCGC-CAGAGTGTCTTGATTGGTTTCG-3'). The PCR product was digested by *NotI* and cloned into the *pART27* plant transformation vector (Gleave, 1992) cut with the same enzyme. For the *ROL5*-GFP fusion constructs, the genomic clone of *ROL5* that was used for complementation was subjected to site-directed mutagenesis (QuikChange; Stratagene) to introduce one *BamHI* site using the primer pair *mutBamHI-midF* (5'-GAATCTCCTCCT-CGGATCCAAAACCTCATAAAAGC-3') and *mutBamHI-midR* (5'-GCTTT-TATGAGTTTTGGATCCGAGGAGGAGATTC-3'). The *GFP* gene was amplified from the vector *pMDC83* (Curtis and Grossniklaus, 2003) using the primer pair *GFP-F* (5'-TATGGATCCATGAGTAAAGGAGAAGAAGT-TTTC-3'), *GFP-R* (5'-AATGGATCCGT-GGTGGTGGTGGTGGTGGTGGTGGT-3') and cloned into the *BamHI* site of the *ROL5* gene. The resulting *ROL5*-*GFP* construct was cloned into the binary vector *pART27* with the restriction enzyme *NotI*. For transient expression in *Arabidopsis* epidermal cells, the *ROL5*-*GFP* construct was ligated into the overexpression cassette of *pART7* (Gleave, 1992) and used for particle bombardment. *CoxIV*-DsRed with a yeast *COXIV* presequence tag for mitochondrial localization (Mollier et al., 2002) was used as the mitochondrial marker protein. The overexpression construct for *ScFKBP12* was obtained by PCR amplification of *FKBP12* from yeast with the primer pair *ScFKBP12\_F* (5'-GAATTCATGTCTGAAGTAATTGAAGGTAAC-3'), *ScFKBP12\_R* (5'-TCTAGATTAGTTGACCTTCAACAATTCGAC-3') and cloning of the PCR product into *pART7* containing a *35S cauliflower mosaic virus promoter:ocs terminator* cassette (Gleave, 1992) by digestion with *EcoRI* and *XbaI*. A correct *pART7*-*ScFKBP12* clone was digested with *NotI*, and the excised *35S:ScFKBP12:ocs* cassette was inserted into the binary vector *pART27* digested with *NotI*. For expression in yeast, the coding sequence of a *ROL5* cDNA was amplified with the primers *ROL5-pFL61\_F* (5'-GCGGCCGCATGGAGGCCAAGAACAAGAAAGC-3') and *ROL5-pFL61\_R* (5'-GCGGCCGCCTTAGAAATCCAGAGATCCACATTG-3') and cloned into the expression vector *pFL61* (Minet et al., 1992) by digestion with *NotI*.

### Yeast Strain and Growth Conditions

Yeast strains used in this study were obtained from EUROSCARF, Frankfurt, Germany. The wild-type strain is BY4741 with the relevant genotype MATa; his3Δ1; leu2Δ0; met15Δ0; ura3Δ0, and the ncs6Δ strain has the relevant genotype MATa; his3Δ1; leu2Δ0; met15Δ0; ura3Δ0; YGL211w::kanMX4. Yeast strains were grown using standard methods. Synthetic yeast media was prepared with 2.4 nM rapamycin where indicated. Yeast strains were grown at 30°C until log phase and drops of an OD<sub>600</sub> of 0.8 and three subsequent 10-fold dilutions were spotted onto synthetic solid medium and grown for 3 d at 30°C.

### Transient Gene Expression in *Arabidopsis* Epidermal Cells

For transient gene expression, *Arabidopsis* leaf epidermal cells were transformed by particle bombardment as described (Escobar-Restrepo

et al., 2007). Bombarded tissue was incubated for 2 d at room temperature and the fluorescence analyzed using confocal microscopy.

### ROS Staining

For ROS staining, 0.1 mg/mL NBT was directly dissolved in 0.1 M K-phosphate buffer, pH 7, stirred for 60 min at room temperature, and filtered through a 0.2-μm pore size filter. Seedlings were incubated at room temperature for 60 min. The reaction was stopped by washing twice with 100% ethanol.

### Microscopy

GFP fluorescence was analyzed by confocal microscopy (DMIRE2; Leica) and analysis of immunolabeling on a LM510 (Zeiss). Phenotypic observations were performed with a Leica LZ M125 stereomicroscope. For cell and root hair length measurements, pictures were taken by differential interference contrast microscopy using an Axioplan microscope (Zeiss). Over 30 data points from ≥5 seedlings were collected. Root length was manually determined, using ≥20 seedlings per data point. The *t* test was used for statistical analysis, and the values are given with ± SD, *P* = 0.05. Confocal microscopy was performed on a DMIRE2 (Leica).

### Immunolabeling

Immunolabeling of surfaces of intact *Arabidopsis* seedling roots was performed using six rat monoclonal antibodies directed to cell wall components. *Arabidopsis* seedlings were prepared for immunofluorescence microscopy as described (McCartney et al., 2003). Seedlings were vertically grown for 4 d prior to immunolabeling. An FITC-linked anti-rat antibody (Sigma-Aldrich) was used as secondary antibody. Seedlings were mounted in a glycerol antifade solution (Citifluor AF1; Agar Scientific) for microscopy observation.

### RNA Extraction and RT-PCR

Seedlings were grown vertically on half-strength MS plates for 2 weeks. Approximately 150 seedlings of each plant line were cut at the hypocotyl to separate shoot and root tissue, and the tissues were used for extraction of total RNA using the TRIzol method (Gibco BRL). The reverse transcription was performed using the SuperScriptTM II RNase reverse transcriptase kit (Invitrogen). The resulting single-stranded DNA was used for PCR with 30 cycles. *ACTIN2* was amplified as a control using the primer pair *Actin2F* (5'-AATGAGCTTCGTATTGCTCC-3') and *Actin2R* (5'-GCACAGTGTGAGACACACC-3'). Levels for the *ROL5* transcript were checked using the primer pair *Rol5NorthF3* (5'-CCAAGATGTAACCTTT-CAAG-3') and *RolNorth2R* (5'-GCTTCTTTGTTTCTTATTATG-3').

### tRNA Extraction and Analysis

Whole seedlings grown for 14 d in half-strength liquid medium containing 1% sucrose and 100 mg/L myo-inositol were collected for tRNA extraction. Plant material was frozen in liquid nitrogen, ground, and stored at -80°C. Total RNA was extracted two times with acidic phenol and chloroform and one time with chloroform. tRNA was purified using Nucleobond AX 100 columns (Macherey-Nagel) and precipitated overnight at -20°C. One microgram of tRNA per sample was separated on an 8% polyacrylamide gel containing 7 M urea and 1 μg/μL APM chloride where indicated. For tRNA visualization, the gel was stained with SYBR Gold (Invitrogen).

### Accession Numbers

Sequence data from this article can be found in the Arabidopsis Genome Initiative or GenBank/EMBL databases under the following accession

numbers: *ROL5*, At2G44270; *LRX1*, At1G12040; *Sc FKBP12*, YNL023C; and *Sc NCS6*, YGL211W.

### Supplemental Data

The following materials are available in the online version of this article.

**Supplemental Figure 1.** Cell Wall Epitopes Not Affected by the *rol5-1* Mutation.

**Supplemental Figure 2.** Effect of Rapamycin on Cell Wall Epitopes.

**Supplemental Figure 3.** ROS Staining in Wild-Type and *rol5-1* Mutant Roots.

### ACKNOWLEDGMENTS

This work was supported by the Swiss National Science Foundation (R.-M.L., F.J., and C.R.) and Marie Curie WallNet MRTN-CT-2004512265 (Y.V.). We thank Sebastian Leidel for providing APM, Geneviève Ephritikhine for the vector encoding the mitochondrial marker protein, Markus Klein for the pFL61 yeast expression vector, and Robert Dudler for helpful discussions and critical reading of the manuscript. The service provided by The Arabidopsis Information Resource (www.arabidopsis.org) was crucial for this work.

Received November 21, 2009; revised May 12, 2010; accepted May 23, 2010; published June 8, 2010.

### REFERENCES

- Anderson, G.H., Veit, B., and Hanson, M.R. (2005). The Arabidopsis *AtRaptor* genes are essential for post-embryonic plant growth. *BMC Biol.* **3**: 12.
- Baron-Epel, O., Gharyal, P.K., and Schindler, M. (1988). Pectins as mediators of wall porosity in soybean cells. *Planta* **175**: 389–395.
- Baumberger, N., Doesseger, B., Guyot, R., Diet, A., Parsons, R.L., Clark, M.A., Simmons, M.P., Bedinger, P., Goff, S.A., Ringli, C., and Keller, B. (2003a). Whole-genome comparison of leucine-rich repeat extensins in Arabidopsis and rice: A conserved family of cell wall proteins form a vegetative and a reproductive clade. *Plant Physiol.* **131**: 1313–1326.
- Baumberger, N., Ringli, C., and Keller, B. (2001). The chimeric leucine-rich repeat/extensin cell wall protein LRX1 is required for root hair morphogenesis in *Arabidopsis thaliana*. *Genes Dev.* **15**: 1128–1139.
- Baumberger, N., Steiner, M., Ryser, U., Keller, B., and Ringli, C. (2003b). Synergistic interaction of the two paralogous Arabidopsis genes *LRX1* and *LRX2* in cell wall formation during root hair development. *Plant J.* **35**: 71–81.
- Björk, G.R., Huang, B., Persson, O.P., and Bystrom, A.S. (2007). A conserved modified wobble nucleoside (mcm(5)s(2)U) in lysyl-tRNA is required for viability in yeast. *RNA* **13**: 1245–1255.
- Bögre, L., Okresz, L., Henriques, R., and Anthony, R.G. (2003). Growth signalling pathways in Arabidopsis and the AGC protein kinases. *Trends Plant Sci.* **8**: 424–431.
- Bork, P., and Koonin, E.V. (1994). A p-loop-like motif in a widespread ATP pyrophosphatase domain: Implications for the evolution of sequence motifs and enzyme-activity. *Proteins* **20**: 347–355.
- Carpita, N.C., and Gibeaut, D.M. (1993). Structural models of primary cell walls in flowering plants: Consistency of molecular structure with the physical properties of the walls during growth. *Plant J.* **3**: 1–30.
- Chan, T.F., Carvalho, J., Riles, L., and Zheng, X.F.S. (2000). A chemical genomics approach toward understanding the global functions of the target of rapamycin protein (TOR). *Proc. Natl. Acad. Sci. USA* **97**: 13227–13232.
- Cunningham, J.T., Rodgers, J.T., Arlow, D.H., Vazquez, F., Mootha, V.K., and Puigserver, P. (2007). mTOR controls mitochondrial oxidative function through a YY1-PGC-1 alpha transcriptional complex. *Nature* **450**: 736–740.
- Curtis, M.D., and Grossniklaus, U. (2003). A gateway cloning vector set for high-throughput functional analysis of genes *in planta*. *Plant Physiol.* **133**: 462–469.
- Deprost, D., Truong, H.N., Robaglia, C., and Meyer, C. (2005). An Arabidopsis homolog of RAPTOR/KOG1 is essential for early embryo development. *Biochem. Biophys. Res. Commun.* **326**: 844–850.
- Deprost, D., Yao, L., Sormani, R., Moreau, M., Leterreux, G., Nicolai, M., Bedu, M., Robaglia, C., and Meyer, C. (2007). The Arabidopsis TOR kinase links plant growth, yield, stress resistance and mRNA translation. *EMBO Rep.* **8**: 864–870.
- Diet, A., Brunner, S., and Ringli, C. (2004). The *enl* mutants enhance the *lrx1* root hair mutant phenotype of *Arabidopsis thaliana*. *Plant Cell Physiol.* **45**: 734–741.
- Diet, A., Link, B., Seifert, G.J., Schellenberg, B., Wagner, U., Pauly, M., Reiter, W.D., and Ringli, C. (2006). The *Arabidopsis* root hair cell wall formation mutant *lrx1* is suppressed by mutations in the *RHM1* gene encoding a UDP-L-rhamnose synthase. *Plant Cell* **18**: 1630–1641.
- Ding, L., and Zhu, J.K. (1997). A role for arabinogalactan-proteins in root epidermal cell expansion. *Planta* **203**: 289–294.
- Dinkova, T.D., de la Cruz, H.R., Garcia-Flores, C., Aguilar, R., Jimenez-Garcia, L.F., and de Jimenez, E.S. (2007). Dissecting the TOR-S6K signal transduction pathway in maize seedlings: Relevance on cell growth regulation. *Physiol. Plant.* **130**: 1–10.
- Dolan, L., Duckett, C.M., Grierson, C., Linstead, P., Schneider, K., Lawson, E., Dean, C., Poethig, S., and Roberts, K. (1994). Clonal relationships and cell patterning in the root epidermis of Arabidopsis. *Development* **120**: 2465–2474.
- Escobar-Restrepo, J.M., Huck, N., Kessler, S., Gagliardini, V., Gheyselinck, J., Yang, W.C., and Grossniklaus, U. (2007). The FERONIA receptor-like kinase mediates male-female interactions during pollen tube reception. *Science* **317**: 656–660.
- Gapper, C., and Dolan, L. (2006). Control of plant development by reactive oxygen species. *Plant Physiol.* **141**: 341–345.
- Gleave, A.P. (1992). A versatile binary vector system with a T-DNA organizational structure conducive to efficient integration of cloned DNA into the plant genome. *Plant Mol. Biol.* **20**: 1203–1207.
- Goehring, A.S., Rivers, D.M., and Sprague, G.F. (2003a). Urmlylation: A ubiquitin-like pathway that functions during invasive growth and budding in yeast. *Mol. Biol. Cell* **14**: 4329–4341.
- Goehring, A.S., Rivers, D.M., and Sprague, G.F. (2003b). Attachment of the ubiquitin-related protein Urm1p to the antioxidant protein Ahp1p. *Eukaryot. Cell* **2**: 930–936.
- Heitman, J., Movva, N.R., and Hall, M.N. (1991). Targets for cell-cycle arrest by the immunosuppressant rapamycin in yeast. *Science* **253**: 905–909.
- Hematy, K., and Höfte, H. (2008). Novel receptor kinases involved in growth regulation. *Curr. Opin. Plant Biol.* **11**: 321–328.
- Hematy, K., Sado, P.E., Van Tuinen, A., Rochange, S., Desnos, T., Balzergue, S., Pelletier, S., Renou, J.P., and Höfte, H. (2007). A receptor-like kinase mediates the response of Arabidopsis cells to the inhibition of cellulose synthesis. *Curr. Biol.* **17**: 922–931.
- Huang, L.S., and Sternberg, P.W. (1995). Genetic dissection of developmental pathways. In *Methods in Cell Biology*, H.F. Epstein and C.C. Shaker, eds (San Diego, CA: Academic Press), pp. 97–122.

- Huang, S., Bjornsti, M.A., and Houghton, P.J. (2003). Rapamycins. *Cancer Biol. Ther.* **2**: 222–232.
- Huh, W.K., Falvo, J.V., Gerke, L.C., Carroll, A.S., Howson, R.W., Weissman, J.S., and O’Shea, E.K. (2003). Global analysis of protein localization in budding yeast. *Nature* **425**: 686–691.
- Humphrey, T.V., Bonetta, D.T., and Goring, D.R. (2007). Sentinels at the wall: Cell wall receptors and sensors. *New Phytol.* **176**: 7–21.
- Ingram, G.C., and Waites, R. (2006). Keeping it together: Co-ordinating plant growth. *Curr. Opin. Plant Biol.* **9**: 12–20.
- Jander, G., Norris, S.R., Rounsley, S.D., Bush, D.F., Levin, I.M., and Last, R.L. (2002). Arabidopsis map-based cloning in the post-genome era. *Plant Physiol.* **129**: 440–450.
- Jones, L., Seymour, G.B., and Knox, J.P. (1997). Localization of pectic galactan in tomato cell walls using a monoclonal antibody specific to (1→4)-beta-D-galactan. *Plant Physiol.* **113**: 1405–1412.
- Knox, J.P., Linstead, P.J., King, J., Cooper, C., and Roberts, K. (1990). Pectin esterification is spatially regulated both within cell walls and between developing tissues of root apices. *Planta* **181**: 512–521.
- Kohorn, B.D., Kobayashi, M., Johansen, S., Riese, J., Huang, L.F., Koch, K., Fu, S., Dotson, A., and Byers, N. (2006). An Arabidopsis cell wall-associated kinase required for invertase activity and cell growth. *Plant J.* **46**: 307–316.
- Krause, K., and Krupinska, K. (2009). Nuclear regulators with a second home in organelles. *Trends Plant Sci.* **14**: 194–199.
- Leidel, S., Pedrioli, P.G.A., Bucher, T., Brost, R., Costanzo, M., Schmidt, A., Aebersold, R., Boone, C., Hofmann, K., and Peter, M. (2009). Ubiquitin-related modifier Urm1 acts as a sulphur carrier in thiolation of eukaryotic transfer RNA. *Nature* **458**: 228–232.
- Levitin, B., Richter, D., Markovich, I., and Zik, M. (2008). Arabinogalactan proteins 6 and 11 are required for stamen and pollen function in Arabidopsis. *Plant J.* **56**: 351–363.
- Liszak, A., van der Zalm, E., and Schopfer, P. (2004). Production of reactive oxygen intermediates ( $O_2^-$ ,  $H_2O_2$ , and  $\cdot OH$ ) by maize roots and their role in wall loosening and elongation growth. *Plant Physiol.* **136**: 3114–3123.
- Loewith, R., Jacinto, E., Wulschleger, S., Lorberg, A., Crespo, J.L., Bonenfant, D., Oppliger, W., Jenoe, P., and Hall, M.N. (2002). Two TOR complexes, only one of which is rapamycin sensitive, have distinct roles in cell growth control. *Mol. Cell* **10**: 457–468.
- Mahfouz, M.M., Kim, S., Delauney, A.J., and Verma, D.P.S. (2006). Arabidopsis TARGET OF RAPAMYCIN interacts with RAPTOR, which regulates the activity of S6 kinase in response to osmotic stress signals. *Plant Cell* **18**: 477–490.
- Majewska-Sawka, A., and Nothnagel, E.A. (2000). The multiple roles of arabinogalactan proteins in plant development. *Plant Physiol.* **122**: 3–9.
- Marcus, S.E., Verherbruggen, Y., Herve, C., Ordaz-Ortiz, J.J., Farkas, V., Pedersen, H.L., Willats, W.G.T., and Knox, J.P. (2008). Pectic homogalacturonan masks abundant sets of xyloglucan epitopes in plant cell walls. *BMC Plant Biol.* **8**: 60.
- Martin, C., Bhatt, K., and Baumann, K. (2001). Shaping in plant cells. *Curr. Opin. Plant Biol.* **4**: 540–549.
- McCartney, L., Ormerod, A.P., Gidley, M.J., and Knox, J.P. (2000). Temporal and spatial regulation of pectic (1→4)-beta-D-galactan in cell walls of developing pea cotyledons: Implications for mechanical properties. *Plant J.* **22**: 105–113.
- McCartney, L., Steele-King, C.G., Jordan, E., and Knox, J.P. (2003). Cell wall pectic (1 → 4)-beta-D-galactan marks the acceleration of cell elongation in the Arabidopsis seedling root meristem. *Plant J.* **33**: 447–454.
- Menand, B., Desnos, T., Nussaume, L., Berger, F., Bouchez, D., Meyer, C., and Robaglia, C. (2002). Expression and disruption of the Arabidopsis TOR (target of rapamycin) gene. *Proc. Natl. Acad. Sci. USA* **99**: 6422–6427.
- Minet, M., Dufour, M.E., and Lacroute, F. (1992). Complementation of *Saccharomyces cerevisiae* auxotrophic mutants by *Arabidopsis thaliana* cDNAs. *Plant J.* **2**: 417–422.
- Mollier, P., Hoffmann, B., Debast, C., and Small, I. (2002). The gene encoding *Arabidopsis thaliana* mitochondrial ribosomal protein S13 is a recent duplication of the gene encoding plastid S13. *Curr. Genet.* **40**: 405–409.
- Noma, A., Sakaguchi, Y., and Suzuki, T. (2009). Mechanistic characterization of the sulfur-relay system for eukaryotic 2-thiouridine biogenesis at tRNA wobble positions. *Nucleic Acids Res.* **37**: 1335–1352.
- Rhoads, D.M., Umbach, A.L., Subbaiah, C.C., and Siedow, J.N. (2006). Mitochondrial reactive oxygen species. Contribution to oxidative stress and interorganellar signaling. *Plant Physiol.* **141**: 357–366.
- Ridley, B.L., O’Neill, M.A., and Mohnen, D.A. (2001). Pectins: Structure, biosynthesis, and oligogalacturonide-related signaling. *Phytochemistry* **57**: 929–967.
- Ringli, C. (2005). The role of extracellular LRR-extensin (LRX) proteins in cell wall formation. *Plant Biosyst.* **139**: 32–35.
- Rosso, M.G., Li, Y., Strizhov, N., Reiss, B., Dekker, K., and Weisshaar, B. (2003). An *Arabidopsis thaliana* T-DNA mutagenized population (GABI-Kat) for flanking sequence tag-based reverse genetics. *Plant Mol. Biol.* **53**: 247–259.
- Schieke, S.M., and Finkel, T. (2006). Mitochondrial signaling, TOR, and life span. *Biol. Chem.* **387**: 1357–1361.
- Schieke, S.M., Phillips, D., McCoy, J.P., Aponte, A.M., Shen, R.F., Balaban, R.S., and Finkel, T. (2006). The mammalian target of rapamycin (mTOR) pathway regulates mitochondrial oxygen consumption and oxidative capacity. *J. Biol. Chem.* **281**: 27643–27652.
- Schlieker, C.D., Van der Veen, A.G., Damon, J.R., Spooner, E., and Ploegh, H.L. (2008). A functional proteomics approach links the ubiquitin-related modifier Urm1 to a tRNA modification pathway. *Proc. Natl. Acad. Sci. USA* **105**: 18255–18260.
- Sormani, R., Yao, L., Menand, B., Ennar, N., Lecampion, C., Meyer, C., and Robaglia, C. (2007). *Saccharomyces cerevisiae* FKBP12 binds *Arabidopsis thaliana* TOR and its expression in plants leads to rapamycin susceptibility. *BMC Plant Biol.* **7**: 26.
- Takeda, S., Gapper, C., Kaya, H., Bell, E., Kuchitsu, K., and Dolan, L. (2008). Local positive feedback regulation determines cell shape in root hair cells. *Science* **319**: 1241–1244.
- Tsao, C.C., Chen, Y.T., and Lan, C.Y. (2009). A small G protein Rhb1 and a GTPase-activating protein Tsc2 involved in nitrogen starvation-induced morphogenesis and cell wall integrity of *Candida albicans*. *Fungal Genet. Biol.* **46**: 126–136.
- van Hengel, A.J., and Roberts, K. (2002). Fucosylated arabinogalactan-proteins are required for full root cell elongation in Arabidopsis. *Plant J.* **32**: 105–113.
- Willats, W.G.T., and Knox, J.P. (1996). A role for arabinogalactan-proteins in plant cell expansion: Evidence from studies on the interaction of beta-glucosyl Yariv reagent with seedlings of *Arabidopsis thaliana*. *Plant J.* **9**: 919–925.
- Willats, W.G.T., Marcus, S.E., and Knox, J.P. (1998). Generation of a monoclonal antibody specific to (1→5)-alpha-L-arabinan. *Carbohydr. Res.* **308**: 149–152.
- Willats, W.G.T., McCartney, L., Mackie, W., and Knox, J.P. (2001). Pectin: Cell biology and prospects for functional analysis. *Plant Mol. Biol.* **47**: 9–27.
- Wulschleger, S., Loewith, R., and Hall, M.N. (2006). TOR signaling in growth and metabolism. *Cell* **127**: 5–19.
- Yates, E.A., Valdor, J.F., Haslam, S.M., Morris, H.R., Dell, A., Mackie, W., and Knox, J.P. (1996). Characterization of carbohydrate structural features recognized by anti-arabinogalactan-protein monoclonal antibodies. *Glycobiology* **6**: 131–139.

## The TOR Pathway Modulates the Structure of Cell Walls in *Arabidopsis*

Ruth-Maria Leiber, Florian John, Yves Verhertbruggen, Anouck Diet, J. Paul Knox and Christoph Ringli  
*Plant Cell* 2010;22;1898-1908; originally published online June 8, 2010;  
DOI 10.1105/tpc.109.073007

This information is current as of June 17, 2019

<b>Supplemental Data</b>	<a href="/content/suppl/2010/05/26/tpc.109.073007.DC1.html">/content/suppl/2010/05/26/tpc.109.073007.DC1.html</a>
<b>References</b>	This article cites 66 articles, 22 of which can be accessed free at: <a href="/content/22/6/1898.full.html#ref-list-1">/content/22/6/1898.full.html#ref-list-1</a>
<b>Permissions</b>	<a href="https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&amp;issn=1532298X&amp;WT.mc_id=pd_hw1532298X">https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&amp;issn=1532298X&amp;WT.mc_id=pd_hw1532298X</a>
<b>eTOCs</b>	Sign up for eTOCs at: <a href="http://www.plantcell.org/cgi/alerts/ctmain">http://www.plantcell.org/cgi/alerts/ctmain</a>
<b>CiteTrack Alerts</b>	Sign up for CiteTrack Alerts at: <a href="http://www.plantcell.org/cgi/alerts/ctmain">http://www.plantcell.org/cgi/alerts/ctmain</a>
<b>Subscription Information</b>	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: <a href="http://www.aspb.org/publications/subscriptions.cfm">http://www.aspb.org/publications/subscriptions.cfm</a>