


IN BRIEF

Fine-Tuning Plant Growth in the Face of Drought ^{OPEN}

Limiting shoot growth is an important survival strategy for plants during times of drought; smaller leaves mean that less water is lost through transpiration and more is retained in the soil. As drought stress restricts both cell division and expansion—processes that rely on specialized microtubule arrays—factors that regulate microtubule organization and stability likely affect a plant’s ability to grow in dry conditions. However, the signaling pathways that link drought stress with changes in microtubule organization are unknown.

Now, **Bhaskara et al. (2017)** report a signaling module in *Arabidopsis thaliana* that influences microtubule organization and regulates plant growth, particularly during prolonged exposure to drought stress. Prompted by the finding that several Type 2C protein phosphatases (PP2Cs) function in drought signaling (Bhaskara et al., 2012; Cutler et al., 2010; Schweighofer et al., 2004), the authors identified three closely related PPC2s (EGRs) that inhibited plant growth and were dramatically induced by drought stress. Phosphoproteomic analysis revealed 17 putative EGR targets, many of which had established roles in cytoskeleton regulation or vesicle trafficking. The authors focused on one of these targets (which they named Microtubule-Associated Stress Protein 1 [MASP1]), due to its massive increase in abundance in drought-stressed plants.

In contrast to the EGR protein phosphatases, MASP1 was found to promote growth, both under control and drought stress conditions. The authors showed that MASP1 associated with EGR proteins at the cell periphery. Given the enrichment of cytoskeleton-related proteins among the putative EGR targets, the authors then used lines expressing GFP-tagged α -tubulin to examine if EGR and MASP1 regulate microtubule organization. In agreement with previous reports (Mei et al., 2012; Fujita et al., 2013), the authors found that cortical microtubule ar-



The EGR-MASP1 module regulates plant growth during times of drought. Model showing the inhibitory effect of EGR and the stimulatory effect of MASP1 on both microtubule (MT) stability and plant growth under drought stress conditions. (Adapted from Bhaskara et al. [2017], Figure 8.)

rays became disorganized within a few hours of exposure to drought stress and partially recovered over the next few days. Strikingly, plants that overexpressed MASP1 or lacked EGR phosphatases had more stable microtubule arrays and exhibited more rapid recovery of microtubule organization under drought stress conditions. Further analysis showed that MASP1 associates with microtubules in vitro. In planta, MASP1 phosphorylation at Ser-670 is critical for growth promotion and microtubule stabilization. Finally, epistasis tests indicated that EGRs and MASP1 function in a common pathway, with MASP1 acting downstream of EGRs.

Based on these findings, the authors developed a model in which EGR phosphatases and MASP1 regulate microtubule stability during drought stress, with the former reducing and the latter promoting microtubule stability and anisotropic cell expansion (see figure). Fine-tuning the activity of these two opposing sets of proteins may improve plant growth and yield during moderate drought.

REFERENCES

Bhaskara, G.B., Nguyen, T.T., and Verslues, P.E. (2012). Unique drought resistance functions of the highly ABA-induced clade A protein phosphatase 2Cs. *Plant Physiol.* **160**: 379–395.

Bhaskara, G.B., Wen, T.-N., Nguyen, T.T., and Verslues, P.E. (2017). Protein phosphatase 2Cs and *Microtubule-Associated Stress Protein 1* control microtubule stability, plant growth, and drought response. *Plant Cell* **29**: 169–191.

Cutler, S.R., Rodriguez, P.L., Finkelstein, R.R., and Abrams, S.R. (2010). Abscisic acid: emergence of a core signaling network. *Annu. Rev. Plant Biol.* **61**: 651–679.

Fujita, S., Pytela, J., Hotta, T., Kato, T., Hamada, T., Akamatsu, R., Ishida, Y., Kutsuna, N., Hasezawa, S., Nomura, Y., Nakagami, H., and Hashimoto, T. (2013). An atypical tubulin kinase mediates stress-induced microtubule depolymerization in *Arabidopsis*. *Curr. Biol.* **23**: 1969–1978.

Mei, Y., Gao, H.B., Yuan, M., and Xue, H.W. (2012). The *Arabidopsis* ARCP protein, CSI1, which is required for microtubule stability, is necessary for root and anther development. *Plant Cell* **24**: 1066–1080.

Schweighofer, A., Hirt, H., and Meskiene, L. (2004). Plant PP2C phosphatases: emerging functions in stress signaling. *Trends Plant Sci.* **9**: 236–243.

Kathleen L. Farquharson
Science Editor
kfarquharson@aspb.org
ORCID ID: 0000-0002-8032-0041

Fine-Tuning Plant Growth in the Face of Drought
Kathleen L. Farquharson
Plant Cell 2017;29;4; originally published online January 18, 2017;
DOI 10.1105/tpc.17.00038

This information is current as of September 28, 2020

Supplemental Data	/content/suppl/2017/01/27/tpc.17.00038.DC1.html
References	This article cites 6 articles, 3 of which can be accessed free at: /content/29/1/4.full.html#ref-list-1
Permissions	https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&issn=1532298X&WT.mc_id=pd_hw1532298X
eTOCs	Sign up for eTOCs at: http://www.plantcell.org/cgi/alerts/ctmain
CiteTrack Alerts	Sign up for CiteTrack Alerts at: http://www.plantcell.org/cgi/alerts/ctmain
Subscription Information	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: http://www.aspb.org/publications/subscriptions.cfm