

IN BRIEF


Secrets of the Forest: Volatiles First Discovered in Pine Trees Propagate Defense Signals within and between Plants ^{OPEN}

Systemic acquired resistance (SAR)—a plant-wide heightened state of defense following localized exposure to a pathogen—is characterized by increased salicylic acid (SA) and reactive oxygen species (ROS) levels and elevated expression of pathogenesis-related genes. SAR depends on ENHANCED DISEASE SUSCEPTIBILITY1 (EDS1), which both generates the SAR signal at the primary infection site and perceives it in distal, uninfected tissues (Breitenbach et al., 2014). Various vascular mobile signals function within SAR, including methyl salicylate (Park et al., 2007), the nonproteogenic amino acid pipecolic acid (Návarová et al., 2012), and the dicarboxylic acid azelaic acid (Jung et al., 2009).

In a new study, Riedlmeier et al. (2017) examined the role of volatile organic compounds (VOCs) in SAR. Using gas chromatography/mass spectrometry, they compared the emissions released by wild-type *Arabidopsis thaliana* plants and SAR-deficient *eds1-2* mutants several hours after induction of SAR. Whereas the wild-type plants released substantial amounts of the bicyclic monoterpenes α -pinene and β -pinene—volatiles best known for giving pine forests their enchanting scent—the mutants with compromised SAR released almost undetectable levels of these VOCs.

Based on these findings, the authors hypothesized that α -pinene and β -pinene induce SAR. To test this possibility, they exposed wild-type plants to a mixture of α -pinene and β -pinene or to the VOC solvent control for 3 d and then challenged the plants with the pathogenic bacterium *Pseudomonas syringae* pv tomato strain DC3000 (*Pst*). They found that bacterial growth was ~10-fold less in the leaves of plants that had been exposed to the monoterpenes and that the effect depended on pinene concentration. Bacterial growth was not limited in the *non-expressor of PR genes1-1* mutant, which has defects in signaling downstream of SA,

^{OPEN}Articles can be viewed without a subscription.
www.plantcell.org/cgi/doi/10.1105/tpc.17.00462



Model explaining the contribution of α -pinene and β -pinene to systemic immunity. Pinenes induce immunity via EDS1, SA biosynthesis (possibly via *CBP60g*), and NPR1 and upregulate the expression of AZI1. In addition, pinenes trigger the production of ROS, which in turn cause SAR-promoting AZA to accumulate. (Adapted from Riedlmeier et al. [2017], Figure 11.)

or in the *sa induction deficient2-1* (*sid2-1*) mutant, which has defects in SA biosynthesis, after 3 d of monoterpene exposure. Furthermore, ROS O_2^- accumulated in the leaves of plants treated with pinenes, but not in those of control plants. Microarray analysis showed that pinene treatment directly influenced key regulatory genes in SA biosynthesis and SAR (e.g., the gene encoding the lipid transfer protein AZELAIC ACID INDUCED1). Therefore, pinenes appear to induce SAR, in a manner that depends on SA signaling.

As monoterpenes are highly volatile, the authors directly tested whether these VOCs could function as mobile plant-to-plant defense signals. Wild-type receiver plants enclosed in a vacuum desiccator for 3 d with SAR-induced (i.e., monoterpene-emitting) sender plants were indeed more resistant to *Pst* infection than were receiver plants enclosed with control plants.

Thus, in addition to functioning in SAR signaling within the plant, α -pinene and β -pinene

act as airborne signals that prime neighboring plants for impending danger (see figure). This exciting finding suggests that the aroma of a pine forest has an important function.

Kathleen L. Farquharson
Science Editor
kfarquharson@aspb.org
ORCID ID: 0000-0002-8032-0041

REFERENCES

- Breitenbach, H.H., et al. (2014). Contrasting roles of the apoplastic aspartyl protease APOPLASTIC, ENHANCED DISEASE SUSCEPTIBILITY1-DEPENDENT1 and LEGUME LECTIN-LIKE PROTEIN1 in Arabidopsis systemic acquired resistance. *Plant Physiol.* **165**: 791–809.
- Jung, H.W., Tschaplinski, T.J., Wang, L., Glazebrook, J., and Greenberg, J.T.

- (2009). Priming in systemic plant immunity. *Science* **324**: 89–91.
- Návarová, H., Bernsdorff, F., Döring, A.C., and Zeier, J.** (2012). Pipecolic acid, an endogenous mediator of defense amplification and priming, is a critical regulator of inducible plant immunity. *Plant Cell* **24**: 5123–5141.
- Park, S.W., Kaimoyo, E., Kumar, D., Mosher, S., and Klessig, D.F.** (2007). Methyl salicylate is a critical mobile signal for plant systemic acquired resistance. *Science* **318**: 113–116.
- Riedlmeier, M., Ghirardo, A., Wenig, M., Knappe, C., Koch, K., Georgii, E., Dey, S., Parker, J.E., Schnitzler, J.-P., and Vlot, C.** (2017). Monoterpenes support systemic acquired resistance within and between plants. *Plant Cell* **29**: 1440–1459.

Secrets of the Forest: Volatiles First Discovered in Pine Trees Propagate Defense Signals within and between Plants

Kathleen L. Farquharson

Plant Cell 2017;29;1181-1182; originally published online June 13, 2017;

DOI 10.1105/tpc.17.00462

This information is current as of September 28, 2020

Supplemental Data	/content/suppl/2017/07/11/tpc.17.00462.DC1.html
References	This article cites 5 articles, 5 of which can be accessed free at: /content/29/6/1181.full.html#ref-list-1
Permissions	https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&issn=1532298X&WT.mc_id=pd_hw1532298X
eTOCs	Sign up for eTOCs at: http://www.plantcell.org/cgi/alerts/ctmain
CiteTrack Alerts	Sign up for CiteTrack Alerts at: http://www.plantcell.org/cgi/alerts/ctmain
Subscription Information	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: http://www.aspb.org/publications/subscriptions.cfm