




IN BRIEF

Nitrate Ahoy! Shoot Cytokinin Signals Integrate Growth Responses with Nitrogen Availability^[OPEN]

Nutrients are rarely distributed homogeneously in soil. Consequently, plants have local and long-distance signaling systems in place to monitor and coordinate both demand and supply of essential macronutrients such as nitrogen (N). The “N-demand” long-distance signal emanates from a section of the root system that may be facing N deprivation, while the “N-supply” signal is activated if N has been found in the vicinity of the root. This allows the plant to adapt its root architecture and ensure optimal N supply by inducing the expression of hundreds of genes and increasing the number of lateral roots to forage for required nutrients (Kiba and Krapp, 2016).

Interestingly, this long-distance signal relay occurs systemically via the shoot. For example, xylem-mobile C-terminally encoded peptides produced in N-starved roots are components of the N-demand signaling pathway (Tabata et al., 2014); upon perception by their cognate receptors in shoots, they can induce nitrate uptake transporters in distal roots. Similarly, once imported, nitrate initiates root-to-shoot signaling involving biosynthesis and movement of the plant growth hormone cytokinin (CK) from the site of production in the roots to the shoots where they control N assimilation. Therefore, in heterogenous nitrate environments, nitrate-inducible *IPT3* and *IPT5* genes can affect overall plant development by controlling biosynthesis of different active forms of CK including *trans*-zeatin (Kieber and Schaller, 2014).

New results from **Poitout et al. (2018)** further push CK to the center stage of the systemic control of nitrate uptake and metabolism. The authors used a hydroponic split-root experimental system wherein *Arabidopsis thaliana* roots



Model proposing the role of cytokinin in plant responses to heterogeneous nitrogen availability. NO_3^- provision leads to *trans*-zeatin (*tZ*) accumulation in roots, which is transported to the shoots and leads to differential control of root responses according to NO_3^- supply to the roots. The CK pathway is further integrated with the CEP-dependent pathway triggered in N-deprived roots. (Adapted from Poitout et al. [2018], Figure 6.)

were partitioned into two compartments. These were set up in three combinations: both compartments containing the same media with nitrate (1 mM KNO_3 ; N supplied homogeneously), without nitrate (1 mM KCl; N deprived homogeneously), or with each of these media in a different compartment (N supplied heterogeneously). Under these conditions, the authors compared the patterns of CK accumulation and associated transcriptional responses of the roots and shoots of wild-type plants to the CK biosynthetic and transport mutants *ipt3,5,7* and *abcg14*, respectively. Importantly, impaired cytokinin biosynthesis and transport did not affect local nitrate perception. However, unlike the wild type, both mutants failed to display a characteristic induction of “sentinel” genes such as *NITRITE REDUCTASE1* that respond rapidly to long-distance N-supply signals. Furthermore, transcriptomic analyses revealed that genes regulated by heterogeneous nitrate provision in shoots of wild-type plants were enriched for genes involved in glutamate synthesis, the amino acid into which N is first assimilated (Forde and Lea, 2007). By contrast, neither mutant showed this transcriptional reprogramming, indicating that N assimilation

pathways in shoots are CK dependent. In parallel, the authors measured the levels of all four isoprenoid CK forms and their derivatives to correlate the transcriptional changes with CK accumulation patterns in roots and shoots. Curiously, in both mutants, only the reduced levels of *trans*-zeatin in shoots and not any of the other isoprenoid CK forms correlated with the dampened shoot transcriptional response. The authors concluded that shoot *trans*-zeatin controls downstream growth responses and further nitrate uptake and assimilation in response to heterogeneous nitrate availability (see figure).

Findings described in this study challenge existing theories on long-distance control of nitrogen uptake and form the basis of future work that will likely determine how cytokinin signaling networks are integrated with CEP signals associated with N foraging.

^[OPEN]Articles can be viewed without a subscription.
www.plantcell.org/cgi/doi/10.1105/tpc.18.00453

REFERENCES

- Forde, B.G., and Lea, P.J.** (2007). Glutamate in plants: metabolism, regulation, and signalling. *J. Exp. Bot.* **58**: 2339–2358.
- Kiba, T., and Krapp, A.** (2016). Plant nitrogen acquisition under low availability: regulation of uptake and root architecture. *Plant Cell Physiol.* **57**: 707–714.
- Kieber, J.J., and Schaller, G.E.** (2014). Cytokinins. *The Arabidopsis Book* **12**: e0168, doi/10.1199/tab.0168.
- Poitout, A., Crabos, A., Petřík, I., Novák, O., Krouk, G., Lacombe, B., and Ruffel, S.** (2018). Responses to systemic nitrogen signaling in Arabidopsis roots involve *trans*-zeatin in shoots. *Plant Cell* **30**: 1243–1257.
- Tabata, R., Sumida, K., Yoshii, T., Ohyama, K., Shinohara, H., and Matsubayashi, Y.** (2014). Perception of root-derived peptides by shoot LRR-RKs mediates systemic N-demand signaling. *Science* **346**: 343–346.

Nitrate Ahoy! Shoot Cytokinin Signals Integrate Growth Responses with Nitrogen Availability

Sonali Roy

Plant Cell 2018;30:1169-1170; originally published online June 12, 2018;

DOI 10.1105/tpc.18.00453

This information is current as of April 10, 2021

References	This article cites 5 articles, 2 of which can be accessed free at: /content/30/6/1169.full.html#ref-list-1
Permissions	https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&issn=1532298X&WT.mc_id=pd_hw1532298X
eTOCs	Sign up for eTOCs at: http://www.plantcell.org/cgi/alerts/ctmain
CiteTrack Alerts	Sign up for CiteTrack Alerts at: http://www.plantcell.org/cgi/alerts/ctmain
Subscription Information	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: http://www.aspb.org/publications/subscriptions.cfm