


IN BRIEF

The ABCCs of Saffron Transportomics^[OPEN]

Saffron spice is prized for its aroma, flavor, and color. The color derives from highly concentrated apocarotenoid lycosides called crocins. Accumulation of specialized metabolites such as crocins often requires their sequestration in the vacuole to prevent cellular toxicity, feedback inhibition of biosynthetic enzymes, and further modification by cytoplasmic enzymes. Crocins can account for as much as 10% of the dry weight within stigmas of *Crocus sativus*, a feat that relies upon transporters embedded within the tonoplast membrane to facilitate selective uptake into the vacuole. However, plant genomes encode hundreds of transporters, complicating the identification of these vacuolar transporters. In this issue, **Demurtas et al. (2019)** implement a transportomics pipeline to successfully characterize crocin transporters.

Several characteristics were used to narrow the list of potential crocin transporters. Preference was given to ABCC- and MATE-type transporters, which have a propensity to localize to the tonoplast. Furthermore, candidates were required to have high transcriptional expression in the stigma and share coexpression networks with crocin metabolites or the first dedicated crocin biosynthetic enzyme, CCD2. ABCC2, ABCC4a, and MATE4 met all these criteria.

Transportomics is a sensitive, high-throughput methodology capable of simultaneously tracking multiple metabolites in complex mixtures at femtomolar concentrations (Krumppochova et al., 2012). To identify active crocin transporters and assess their specificity, microsomes (i.e., membrane fragments) isolated from yeast (*Saccharomyces cerevisiae*) expressing the candidate transporters were incubated in saffron



Transportomics Assay Workflow to Identify Crocin Transporters.

Schematic showing that candidate transporters from *C. sativus* are expressed heterologously in yeast, run through transport assays using yeast microsomes and saffron metabolite extracts, and analyzed by metabolomics to identify transporter activity. (Adapted from Demurtas et al. [2019], Figure 4A)

stigma hydroalcoholic extracts (see figure). Metabolomics was then used to identify which metabolites from the extract were imported into the microsome, to determine substrate specificity. These assays demonstrated that MATE4 is a flavonoid transporter, whereas ABCC2 and ABCC4a are bona fide ATP-dependent crocin transporters with varying stereoisomer preferences. ABCC4a transported both *cis*- and *trans*-crocin 3 in complex saffron stigma extracts but was able to transport only the *cis* version when individually purified metabolites were assayed. This unusual result suggests that a cooperativity mechanism is necessary for uptake of the *trans* form. Stereoselectivity might serve as an isomerization trapping mechanism (Matern et al., 1983) where acidic conditions within the vacuole favor the formation of the immobile *trans* form, thereby encouraging its accumulation by preventing crocin escape from the vacuole.

Notably, expression of ABCC4a together with CCD2 in *Nicotiana benthamiana* leaves stimulated higher crocin production than CCD2 expression alone—evidence that vacuolar crocin transporters can enhance net crocin accumulation. Importantly, the ability to produce

appreciable quantities of crocins in heterologous systems could mean higher supply and lower prices for saffron in the future. Collectively, this pioneering work lays a foundation for how transportomics can be harnessed to discover transporters of other important plant natural products.

Robert C. Augustine
 Department of Biology
 Vassar College
 raugustine@vassar.edu
 ORCID ID: 0000-0002-9614-1558

REFERENCES

- Demurtas, O.C., et al.** (2019). ABCC transporters mediate the vacuolar accumulation of crocins in saffron stigmas. *Plant Cell* **31**: 2789–2804.
- Krumppochova, P., Sapth, S., Brouwers, J.F., de Haas, M., de Vos, R., Borst, P., and van de Wetering, K.** (2012). Transportomics: Screening for substrates of ABC transporters in body fluids using vesicular transport assays. *FASEB J.* **26**: 738–747.
- Matern, U., Heller, W., and Himmelsbach, K.** (1983). Conformational changes of apigenin 7-O-(6-O-malonylglucoside), a vacuolar pigment from parsley, with solvent composition and proton concentration. *Eur. J. Biochem.* **133**: 439–448.

The ABCCs of Saffron Transportomics

Robert C. Augustine

Plant Cell 2019;31;2558; originally published online September 23, 2019;

DOI 10.1105/tpc.19.00720

This information is current as of April 10, 2021

References	This article cites 3 articles, 1 of which can be accessed free at: /content/31/11/2558.full.html#ref-list-1
Permissions	https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&issn=1532298X&WT.mc_id=pd_hw1532298X
eTOCs	Sign up for eTOCs at: http://www.plantcell.org/cgi/alerts/ctmain
CiteTrack Alerts	Sign up for CiteTrack Alerts at: http://www.plantcell.org/cgi/alerts/ctmain
Subscription Information	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: http://www.aspb.org/publications/subscriptions.cfm