



## CORRECTION


Bernoux, M., Timmers, T., Jauneau, A., Brière, C., de Wit, P.J.G.M., Marco, Y., Deslandes, L. (2008). RD19, an *Arabidopsis* cysteine protease required for RRS1-R-mediated resistance, is relocalized to the nucleus by the *Ralstonia solanacearum* PopP2 effector. *Plant Cell* 20: 2252–2264.

The authors of the above article request the correction of Supplemental Figure 6, as it has come to our attention that the bottom lanes labeled “WB:  $\alpha$ -HA” are not correct as indicated, but came from a different immunoblot. This was an inadvertent mistake, as the correct blots look very similar. We apologize for this mistake.

The original figure and a corrected version are shown below. This correction does not change the conclusion of the figure, as the purpose of this immunoblot was to verify the accumulation of intact RD-YFP fusion proteins (RD19, RDL1, and RDL2) in presence or absence of PopP2-3HA protein.


The revised figure corresponds to data obtained in 2008, and shows an anti-GFP immunoblot (top) showing the accumulation levels of RD19-YFPv, RDL1-YFPv, and RDL2-YFPv proteins expressed either alone or in the presence of PopP2-3HA, and an anti-PopP2 immunoblot (bottom) showing the accumulation levels of PopP2-3HA from the same protein extracts. Unfortunately, we no longer have the images of Ponceau staining corresponding to these immunoblots. Nevertheless, we believe that the non-specific signals visible on each of the two immunoblots allows us to conclude that similar amounts of protein were loaded in the different lanes, sufficient to make the stated conclusion.

*Note: This correction was reviewed by members of The Plant Cell editorial board. The authors are responsible for providing a complete listing and accurate explanations for all known errors or instances of inappropriate data handling or image manipulation associated with the original publication.*



**Supplemental Figure 6 (corrected).** Detection of Full Length YFPv-Tagged Proteins.

RD19-YFPv, RDL1-YFPv and RDL2-YFPv were transiently expressed alone or in the presence of PopP2-3HA in *N. benthamiana*. YFPv-tagged proteins and PopP2-3HA were detected using anti-GFP and anti-PopP2 antibodies, respectively. \*Similar amounts of total protein extracts were loaded as indicated by the intensity of non-specific bands detected on each immunoblot.



**Supplemental Figure 6 (original).** Detection of Full Length YFPv-Tagged Proteins.

RD19-YFPv (63.8 kD), RDL1-YFPv (68 kD) and RDL2-YFPv (69.6 kD) were transiently expressed alone or in the presence of PopP2-3HA in *N. benthamiana*. YFPv- and HA-tagged proteins were detected using anti-GFP and anti-HA antibodies, respectively. Equal amounts of total protein extracts were loaded as indicated by Ponceau staining.

## CORRECTION

*Plant Cell* 2020;32;3036-3037; originally published online July 2, 2020;  
DOI 10.1105/tpc.20.00482

This information is current as of April 18, 2021

<b>Permissions</b>	<a href="https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&amp;issn=1532298X&amp;WT.mc_id=pd_hw1532298X">https://www.copyright.com/ccc/openurl.do?sid=pd_hw1532298X&amp;issn=1532298X&amp;WT.mc_id=pd_hw1532298X</a>
<b>eTOCs</b>	Sign up for eTOCs at: <a href="http://www.plantcell.org/cgi/alerts/ctmain">http://www.plantcell.org/cgi/alerts/ctmain</a>
<b>CiteTrack Alerts</b>	Sign up for CiteTrack Alerts at: <a href="http://www.plantcell.org/cgi/alerts/ctmain">http://www.plantcell.org/cgi/alerts/ctmain</a>
<b>Subscription Information</b>	Subscription Information for <i>The Plant Cell</i> and <i>Plant Physiology</i> is available at: <a href="http://www.aspb.org/publications/subscriptions.cfm">http://www.aspb.org/publications/subscriptions.cfm</a>